# Texas School Survey of Substance Use Methodology Report 2010

# For the Texas Department of State Health Services

Prepared by
James A. Dyer, Ph.D.
Allison Seibert
Ashweeta Patnaik
Pratik Mhatre

Public Policy Research Institute
Texas A&M University
College Station, TX 77843-4476

# **Table of Contents**

INTRODUCTION	1
SURVEY INSTRUMENT	2
Survey Content	2
SURVEY SAMPLE	3
SELECTION OF DISTRICTS  SOLICITING DISTRICT PARTICIPATION  PARTICIPATION RATES	5 5
ALLOCATION OF SURVEYS AMONG DISTRICTS	6 6
SURVEY ADMINISTRATION PROCEDURES	8
DATA ENTRY AND ANALYSES	
Exaggerated ResponsesUnreported Grade Levels	
QUALITY CONTROL MEASURES	10
WEIGHTS, STANDARD ERRORS, AND CONFIDENCE INTERVALS	11
CONCLUSION	12
APPENDIX A - SURVEY INSTRUMENTS	A-1
APPENDIX B - PARTICIPATING DISTRICTS	B-1
APPENDIX C - CLASSROOM SELECTION GUIDELINES	C-1
APPENDIX D - SCHOOL SURVEY ADMINISTRATION	D-1
APPENDIX E - COMPUTATION OF SAMPLE WEIGHTS	
APPENDIX F - STANDARD ERROR COMPUTATION	
APPENDIX G - STANDARD ERROR & CONFIDENCE INTERVAL TABL	ES G-1

#### Introduction

The Public Policy Research Institute (PPRI), in conjunction with the Texas Department of State Health Services (DSHS) conducted the twelfth statewide survey of drug and alcohol use among Texas elementary and secondary students in the Spring of 2010. Originally implemented in 1988 as a component of a larger survey assessing substance use among the state's general population, the school survey has since become an ongoing, independent project. District surveys are offered every year with a statewide survey conducted every two years. The 2010 effort provides follow-up data reflecting changes over the past twenty two years in grades four through twelve.

The *Texas School Survey* project has two primary objectives. First, it serves to inform state and local policy-makers about the extent and nature of the substance use problem in Texas schools. Second, the statewide survey provides a standard of comparison for districts conducting local assessments of drug and alcohol use.

The purpose of this document is to describe the methodology used to administer the 2010 *Texas School Survey of Substance Use*. Following a brief introduction to the survey instrument itself, attention is then focused on sample selection and survey administration procedures. Next, methods for data processing and quality control are described and the report concludes with a review of standard error estimates.

## Survey Instrument

Two versions of the 2010 Texas School Survey of Substance Use were developed and administered. The first was a six-page questionnaire designed for students in grades seven through twelve. The second was a simplified three-page instrument created for students in grades four through six. The elementary survey differs from the secondary survey in that it has simplified language and some complex questions were omitted. Elementary students were asked about only four types of substances including tobacco (cigarettes, snuff, and chewing tobacco), alcohol (beer, wine, wine coolers, and liquor), inhalants, and marijuana. Secondary students were asked about the same substances, as well as a broader range of illicit drugs including powdered cocaine, crack, hallucinogens, uppers, downers, steroids, ecstasy, Rohypnol, DXM, and heroin. Other sets of questions on both the elementary and secondary instruments were designed to assess behavioral correlates of substance use and students' perceptions of support available to help them cope with substance-related problems.

The questionnaire was in a format that could be scanned optically, similar to that used for standardized testing. It was designed for anonymous self-administration by students with the aid of a staff member to pass out the survey, read a common set of instructions, monitor the class during survey administration, and collect the instruments after they are completed. The survey instruments are included in Appendix A.

## **Survey Content**

The *Texas School Survey of Substance Use* content was updated in 2010. Both the secondary and elementary instruments include a false drug designed to identify exaggerated responses. In previous versions of the surveys, this drug was called "Cosma" or "Shimeron". The current versions list this drug as "Oxaril".

Revisions were made to the elementary and secondary survey instruments in 2010. For the elementary survey, revisions included the addition of "magic markers" and "computer dusting sprays" as a category in the question related to lifetime inhalant use. For the secondary survey, revisions included: (1) removal of question related to where students had received information about alcohol and drugs out of school, (2) removal of question related to the purchase of Texas lottery tickets, and (3) addition of questions related to specific inhalant use. Revisions were made to improve response quality and to capture data related to new use trends. Measures were taken to ensure compatibility with previous survey data.

## Survey Sample

The sample of students for the 2010 survey was designed to be a stratified, weighted, random sample of all public school students between the fourth and twelfth grades in the state. In order to make administration practical, students were selected using a multi-stage stratified sampling procedure. This involved sampling districts, schools within districts, and classrooms within districts. All students in a sampled classroom were asked to participate in the survey.

#### **Selection of Districts**

The primary analytic cluster was the school district since the approval needed to administer the survey had to be obtained at that level. First, the districts were stratified according to how urban the counties were in which they were located. Stratification along a variable deemed to be highly correlated with the characteristics of interest is a means of increasing the precision of estimates in complex sampling designs. The strata were formed as follows. First, 28 border counties formed a separate border stratum. Of the remaining counties, the most urban stratum consisted of counties with metropolitan populations of 1,000,000 or more; the next stratum containing those between 250,000 and 1,000,000; and the third stratum was made up of those metropolitan areas with less than 250,000. The remainder of the state constituted a final major stratum.

The strata were further subdivided by relative size of the districts, so that each stratum had a combination of large and small districts. Due to their large size relative to other districts, a total of nine districts were sampled with a probability of one. This means that these districts are always selected as part of the sample. They formed two of the substrata. The strata are listed in Table 1.

Districts were selected for the state sample from the 11 strata, above, in the following manner. The 14 districts that formed strata 1-A and 2-A were included with probability one. From each of the remaining strata, excluding the border stratum, a simple random sample of districts was selected. The districts in the border stratum were also selected with probability one in that all districts constituting that stratum were invited to participate. The districts that were selected in this manner are listed in Table 2.

If a non-probability one district refused to participate, it was replaced with another district selected at random from within the stratum.

TABLE 1. Distribution of Selected Districts by Urban Class Size

Stratum	Group
1-A 1-B 1-C	Large Urban Counties- 55,000 < enrollment (probability one districts)  Large Urban Counties- 20,000 < enrollment < 55,000  Large Urban Counties- 10,000 < enrollment <20,000
1-D 2-A 2-B	Large Urban Counties- enrollment <10,000  Medium Urban Counties- 20,000 < enrollment (probability one districts)  Medium Urban Counties- enrollment <20,000
3-A 3-B	Small Urban Counties- 10,000 < enrollment Small Urban Counties- enrollment <10,000
4-A 4-B	Non-Urban Counties- 5,000 < enrollment Non-Urban Counties- enrollment < 5,000
5-A	Border (28 counties)

TABLE 2. State Sample by Strata

Original	State	Sample
(n	-237	)

Actual State Sample (n=78)

Stratum 1-A: District Name	Stratum 1-A: District Name
GARLAND ISD	ARLINGTON ISD
ARLINGTON ISD	FORT WORTH ISD
CYPRESS-FAIRBANKS ISD	NORTHSIDE ISD
ALDINE ISD	
NORTH EAST ISD	
KATY ISD	
FORT BEND ISD	
HOUSTON ISD	
FORT WORTH ISD	
NORTHSIDE ISD	
DALLAS ISD	

Stratum 1-B: District Name	Stratum 1-B: District Name
HUMBLE ISD	CARROLLTON-FARMERS BRANCH ISD
KELLER ISD	CONROE ISD
GALENA PARK ISD	GALENA PARK ISD
SAN ANTONIO ISD	KELLER ISD
CONROE ISD	LAMAR CISD
SPRING ISD	SAN ANTONIO ISD
MCKINNEY ISD	SPRING ISD
JUDSON ISD	
DENTON ISD	
CARROLLTON-FARMERS BRANCH ISD	
PASADENA ISD	
RICHARDSON ISD	

GOOSE CREEK CISD **IRVING ISD ALIEF ISD** MESQUITE ISD LAMAR CISD

## **Stratum 1-C: District Name**

## **Stratum 1-C: District Name**

**COMAL ISD** 

MAGNOLIA ISD

NORTHWEST ISD

HARLANDALE ISD

SCHERTZ-CIBOLO-U CITY ISD

**SOUTHWEST ISD** 

**ROCKWALL ISD** 

**CROWLEY ISD** 

**DEER PARK ISD** 

**EDGEWOOD ISD** 

**DUNCANVILLE ISD** 

**DEER PARK ISD** 

#### **Stratum 1-D: District Name**

#### **Stratum 1-D: District Name**

FT SAM HOUSTON ISD **COMMUNITY ISD** NAVARRO ISD **BLUE RIDGE ISD** LACKLAND ISD FORNEY ISD STAFFORD MSD

ALAMO HEIGHTS ISD

**EVERMAN ISD** 

**ALVARADO ISD** 

WALLER ISD

**GARNER ISD** 

PEASTER ISD

**AZLE ISD** 

KRUM ISD

**ROYSE CITY ISD** 

PILOT POINT ISD

**BROCK ISD** 

MONTGOMERY ISD

ALVARADO ISD **EVERMAN ISD KRUM ISD** PEASTER ISD **ROYSE CITY ISD** STAFFORD MSD

#### **Stratum 2-A: District Name**

#### **Stratum 2-A: District Name**

LEANDER ISD

PFLUGERVILLE ISD

**AUSTIN ISD** 

CORPUS CHRISTI ISD

ROUND ROCK ISD

LEANDER ISD

Stratum 2-B: District Name	Stratum 2-B: District Name
DRISCOLL ISD WEST OSO ISD MATHIS ISD GREGORY-PORTLAND ISD LUMBERTON ISD JARRELL ISD BRIDGE CITY ISD EANES ISD DEL VALLE ISD TAFT ISD LITTLE CYPRESS-MAURICEVILLE CISD LAGO VISTA ISD ARANSAS PASS ISD HUTTO ISD PORT ARANSAS ISD	DRISCOLL ISD LITTLE CYPRESS-MAURICEVILLE CISD LUMBERTON ISD
Stratum 3-A: District Name	Stratum 3-A: District Name
WICHITA FALLS ISD MIDLAND ISD ECTOR COUNTY ISD AMARILLO ISD KILLEEN ISD BRYAN ISD ALVIN ISD TYLER ISD BRAZOSPORT ISD	ALVIN ISD BRYAN ISD ECTOR COUNTY ISD
Stratum 3-B: District Name	Stratum 3-B: District Name
IDALOU ISD RED LICK ISD SHALLOWATER ISD WHITEHOUSE ISD COPPERAS COVE ISD KILGORE ISD FRIENDSWOOD ISD SLATON ISD WASKOM ISD BULLARD ISD MAUD ISD	FRIENDSWOOD ISD IDALOU ISD MAUD ISD SHALLOWATER ISD WHITEHOUSE ISD
Stratum 4-A: District Name	Stratum 4-A: District Name
CORSICANA ISD HUNTSVILLE ISD SAN ANGELO ISD	CORSICANA ISD GRANBURY ISD MIDWAY ISD

NACOGDOCHES ISD MIDWAY ISD GRANBURY ISD MOUNT PLEASANT ISD SHERMAN ISD PLAINVIEW ISD

MALONE ISD

MOUNT PLEASANT ISD NACOGDOCHES ISD

Stratum 4-B: District Name	Stratum 4-B: District Name	
THORNDALE ISD	BOOKER ISD	
LINDEN-KILDARE CISD	CELESTE ISD	
SILVERTON ISD	LANEVILLE ISD	
GLEN ROSE ISD	LINDEN-KILDARE CISD	
BOLING ISD	MARLIN ISD	
BLUFF DALE ISD	WESTBROOK ISD	
LANEVILLE ISD		
BARBERS HILL ISD		
WESTBROOK ISD		
CHESTER ISD		
WHITHARRAL ISD		
MULESHOE ISD		
HARTLEY ISD		
MARLIN ISD		
DIVIDE ISD		
CELESTE ISD		
CONNALLY ISD		
GROVETON ISD		
RICHLAND SPRINGS ISD		
GUSTINE ISD		
BOOKER ISD		
KINGSVILLE ISD		

Stratum 5-A: All ISDs in counties listed (n=107)		d Stratum 5-A: District Name	
BREWSTER	KINNEY	ALPINE ISD	MCALLEN ISD
BROOKS	LA SALLE	BALMORHEA ISD	MERCEDES ISD
CAMERON	MAVERICK	BENAVIDES ISD BROOKS COUNTY	OLTON ISD
CULBERSON	PECOS	ISD	PALMER ISD PECOS-BARSTOW-
DIMMIT	PRESIDIO	CANUTILLO ISD	TOYAH ISD PHARR-SAN JUAN-
DUVAL	REEVES	COTULLA ISD	ALAMO ISD
EDWARDS	STARR	DONNA ISD EDCOUCH-ELSA	POINT ISABEL ISD
EL PASO	TERRELL	ISD	RIO HONDO ISD

HIDALGO	UVALDE	FERRIS ISD	ROMA ISD
HUDSPETH	VAL VERDE	FT HANCOCK ISD	SAN BENITO CISD
JEFF DAVIS	WEBB	HIDALGO ISD	SAN ELIZARIO ISD
		IRAAN-	
JIM HOGG	WILLACY	SHEFFIELD ISD	SAN PERLITA ISD
JIM WELLS	ZAPATA	ITALY ISD	SANTA ROSA ISD
KENNEDY	ZAVALA	LA JOYA ISD	SHARYLAND ISD
		LITTLEFIELD ISD	SOUTH TEXAS ISD
		LOS FRESNOS	
		CISD	UNITED ISD
		LYFORD CISD	UTOPIA ISD
		MARFA ISD	VALLEY VIEW ISD
		MAYPEARL ISD	YSLETA ISD

## **Soliciting District Participation**

Obtaining cooperation from those districts that were randomly selected for the state sample when the selected district did not plan to do a local survey was sometimes a problem. Yet, it was critical to get data from as many of the originally selected districts as possible. Some state sample districts that were initially hesitant were persuaded to cooperate by the use of incentives. The various incentives used included waiving participation and sampling fees, offering to discount the fees for participating the following year, paying all shipping costs, and discounting campus level analyses fees.

In order to allow further analysis of substance use among students living on the Texas-Mexico border, school districts along the border were strongly encouraged to participate in the 2010 Texas School Survey. The survey was offered free of charge to border districts, and data was collected from a broadly defined 28-county area. The border sample was designed to collect data from approximately 500,000 students. One-hundred and eight school districts from 28 counties were invited to participate in the 2010 survey. Each border district surveyed was included in the state survey sample.

Seventy-eight of the original 237 selected districts participated in the study. Most districts declined due to the lack of time and resources involved in survey administration. The selected districts included 2.99 million elementary and secondary students and those that were included in the final sample included 914, 898 students. A total of 31 percent of the students in the original sample were in the final sampling frame.

Seventy-five of the seventy-eight districts sampled submitted both elementary and secondary surveys. Three districts -Maud, South Texas, and Conroe - did not administer surveys to elementary students.

## **Allocation of Surveys among Districts**

The state survey sample was designed to collect data from a minimum sample of about 5,555 students per grade, however, many districts chose to survey more than the minimum number of students specified in the state sampling plan. Some extremely small districts received somewhat more than a strict proportional allocation because, while the data was technically only needed from one or two students per grade, the survey was administered to the entire classroom. Similarly, in a few extremely large (urban) districts, fewer students were need for accuracy than would result from a true proportional allocation. All surveys submitted from a cooperating district were included in the sample. Accordingly, in the final analyses, the data were weighted to provide an accurate proportional allocation.

Had districts provided data for the minimum number of sample students, the sample would have included approximately 50,000 students; it actually included 57,328 elementary students and 96,271 secondary students (See Table 3). This significantly improves the accuracy of estimates.

TABLE 3. Number of Surveys Included in State Sample

	Total Non-Blank Surveys	Usable Surveys	Number rejected	Percent rejected
Secondary	101,595	96,271	5,324	5.24%
Elementary	58,338	57,328	1,010	1.73%
Total	159,933	153,599	6,334	3.96%

<sup>\*</sup>Surveys were rejected because the responses indicated exaggeration or the survey could not be matched to a sampled school and grade.

## **Allocation of Surveys among Classrooms and Campuses**

Once the number of surveys to be administered in each district was established, the next step was to determine the number of classrooms to be surveyed per grade. This was achieved by dividing the number of questionnaires per grade (ascertained for each district using proportional population calculations) by the average number of students per class---20 for grades four through six, 22 for grades seven through twelve. The result of this computation indicated the total number of classes to be surveyed. These classes were selected so that as many different campuses as possible were in the final sample. Ideally, the classrooms surveyed were evenly distributed across all campuses in the district. If there were more campuses containing a given grade than classrooms needed, then a simple random selection procedure was used to determine which campuses would be sampled. In general, once a campus was selected, all relevant grades at that campus were surveyed. Therefore, campus selection was not independent between grades.

TABLE 4. Survey Distribution by Grade

		•	
	Grade	Number of Usable Surveys	Percentage
Elementary	Grade 4	18,328	32%
	Grade 5	19,225	34%
	Grade 6	19,775	35%
	Total	57,328	100%
Secondary	Grade 7	18,097	19%
	Grade 8	15,939	17%
	Grade 9	18,304	19%
	Grade 10	15,777	16%
	Grade 11	14,893	16%
	Grade 12	13,261	14%
	Total	96,271	100%

#### **Selection of Classrooms within Campuses**

After the total number of classrooms to be surveyed in each grade at each campus was determined, it was necessary to identify specific classrooms. This selection procedure was performed by campus personnel based on a set of guidelines provided by PPRI (illustrated in Appendix C). Members of campus staff were asked to make a list by grade (according to

teacher's last name or some other convenient method) of all classes held during a selected class period. They were then instructed to use a random number table to select the exact classes to survey in each grade.

## **Other Sampling Considerations**

Some school districts sampled all students in all or some of the grades. In these districts, the methodology outlined above did not apply to the grades sampled at 100 percent.

## Survey Administration Procedures

Districts selected for inclusion in the state sample were notified about the project via letter and were sent a descriptive brochure. State sample districts that planned to administer a local drug and alcohol survey had virtually no procedural changes resulting from their involvement in the statewide project. In those districts that surveyed grades four through twelve, sufficient data was collected from all relevant campuses to meet the data collection needs of the statewide survey. These districts benefited from their inclusion in the state survey project because they were not charged for the surveys that became part of the state database. The larger number of surveys from these districts was weighted down so that their contribution to the final sample was in correct proportion.

In those instances where state sample districts were collecting local data for an incomplete combination of grades, or where they were not conducting local surveys at all, the campus and classroom selection procedures described above were applied. Arrangements for giving the survey were established on an individual basis with these districts. Since those not doing local surveys did not stand to gain directly from having the survey administered in their district, an effort was made to be as accommodating as possible. PPRI was able to arrange survey administration in the selected schools and classes by school personnel.

Starting with the 2010 Texas School Survey, each participating school district had the ability to administer either an online version of the survey via a secure website, or the traditional paperpencil administration. School districts selected which method of administration they preferred when signing up for the survey. Out of the seventy eight participating districts, only one school district (Bryan ISD) administered the online version of the survey.

Relevant personnel in the selected districts and campuses were provided with complete instructions and materials necessary to administer the survey (see Appendix D). Classrooms were selected randomly by PPRI based on information from a computer printout from the district or Campus Information Form. Teachers in selected classrooms were given a script to read so that all students would receive a standardized set of instructions. Teachers were also asked to complete a Classroom Identification Form that provided data on the number of students that should have taken the survey but were absent, and the number of students that were present but failed to complete the survey. This information was useful for computing error estimates.

Survey administration procedures for the online survey followed the same protocols as the paper-pencil administration, except for minor modifications. Instead of receiving paper copies of the survey instrument, students in sampled classrooms were provided with unique, randomly generated six digit alpha-numeric codes, referred to as survey tokens. These survey tokens provided access to the online survey via a secure website. Each survey token could only be utilized once. In addition, to ensure only sampled classrooms completed the online survey, all unused and used survey tokens were collected by the classroom teachers when surveys were completed. Lastly, PPRI limited access to the online survey during the district's scheduled survey administration period.

The envelopes containing either the survey tokens or survey instruments from all participating classrooms were collected and returned to PPRI.

## Data Entry and Analyses

As noted earlier, the format of the survey instruments enabled them to be scanned optically. Upon receipt at PPRI, the instruments were logged in, coded and scanned by staff or trained personnel.

## **Exaggerated Responses**

Because the *Texas School Survey* data is based entirely upon respondents' description of their own behavior, it is inevitable that some students will under- or over-report their use of drugs or alcohol, and to the extent possible PPRI attempted to identify and eliminate data from those respondents. Two checks were incorporated into the data analysis program to identify exaggerators. First, both elementary and secondary students were asked about their use of a false drug call "Oxaril". Data from students claiming to have used this substance were considered suspect and dropped from the analyses.

Second, checks were run to identify any students claiming extremely high levels of drug and alcohol use. Unbelievable high substance use for elementary students was defined as the use of five or more substances, 11 or more times in the past school year or over a lifetime. Secondary students were defined as exaggerators based on the following criteria: (1) students reported that they had five or more drinks of two or more beverages every day; (2) students reported that they had consumed three or more alcoholic beverages every day; or (3) students reported that they used four or more drugs (other than cigarettes, alcohol, or steroids) eleven or more times in the past month. As in those cases where students reported using "Oxaril", data from students reporting exaggerated use were also dropped from the analyses.

## **Unreported Grade Levels**

When students failed to report their grade level, it was impossible to determine unequivocally in which grade these students' data should be analyzed. When a grade level was missing, an estimate of the grade was made based on the students' age and the data were retained. Table 4 identifies the range of students' ages and the corresponding grade levels that were assigned. If both grade and age were missing, the data were dropped from the analyses.

TABLE 4. Age-Based Grade Assignments.

Age	Elementary Level	Age	Secondary Level
9	4 <sup>th</sup> Grade	12	7 <sup>th</sup> Grade
10	5 <sup>th</sup> Grade	13	8 <sup>th</sup> Grade
11	6 <sup>th</sup> Grade	14	9 <sup>th</sup> Grade
		15	10 <sup>th</sup> Grade
		16	11 <sup>th</sup> Grade
		17 or older	12 <sup>th</sup> Grade

## **Quality Control Measures**

To ensure the quality of the statewide survey data, a number of internal checks were put into place to guide survey processing. First, a quality control analyst oversaw the implementation of all pre- and post-analysis quality control procedures. As the following paragraphs describe, many aspects of PPRI's plan for quality control were embedded in automated procedures. However, there is no replacement for human oversight. The quality control analyst monitored and tracked the processing of each district's surveys from the initial mailing through the production of the final state report. Responsibilities included ensuring that surveys were properly coded and scanned and checking for anomalies in the final table of results.

In addition to the safeguards resulting from careful project oversight, there were also a number of procedural checks against error. For example, there was a possibility, however remote, that after the bindings of a set of survey instruments were cut, the instruments could be dropped or otherwise placed out of order. If this occurred, it is conceivable that some pages of data could have been read into the incorrect computer record. To resolve this problem, each instrument used in the 2010 survey was printed with a five-digit "litho-code" number. With this coding process, every page of a given instrument is printed with the same scannable number, but a unique number is assigned to every instrument. By using the litho-code, when each page of an instrument is scanned it will automatically be read into the correct computer record. In this way, even if the pages from different instruments were shuffled together and read randomly, all data derived from the same instrument would automatically be read to the same data record.

Litho-coding also enabled PPRI to confirm that data from every survey instrument read was associated with the correct district. Survey instruments were mailed to participating districts in consecutive order. By recording the beginning and ending instrument numbers going to each district, PPRI was able to check the litho-codes scanned for a given district. In this way, any stacks of data that could potentially have been inadvertently mislabeled could be easily identified.

Programming checks were also incorporated into the data analysis program by cross-analysis. That is, the same data was run in several different ways using existing programs, and program outputs were then compared for consistency. Confidence is high that these quality control features will ensure valid and reliable survey findings.

## Weights, Standard Errors, and Confidence Intervals

Weights were applied to each case based on the strata, district, and campus. The weights were applied so that the aggregation of students in each campus, district, and strata reflected their proportions in the actual district, campus, and strata populations. The formulae use to determine these weights are presented in Appendix E.

Standard errors and confidence intervals were estimated for each grade and the aggregation. The formulae used are presented in Appendix F. The table of standard errors and confidence intervals for past year and lifetime use of substances by grades are presented in Appendix G.

## **Conclusion**

The *Texas School Survey* has become a valuable policy tool for both state and local educators and policy-makers. The survey, performed every two years, provides timely and relevant information about current drug and alcohol use patterns among young people enrolled in Texas' public schools. Furthermore, longitudinal analysis can provide insight into changes in drug and alcohol prevalence over time. As was noted in the introduction, every state survey culminates in a Texas Department of State Health Services publication providing an overview of findings to date. Data is also available for independent analysis by policy-makers and academicians.

Appendix A Survey Instruments

## TEXAS SCHOOL SURVEY OF SUBSTANCE USE

## ELEMENTARY SURVEY INSTRUMENT 2010 VERSION

#### INTRODUCTION

Before you answer any questions in this survey, you need to know the following:

- This is a survey that measures tobacco, drug and alcohol use and other information relating to tobacco, alcohol, inhalant and drug use;
- This survey is being given to as many as 300,000 students in the state of Texas;
- There is no way for anyone to know how you respond to the questions; and
- You can skip any question or not take the survey at all if you do not want to. There is no penalty if you skip questions or leave the survey blank.

If you feel you need to talk with someone about problems with tobacco, inhalants, alcohol or drugs, counselors from the Worker's Assistant Program of Texas are available by calling this toll free number: 1-877-9-NO-DRUG (1-877-966-3784).

This research study has been reviewed and approved by the Institutional Review Board — Humas Subjects in Research, Texas A&M University. For research-related questions regarding subjects' rights, the Institutional Review Board may be contacted at (979) 458-4087 or intotacted at (979) 4

#### DIRECTIONS

- · DO NOT write your name anywhere on this booklet.
- · Use a NUMBER 2 PENCIL only.
- · Fill in only ONE BUBBLE for each part of a question.
- · Be sure to read each question carefully.

#### **EXAMPLE QUESTION**







5 C I N T R O N EM-98505-10-654321

PLEASE DO NOT WRITE IN THIS AREA

500055

1. Are you a:	During the current school year, do you qualify for a free or reduced price school lunch?
Boy Girl	
0 0	Yes No Don't Know
	0 0 0
2. What grade are you in?	10. Do one or both of your parents usually attend
	school-sponsored open houses or PTA meetings?
4 5 6	Yes No
0 0 0	0 0
Do you live with both of your parents?	Below is a list of things some people sniff to get high. These are called inhalants.
Yes No	11. Have you EVER sniffed any of the following inhalants to get high?
0 0	(DARKEN ONE BUBBLE FOR EACH DRUG)
	Never Used Used
4. How old are you?	a. Gasoline to get high
4. How old are your	b. Paint thinner to get high
	c. Glue to get high
8 or younger 9 10 11 12 older	d. Whiteout, correction fluid, or magic markers to get high
	magic markers to get night
	a: Stray noint to not high
0 0 0 0 0	e. Spray paint to get high  f Computer dusting sprays to get high
5. Are you:  African Mexican Asian Native	f Computer dusting sprays to get high og. Other inhalants to get high of the g
5. Are you:  African Mexican Asian Native	g. Other inhalants to get high O  The following questions ask about drugs people use.  12. How many times have you used any of these drugs (DARKEN ONE BUBBLE FOR EACH DRUG)
5. Are you:  White American American American American Other  O O O  6. What grades do you usually get?  (ONLY CHOOSE ONE)	g. Other inhalants to get high O  The following questions ask about drugs people use.  12. How many times have you used any of these drugs (DARKEN ONE BUBBLE FOR EACH DRUG)
5. Are you: White American American American Other  O O O O  6. What grades do you usually get?	g. Other inhalants to get high  The following questions ask about drugs people use.  12. How many times have you used any of these drugs (DARKEN ONE BUBBLE FOR EACH DRUG)  (DARKEN ONE BUBBLE FOR EACH DRUG)
5. Are you:  White American American American American Other  O O O O  6. What grades do you usually get? (ONLY CHOOSE ONE)  Mostly Mostly Mostly Mostly Mostly	g. Other inhalants to get high  The following questions ask about drugs people use.  12. How many times have you used any of these drugs (DARKEN ONE BUBBLE FOR EACH DRUG)  (DARKEN ONE BUBBLE FOR EACH DRUG)  a. Cigarettes?
5. Are you:  White American American American American Other  O O O O  6. What grades do you usually get?  (ONLY CHOOSE ONE)  Mostly Mostly Mostly Mostly Mostly A's B's C's D's F's	g. Other inhalants to get high  The following questions ask about drugs people use.  12. How many times have you used any of these drugs (DARKEN ONE BUBBLE FOR EACH DRUG)  a. Cigarettes?  b. Snuff or Chewing
5. Are you:  White American American American American Other  O O O O  6. What grades do you usually get?  (ONLY CHOOSE ONE)  Mostly Mostly Mostly Mostly Mostly A's B's C's D's F's	g. Other inhalants to get high  The following questions ask about drugs people use.  12. How many times have you used any of these drugs (DARKEN ONE BUBBLE FOR EACH DRUG)  a. Cigarettes?  b. Snuff or Chewing Tobacco?
5. Are you:  White American American American American Other  O O O O  6. What grades do you usually get?  (ONLY CHOOSE ONE)  Mostly Mostly Mostly Mostly Mostly A's B's C's D's F's	The following questions ask about drugs people use.  12. How many times have you used any of these drugs (DARKEN ONE BUBBLE FOR EACH DRUG)  a. Cigarettes?  b. Snuff or Chewing
5. Are you:  White American American American American Other  6. What grades do you usually get?  (ONLY CHOOSE ONE)  Mostly Mostly Mostly Mostly Mostly A's B's C's D's F's  O O O	g. Other inhalants to get high  The following questions ask about drugs people use.  12. How many times have you used any of these drugs (DARKEN ONE BUBBLE FOR EACH DRUG)  a. Cigarettes?  b. Snuff or Chewing Tobacco?  c. Beer?
5. Are you:  White African American American American American Other  O O O O  6. What grades do you usually get? (ONLY CHOOSE ONE)  Mostly Mostly Mostly Mostly Mostly A's B's C's D's F's O O O O  7. Have you lived in this town for more than 3 years?  Yes No Don't Know	g. Other inhalants to get high  The following questions ask about drugs people use.  12. How many times have you used any of these drugs (DARKEN ONE BUBBLE FOR EACH DRUG)  a. Cigarettes? b. Snuff or Chewing Tobacco? c. Beer? d. Wine Coolers (hard lemonade, hard sodas, etc.)?
5. Are you:  White African Mexican American American American Other  6. What grades do you usually get?  (ONLY CHOOSE ONE)  Mostly Mostly Mostly Mostly Mostly A's B's C's D's F's   O O O O  7. Have you lived in this town for more than 3 years?	g. Other inhalants to get high  The following questions ask about drugs people use.  12. How many times have you used any of these drugs (DARKEN ONE BUBBLE FOR EACH DRUG)  a. Cigarettes? b. Snuff or Chewing Tobacco? c. Beer? d. Wine Coolers (hard lemonade, hard sodas, etc.)? e. Wine?
5. Are you:  White African American American American Other  O O O O  6. What grades do you usually get? (ONLY CHOOSE ONE)  Mostly Mostly Mostly Mostly Mostly A's B's C's D's F's O O O O  7. Have you lived in this town for more than 3 years?  Yes No Don't Know	g. Other inhalants to get high  The following questions ask about drugs people use.  12. How many times have you used any of these drugs (DARKEN ONE BUBBLE FOR EACH DRUG)  a. Cigarettes? b. Snuff or Chewing Tobacco? c. Beer? d. Wine Coolers (hard lemonade, hard sodas, etc.)? e. Wine? f. Liquor (whiskey, vodka,
5. Are you:  White African American American American American American American American Other  6. What grades do you usually get?  (ONLY CHOOSE ONE)  Mostly Mostly Mostly Mostly Mostly A's B's C's D's F's  O O O O  7. Have you lived in this town for more than 3 years?  Yes No Don't Know O O	g. Other inhalants to get high  The following questions ask about drugs people use.  12. How many times have you used any of these drugs (DARKEN ONE BUBBLE FOR EACH DRUG)  a. Cigarettes? b. Snuff or Chewing Tobacco? c. Beer? d. Wine Coolers (hard lemonade, hard sodas, etc.)? e. Wine? f. Liquor (whiskey, vodka, tequila, etc.)?
5. Are you:  White African American American American Other  O O O O  6. What grades do you usually get? (ONLY CHOOSE ONE)  Mostly Mostly Mostly Mostly Mostly A's B's C's D's F's O O O O  7. Have you lived in this town for more than 3 years?  Yes No Don't Know	g. Other inhalants to get high  The following questions ask about drugs people use.  12. How many times have you used any of these drugs (DARKEN ONE BUBBLE FOR EACH DRUG)  a. Cigarettes? b. Snuff or Chewing Tobacco? c. Beer? d. Wine Coolers (hard lemonade, hard sodas, etc.)? e. Wine? f. Liquor (whiskey, vodka,
5. Are you:  White American American American American Other  6. What grades do you usually get? (ONLY CHOOSE ONE)  Mostly Mostly Mostly Mostly Mostly A's B's C's D's F's  O O O O  7. Have you lived in this town for more than 3 years?  Yes No Don't Know O O O  8. Did either of your parents graduate from college?	g. Other inhalants to get high  The following questions ask about drugs people use.  12. How many times have you used any of these drugs (DARKEN ONE BUBBLE FOR EACH DRUG)  a. Cigarettes? b. Snuff or Chewing Tobacco? c. Beer? d. Wine Coolers (hard lemonade, hard sodas, etc.)? e. Wine? f. Liquor (whiskey, vodka, tequila, etc.)? g. Ozaril?
5. Are you:  White American American American American American American American Other  6. What grades do you usually get?  (ONLY CHOOSE ONE)  Mostly Mostly Mostly Mostly Mostly A's B's C's D's F's  O O O O  7. Have you lived in this town for more than 3 years?  Yes No Don't Know O O	g. Other inhalants to get high  The following questions ask about drugs people use.  12. How many times have you used any of these drugs (DARKEN ONE BUBBLE FOR EACH DRUG)  a. Cigarettes? b. Snuff or Chewing Tobacco? c. Beer? d. Wine Coolers (hard lemonade, hard sodas, etc.)? e. Wine? f. Liquor (whiskey, vodka, tequila, etc.)? g. Ozaril? h. Inhalants (whiteout,

13. SINCE SCHOOL BEGAN IN THE FALL, how many times have you used:  (DARKEN ONE BUBBLE FOR EACH DRUG)  Times used since school began in the Fall.  11. Sinuff or Chewing Tobacco?  C. Beer?	a. Cigarettes? b. Snuff or Chewing Tobacco? C. Beer? d. Wine Coolers (hard
d. Wine Coolers (hard lemonade, hard sodas, etc.)? e. Wine? f. Liquor (whiskey, vodka, tequila, etc.)? G. Oxaril? h. Inhalants (whiteout, spray paint, glue, gas, etc.) to get high? i. Marijuana (pot)?  14. How old were you when you first used: (DARKEN ONE BUBBLE FOR EACH DRUG)	lemonade, hard sodas, etc.)? e. Wine? f. Liquor (whiskey, vodka, tequila etc.)? g. Oxaril? h. Inhalants (whiteout, spray paint, glue, gas, etc.) to get high? i. Marijuana (pot)?  16. Has anyone ever tried to give you: (DARKEN ONE BUBBLE FOR EACH DRUG)
a. Cigarettes? b. Snuff or Chewing Tobacco? c. Beer? d. Wine Coolers (hard lemonade, hard sodas, etc.)? e. Wine? f. Liquor (whiskey, vodka, tequila, etc.)? g. Oxaril? h. Inhalants (whiteout, spray paint, glue, gas, etc.) to get high? i. Marijuana (pot)?	a. Cigarettes? b. Snuff cr Chewing Tobacco? c. Beer? d. Wine Coolers (hard lemonade, hard sodas, etc.)? e. Wine? f. Liquor (whiskey, vodka, tequila, etc.)? g. Oxaril? h. Inhalants (whiteout, spray paint, glue, gas, etc.) to get high? i. Marijuana (pot)?

PLEASE GO ON TO THE NEXT PAGE

17. When you drink alcohol (beer, wine coolers liquor), where do you get it?  (DARKEN ONE BUBBLE FOR EACH LINE)  Don't Drink V  a. Get it from home b. Get it from friends c. Get it from somewhere else	No a Boer O b. Wine Coolers O c. Wine O c. Win	ses or
18. How dangerous do you think it is for kids y age to use:  (DARKEN ONE BUB3LE FOR EACH DRUG)	z1. Since school began in the Fall, have you skipped school when your parents did know?	
a. Cigarettes?  b. Snuff or Chewing Tobacco?  c. Beer?	22. Since school began in the Fall, have you sent to anyone like the principal becausomething against the rules?	
d. Wine Coolers (hard lemonade, hard sodas, etc.)?  # Wine?  # Liquor (whiskey, vodka, tequila, etc.)?  # Oxaril?  h. Inhalants (whiteout, spray paint, glue, gas,	23. How do your parents feel about kids y smoking cigarettes?  They don't like it. They don't care. They think it's O.K. I don't know.	our age
etc.) to get high?  i. Marijuana (pot)?  OOOO  OOOOOOOOOOOOOOOOOOOOOOOOOOOOO	24. How do your parents feel about kids y drinking beer?  They don't like it. They don't care. They think it's O.K. I don't know.	O U O C
a. Your teacher? b. A visitor to your class? c. An assembly program? d. A school counselor?	No 25. How do your parents feel about kids y using marijuana? They don't like it. They don't care. They think it's O.K. I don't know.	your age

## TEXAS SCHOOL SURVEY OF SUBSTANCE USE

## SECONDARY SURVEY INSTRUMENT 2010 VERSION

#### INTRODUCTION

Before answering the survey questions in this booklet, please read the following:

- I am aware that the survey questions I am answering are designed to measure drug and alcohol use in my school district and the state of Texas, that I will be asked questions about student attitudes and behaviors that relate to substance abuse issues, and that over 300,000 students all over Texas will be taking this survey;
- I know that I do not have to take this survey and that I can skip any question that I do not choose to answer for any reason without any consequences;
- I know that there is no way for anyone to find out which survey that I answered, that the numbers on the questionnaire books are used only to identify which pages go together after the booklets are cut and cannot be used to tell who answered the questions; and
- > I am voluntarily participating in the Texas School Survey of Drug and Alcohol Use.


If you feel you need to talk with someone about problems with tobacco, inhalants, alcohol or drugs, counselors from the Worker's Assistant Program of Texas are available by calling this toll free number: 1-877-9-NO-DRUG (1-877-966-3784).

This research study has been reviewed and approved by the Institutional Review Board --- Human Subjects in Research, Texas A&M University. For research-related questions regarding subjects' rights, the Institutional Review Board may be contacted at (979) 458-4067 or inb@tamu.edu.

#### DIRECTIONS

- · DO NOT write your name anywhere on this booklet.
- · Use a NUMBER 2 PENCIL only.
- · Fill in only ONE BUBBLE for each part of a question.
- · Be sure to read each question carefully.

#### **EXAMPLE QUESTION**





FOR OFFICE

USE ONLY



5 C A N T I O N EM-98598-11:854321

PLEASE DO NOT WRITE IN THIS AREA

1762947

1. Are you:  Male Female	7. How long have you attended school in this District?  1 2-3 4 or year or less years more years.  O O O
2. What grade are you in?  7 8 9 10 11 12  O O O O O  3. Do you live with both of your parents?	8. Do you have a job? Yes No O  9. Do you get an allowance? Yes No O O O
Yes No O	10. Did either of your parents graduate from college?  Yes No Don't Know  O O O
11 or younger 12 13 14 15 16 17 18 19+	11. During the current school year, do you qualify for a free or reduced price school lunch?  Yes  No  Don't Know
African Mexican Asian Native White American American American American Other  6. On average what grades do you get? (DARKEN ONE BUBBLE ONLY)  Mostly Mostly Mostly Mostly Mostly A's B's C's D's F's  O O O	12. Do you regularly participate in any of the following extra-curricular activities? (DARKEN ONE BUBBLE FOR EACH LINE: a-m.)  Yes No  a. O School Athletics b. O School Band/Orchestra c. O School Choir d. O Drama, Speech, or Art Clubs e. O Drill Team or Cheerleading Squad f. O Student Government g. O Student Newspaper or Yearbook h. O Academic Clubs or Societies (math, science, foreign language, etc.) i. O Academic Competition Groups (UIL, science bowl, etc.) j. O Service Clubs at School (Key Club, Peer Mediators, PALS) k. O Other School Clubs or Student Groups l. O Athletic Teams OUTSIDE of School m. O Other Clubs or Groups OUTSIDE of School

13.	How safe do you feel when you are: (DARKEN ONE BUBBLE FOR EACH LINE: a-c.)	Very	Somewhat	Not Very	Not Safe	Don't
	a. in your home b. out in your neighborhood c. at school	Safe	Safe	Safe	At All	Know
14.	How many of your FRIENDS would you say: (DARKEN ONE BUBBLE FOR EACH LINE: a-e.)					
	Feel close to their parents?     Sometimes carry weapons like a knife or gure.     Care about making good grades?     Belong to a gang or are interested in become a gang member?	Ŏ	A Few	Some	Most	000
	e. Wish they could drop out of school?	ŏ	ŏ	Ö	ŏ	ŏ
15.	Do one or both of your parents usually attend so or PTA meetings?	chool-spor	nsored open hou		Yes (	⊃ No
Belo	ow is a list of things some people sniff to get high	n. They are	called INHALA	NTS.	Unidad	
16.	How recently, if ever, have you sniffed, huffed, o following INHALANTS for "kicks" or to get "hig (DARKEN ONE BUBBLE FOR EACH LINE: a-h.)		the Never heard of/Never used it	Used at least once in the Past Month	Used at least once Since Schoo Began in the Fall	Used at least once In Your Lifetime
	<ul><li>a. Spray paint to get high?</li><li>b. Whiteout, correction fluid, magic markers to</li><li>c. Computer dusting sprays to get high?</li></ul>		000	000	000	000
	<ul> <li>d. Helium, butane, propane, whippets (nitrous refrigerants/Freon gas to get high?</li> <li>e. Glue to get high?</li> <li>f. Toluene/tolly, paint thinner, other solvents to g. Gasoline, octane booster, carburetor cleaner</li> </ul>	o get high	_	0000	0000	0000
	h. Other aerosols/sprays (deodorant spray, Par room spray, etc.) to get high?	n, hair spr	ау,	0	0	0
17.	How recently, if ever, have you used the followin TOBACCO, ALCOHOL, and INHALANT products one time:  (DARKEN ONE BUBBLE FOR EACH LINE: a-g.)	even	18. How recen have you u following I even one t (DARKEN OI FOR EACH L	ised the	Wever neard of, Used at least Past No.	Used at least one Began in the Fall In Your Lifetime
a.	Cigarettes?	$\circ$	a. Marijuana? b. Cocaine (no c. Crack? d. Oxaril?	ot crack)?		0000
b. c. d.	Beer? O O O O O O O O O O O O O O O O O O O	8	Mushrooms f. Uppers? g. Downers?		) o l	0000
e. f.	Wine? O O O Liquor (vodka, rum,	0	h. Rohypnol (i roofies, etc i. Steroids? j. Ecstasy? k. Heroin (black	.)?	0 0	0000
a.	spray paint, glue, gas,	0	cheese, chi			0 0

A-8


17629	47	(	000		PLE	ASE DO NOT WRITE IN THIS AREA				
19. IN THE PAST 30 DAYS, you used: (DARKEN ONE BUBBL	E FOR	EACH LI	NE: a-f	20. About how many of your close friends use: (DARKEN ONE BUBBLE FOR EACH LINE: a-h.)						
a. Tobacco? b. Alcohol? c. Inhalants (whiteout, spray paint, glue, gas etc.) to get high? d. Marijuana? e. Cocaine (not crack)? f. Crack?	OOOO Oof Never heard	0000 001.21	.000		0000 0011+ Ilmes	a. Cigarettes? b. Smokeless Tobacco? c. Beer? d. Wine Coolers (hard lemonade, hard sodas, etc.)? e. Wine? f. Liquor (vodka, rum, whiskey, etc.)? g. Inhalants (whiteout, spray paint, glue, gas, etc.)? h. Marijuana?				
21. If you wanted some, h (DARKEN ONE BUBBL			NE: a-r	/.	1	22. How old were you when you first used: (DARKEN ONE BUBBLE FOR EACH LINE: a-r.)				
a. Cigarettes?	0000	0000	000	000 0	000 0	a. Cigarettes? b. Smokeless Tobacco? O O O O O O O O O O O O O O O O O O O				
e. Wine? f. Liquor (vodka, rum, whiskey, etc.)? g. Inhalants (whiteout,	0	0 0	00 0	0 0	0	e. Wine? OOOOOOO  f. Liquor (vodka, rum, whiskey, etc.)? g. Inhalants (whiteout, spray paint, glue,				
spray paint, glue, gas, etc.)? h. Marijuana? i. Cocaine (not crack)? j. Crack? k. Oxaril? l. Hallucinogens (LSD, PCP, Mushrooms,	0 1	00000	00000	00000	00000	gas, etc.)?  h. Marijuana? i. Cocaine (not crack)?				
etc.)? m. Uppers? n. Downers? o. Rohypnol (roches, roofles, etc.)?	00 0	000 000	000 00	000 00	000 00	etc.)?				
p. Steroids? q. Ecstasy? r. Heroin (black tar, cheese, chiva, brown heroin)?	0	000	00 0	000 0	00 0	p. Steroids? OOOOOOOO  q. Ecstasy? COOOOOOO  r. Herein (black tar, cheese, chiva, brown herein)?				

	1762947 OODSON ON WRITE IN THIS AREA									
23.	(DARKEN ONE BUBBLE FOR EACH LINE: a-r.) had 5 or more drinks of alcohol AT ONE T						IN THE PAST 30 DAYS, on how many days have you had 5 or more drinks of alcohol AT ONE TIME?			
	Cigarettes?									Never had 5 or more Alone None  2 days  2 to 5 days  10 or more days
a. b.		000	000	000	000	000	000		a.	Alcohol O O O O O O
d.	The Control of the Co									
e.	sodas. etc.)? () Wine? ()	00	00	00	00	00	00			
f.	Liquor (vcdka, rum, whiskey, etc.)?	0	0	0	0	0	0		26.	How often do you get alcoholic beverages from
	spray paint, glue, gas, etc.)? Oracine (not crack)? Crack?	0000	00000	00000	00000	00000	00000		20.	the following sources? (DARKEN ONE BUBBLE FOR EACH LINE: a-e.)
k. I. m.	Oxaril? Hallucinogens (LSD, PCP, Mushrooms, etc.)? Uppers? Downers?	0 000	0 000	0 000	0 000	0 000	00		a. b. c. d.	At home   Compared to the comp
n. o.	Rohypnol (roches, roofies, etc.)?						0		e.	Other source
p. q.	aa	000	000	000	000	000	000			
r.	Heroin (black tar, cheese, chiva, brown heroin)?	0	0	0	0	0	0			
									27.	How often do you use marijuana in the following
24.	When you drink alcoho you usually have AT O (DARKEN ONE BUBBLI	NE TIN	ΛE, on	avera	age: : a-d.)	ny dri	, ,			ways? (DARKEN ONE BUBBLE FOR EACH LINE: a-e.)
a. b. c. d.	b. Wine Coolers (hard lemonade, hard sodas, etc.)  c. Wine  D. Blunts  C. Bongs or water pipes  D. C. Bongs or water pipes									

= 28. =	How recently, if ever, have you used methamphetamine (speed, crystal meth, ice, or crank) even one time?	Never heard of/Never	Used least of in the Past N	once he	Used at least once Since School Began in	In Your
	a. Methamphetamine (speed, crystal meth, ice, or crank)	used it	Pastiv		the Fall	Lifetime
29.	How recently, if ever, have you taken the following PRESCRIPTION DRUGS only for the experience or feeling it caused even one time: (DARKEN ONE BUBBLE FOR EACH LINE: a-e.)		Used least of in to Past M	once he	Used at least once Since School Began in the Fall	Used at
-	a. Codeine cough syrup (Lean, Nods, AC/DC)?	0	(	-	0	0
-	b. OxyContin, Percodan, or Percocet?	Ö	9	2	O	0
	c. Vicodin, Lortab, or Lorcet?	0	Š	?	0	0
	d. Valium or Diazepam? e. Xanax or Alprazolam?	8	2	<	. 0	8
_					Used at	
30.	How recently, if ever, have you taken the following OVER- THE-COUNTER DRUGS only for the experience or feeling it caused even one time:	Never heard of/Never used it	Used least of In to Past N	once he fonth	least once Since School Began In the Fall	Used at least once In Your Lifetime
	a. DXM, Triple C's, Skittles, or Coricidin?	0		)	0	0
31.	How dangerous do you think it is for kids your age to use: (DARKEN ONE BUBBLE FOR EACH LINE: a-j.)  a. Tobacco? b. Alcohol? c. Inhalants (whiteout, spray paint, glue, gas, etc.)/ d. Marijuana? e. Cocaine (not crack)? f. Crack? g. Ecstasy? h. Steroids? i. Heroin (black tar, cheese, chiva, brown heroin)? j. Methamphetamine?  SINCE SCHOOL BEGAN IN THE FALL, on how many DAYS have you (DARKEN ONE BUBBLE FOR EACH LINE: a-g.)			17 Part Part	4-9	10+
3	a. missed a whole day of school because you "skipped" or "cut"?     b. missed a whole day of school because you were III?		None	days	days	days
	c. missed a whole day of school for some other reason?		ŏ	ŏ	0	ŏ
	d. been sent by a teacher to someone like the Principal, Dean, or 0	Suidance	_			
	Counselor because of your conduct or attitude?		0	0	0	0
	<ul> <li>had someone from your home be called to school because of your attitude?</li> </ul>	our condu	ict O	0	0	0
	f. gotten into trouble with your teachers because of your drinking	12	8	8	ŏ	8
	g. gotten into trouble with your teachers because of your drinking		ŏ	ŏ	ŏ	ŏ
33.	DURING THE PAST 12 MONTHS, how many times (if any) have you (DARKEN ONE BUBBLE FOR EACH LINE: a-h.)		None	1-3 times	4-9 times	10+ times
-	a. gotten into difficulties of any kind with your friends because of	your				
	drinking?		0	0	0	0
	b. driven a car when you've had a good bit to drink?		0	Ö	Ö	0
3	<ul> <li>been criticized by someone you were dating because of drinkin</li> <li>gotten into trouble with the police because of drinking?</li> <li>gotten into difficulties of any kind with your friends because of</li> </ul>		8	8	00	0
-	use?	, our uru		0	0	0
	f. driven a car when you've felt high from drugs? g. been criticized by someone you were dating because of drug us. h. gotten into trouble with the police because of drug use?	se?	000	000	000	000

A-11

34.	SINCE SCHOOL BEGAN IN THE FALL, o "drunk" or "stoned" on (DARKEN ONE BUBBLE FOR EACH LINE		any DAYS (if			d at least one	4-9	10+		
	a. beer, wine coolers, wine, or hard lin				O	,	days	days		
	,,,,	uorr			Ŏ	Ö	Õ	Õ		
	b. marijuana?				O .	Ó	Õ	0		
	c. inhalants?				0	0	0	0		
	d. some other drug(s)?				0	0	0	0		
6.8	THE COURT OF STREET									
25	Alcohol or other drugs are sometimes u	and at to	anana nastia	- Thinkin	an of the west		L- 4 A L- 7			
30.	(DARKEN ONE BUBBLE FOR EACH LINE	ised at te	enage partie	s. i ninkii	ng or the parti	es you attend	lea this sc	nool year	٠.	
	(DARKEN ONE BUBBLE FOR EACH LINE			Half	Most of			dn't attend		
		Never	Seldom	the time	the time	Always	Know a	ny parties	- 8	
	a. how often was alcohol used?	0	0	0	0	0	0	0		
	<ul> <li>b. how often were marijuana</li> </ul>									
	and/or other drugs used?	0	0	0	0	0	Ó	0		
<u> </u>		DIVERSE.	- Hilliam	all the same of th	Name of the local					
			ACM MARK	to this		the same and	Control of	The said of the	٩,	
36.	SINCE SCHOOL BEGAN IN THE FALL, h	ave you s	ought help,	other tha	n from family		1	res No		
	or friends, for problems in any way con	nected w	ith your use	(if any) o	f alcohol,		9	0 0		
	marijuana, or other drugs?							21 0.27		
To Vie			DAME OF SUR		navi on suce		No.			
5,00	THE REPORT OF SHARP STATES	1002							Η,	
37.	If you had a drug or alcohol problem an	d needed		38. SII	NCE SCHOOL	BEGAN IN TH	E FALL, ha	ive you		
	help, who would you go to?			go	tten any infor	mation on dr	igs or alco	hol from		
	(DARKEN ONE BUBBLE FOR EACH LINE	: a-h.) Y	es No	the following SCHOOL sources?						
	a. A counselor or program in school?	(	0		(DARKEN ONE BUBBLE FOR EACH LINE: a-h.)					
	b. Another adult in school (such as a n	urse		Yes No						
	or teacher)?		0		A cohool boo	léh alasa2				
				a. A school health class?  b. An assembly program at school?						
	c. A counselor or program outside of s		0							
	d. Your parents?		0	C.		dance counse	lor?	0 0		
	e. A medical doctor?	(	0 0	d.	A science cla	557	(	0 0		
	f. Your friends?	. (	0 0	e.	A social stud	y class?	(			
	g. Another adult (such as relative,			f.	A student group	or club meeting	at school? (	Ŏ Ŏ		
	clergyman, or other family friend)?	(	0 0	g.	An invited so					
	h. I wouldn't go to anyone.	ò	ň ň	h.		rce at school?		ŏŏ		
		`			Allottici soul	ice at selloon	,	) 0		
10-91						C. C. C.				
39.	How do your parents feel about kids yo	ur age.	(DARKEN C	NE BUB	BLE FOR FAC	H LINE: a.c.)			-	
		an again.	TET THE STATE OF		DEE I OII ENO	THE COUNTY			•	
	Strongly	Mildly		r Approve	Mildly	Stro	ngly	Don't		
	Disapprove	Disappro	ove nor Di	isapprove	Approve	Appr	ove	Know		
	a. using tobacco?	0		0	0	C	)	0		
	b. drinking alcohol?	0		Ó	Ŏ	C	)	Ö		
	c. using marijuana?	Ö		Ŏ	Ŏ	Č	)	Ŏ		
	101 - 101 -			1000			55	-	-	
		100			BANTIN	SEVER STATE	100000	THE STREET	ij-	
									-	
40.	IN THE PAST SCHOOL YEAR, have you	been test	ed for drug u	se by so	meone at scho	ool?	) Yes	O No	-	
			1115AC (27577)	THE PERSON		153		10001049	-	
					1 M 6 -	Mary Comment		E-V-	-	



Appendix B
Participating Districts

Diampica VIII	DIGEDICE ID	CER A EL A
DISTRICT NAME	DISTRICT ID	STRATA
Alpine	022-901	5A
Alvarado	126-901	1D
Alvin	020-901	3A
Arlington	220-901	1A
Balmorhea	195-902	5A
Benavides	066-901	5A
Booker	148-901	4B
Brooks County	024-901	5A
Bryan	021-902	3A
Canutillo	071-907	5A
Carrollton-Farmers Branch	057-903	1B
Celeste	116-902	4B
Conroe	170-902	1B
Corsicana	175-903	4A
Cotulla	142-901	5A
Deer Park	101-908	1C
Donna	108-902	5A
Driscoll	178-905	2B
Ector County	068-901	3A
Edcouch-Elsa	108-903	5A
Everman	220-904	1D
Ferris	070-905	5A
Fort Worth	220-905	1A
Friendswood	084-911	3B
Ft Hancock	115-901	5A
Galena Park	101-910	1B
Granbury	111-901	4A
Hidalgo	108-905	5A
Idalou	152-910	3B
Iraan-Sheffield	186-903	5A
Italy	070-907	5A
Keller	220-907	1B
Krum	061-905	1D
La Joya	108-912	5A
Lamar	079-901	1B
Laneville	201-903	4B
Leander	246-913	2A
Linden-Kildare	034-905	4B
Little Cypress-Mauriceville	181-908	2B
••		

Littlefield	140-904	5A
Los Fresnos	031-906	5A
Lumberton	100-907	2B
Lyford	245-902	5A
Marfa	189-901	5A
Marlin	073-903	4B
Maud	019-903	3B
Maypearl	070-915	5A
McAllen	108-906	5A
Mercedes	108-907	5A
Midway	161-903	4A
Mount Pleasant	225-902	4A
Nacogdoches	174-904	4A
Northside	015-915	1A
Olton	140-905	5A
Palmer	070-910	5A
Peaster	184-908	1D
Pecos-Barstow-Toyah	195-901	5A
Pharr-San Juan-Alamo	108-909	5A
Point Isabel	031-909	5A
Rio Hondo	031-911	5A
Roma	214-903	5A
Royse City	199-902	1D
San Antonio	015-907	1B
San Benito	031-912	5A
San Elizario	071-904	5A
San Perlita	245-904	5A
Santa Rosa	031-914	5A
Shallowater	152-909	3B
Sharyland	108-911	5A
South Texas	031-916	5A
Spring	101-919	1B
Stafford	079-910	1D
United	240-903	5A
Utopia	232-904	5A
Valley View	108-916	5A
Westbrook	168-903	4B
Whitehouse	212-906	3B
Ysleta	071-905	5A

Appendix C Classroom Selection Guidelines

## Texas School Survey of Substance Use

## **Teacher Lists for Random Sample Selection**

#### For elementary

- List <u>all</u> the teachers' names that teach in grades 4, 5 and 6;
- List the grade level that each teaches; and
- List the number of students per class.

#### For secondary

- Choose *either* a class period in which <u>all</u> students are present *or* a subject all students must be enrolled;
- List <u>all</u> teachers for that class period or that subject; and
- List the number of students *per grade level* in each of those classes.

#### Note:

If a subject has been chosen rather than a particular class period, and where a single teacher teaches multiple classes, a classroom identifier of some sort (e.g., the period) should be included in the column under the heading of "period".

PPRI will randomly pick entire classrooms until we reach the target numbers per grade level for the sample. Then, we'll send you the survey materials along with a Master List of those classes randomly selected to be surveyed.

## Helpful Hint:

Electronic submissions (i.e., as an attachment to an email) are encouraged for speedy turnaround, as the random sample selection is a computer program.

#### Remember:

Sixth graders are considered elementary students for the purposes of this survey.

Appendix D
School Survey Administration
District Coordinator's Responsibilities
Campus Coordinators' Responsibilities
Teachers' Guide

# Texas School Survey of Drug and Alcohol Abuse District Coordinator's Responsibilities- Paper-Pencil Administration

## 1. Distribution of the Survey Administration Guide

The Survey Administration Guide is designed to instruct the individuals who will administer the Texas School Survey at each campus. Each individual involved in the survey administration should receive a copy. All survey packets have a copy of the Survey Administration Guide enclosed. We have provided additional copies of this document in this mailing.

The guides should be sent to the campuses in the time to be distributed one week prior to the scheduled survey administration date. This will allow time for everyone involved to read through the information and resolve any questions.

#### 2. Distribution of Survey Instruments and Envelopes

Enclosed is a set of envelopes. There should be approximately one envelope per participating classroom. These envelopes will be used to seal the completed instruments after survey administration (see Campus Coordinators' Responsibilities).

At least one week prior to the scheduled survey date, District Coordinators should forward the appropriate envelopes to each campus. Each campus will need as many envelopes as there are classes participating.

#### 3. Packaging the Completed Surveys for Delivery to PPRI

Completed survey instruments will be sealed in envelopes and collected from each classroom. As the District Survey Coordinator, you should clearly designate to each Campus Coordinator where the surveys should be sent from there. All completed surveys should be collected at a central location. Completed surveys should remain in the sealed envelopes, and the envelopes from each campus should be grouped together and **CLEARLY LABELED** with the campus name. **BLANK SURVEYS SHOULD BE** returned to PPRI along with the classroom envelopes.

For shipping completed survey instruments to PPRI, District Coordinators should keep the original boxes in which the blank surveys and envelopes were shipped in. This will allow PPRI to send shipping labels to the district, without District Coordinators having to weigh each box. The survey materials should be boxed, and the boxes should be labeled sequentially (i.e., "one of three, two of three," etc.). Materials should then be returned to PPRI in a single mailing. Ship to:

Texas School Survey
PPRI / Texas A&M University
317 Dulie Bell Bldg.
College Station, TX 77843-4476

## Texas School Survey of Drug and Alcohol Abuse District Coordinator's Responsibilities- Online Administration

#### 1. Distribution of the Survey Administration Guide

The *Survey Administration Guide* enclosed in this mailing list is designed to instruct the individuals who will administer the *Texas School Survey* at each campus. Each individual involved in the survey administration should receive a copy.

The guides should be sent to the campuses in the time to be distributed one week prior to the scheduled survey administration date. This will allow time for everyone involved to read through the information and resolve any questions.

#### 2. Distribution of Online Survey Tokens and Envelopes

Enclosed is a set of envelopes. There should be approximately one envelope per classroom. Secondary (Grades 7–12) envelopes will contain 30 online survey tokens and Elementary (Grades 4-6) envelopes will contain 25 online survey tokens. These envelopes will be used to seal the online survey tokens after survey administration (see Campus Coordinators' Responsibilities).

At least one week prior to the scheduled survey date, District Coordinators should forward the appropriate envelopes to each campus.

#### 3. Packaging the Survey Materials for Delivery to PPRI

The online survey tokens will be sealed in envelopes and collected from each classroom. As the District Survey Coordinator, you should clearly designate to each Campus Coordinator where the classroom envelopes should be sent from there. All classroom envelopes should be collected at a central location. The classroom envelopes from each campus should be grouped together and CLEARLY LABELED with the campus name. UNUSED ONLINE SURVEY TOKENS SHOULD BE returned to PPRI along with the classroom envelopes.

For shipping classroom envelopes to PPRI, District Coordinators should keep the original boxes in which the survey materials and envelopes were shipped in. This will allow PPRI to send shipping labels to the district, without District Coordinators having to weigh each box. The survey materials should be boxed, and the boxes should be labeled sequentially (i.e., "one of three, two of three," etc.). Materials should then be returned to PPRI in a single mailing. Ship to:

Texas School Survey
PPRI / Texas A&M University
317 Dulie Bell Bldg.
College Station, TX 77843-4476

## Texas School Survey of Drug and Alcohol Use Campus Coordinators' Responsibilities- Paper-Pencil Administration

## 1. Distribution of the Guide to Survey Administration

Each Campus Coordinator will receive copies of the Survey Administration Guide and Procedure from the District Coordinator. The Guide should be distributed to all survey administrators approximately one week prior to the scheduled administration date so that everyone will have adequate time to read through the information and resolve any questions.

## 2. Distribution of Survey Instruments to Classes

The District Coordinator will send to each participating campus the appropriate number of survey instruments in a large envelope. Please do **not** seal the envelopes.

On the day the survey is to be given, envelopes filled with the survey instruments should be distributed to each survey administrator. The individuals administering the survey should have reviewed the *Survey Administration Guide and Procedure* (described above) prior to this point and should already be aware of their responsibilities.

## 3. Collecting Completed Survey Instruments

After the survey has been administered in each class, teachers will be instructed to seal the completed survey forms in the envelopes. Unused blank forms should be kept together **OUTSIDE THE ENVELOPE**. It is recommended that teachers not leave the room with the sealed envelopes. Rather, someone who did not participate in the actual survey administration should be sent around to collect the envelopes.

Once envelopes have been collected from every classroom, they should be forwarded to the location designated by the District Survey Coordinator where they will be collected and returned to PPRI in a **single mailing**.

## Texas School Survey of Drug and Alcohol Use Campus Coordinators' Responsibilities- Online Administration

## 1. Distribution of the Guide to Survey Administration

Each Campus Coordinator will receive copies of the Survey Administration Guide and Procedure from the District Coordinator. The Guide should be distributed to all survey administrators approximately one week prior to the scheduled administration date so that everyone will have adequate time to read through the information and resolve any questions.

## 2. Distribution of Survey Instruments to Classes

The District Coordinator will send to each participating campus the appropriate number of online survey tokens in a large envelope. When the materials arrive, Campus Coordinators should label each envelope to indicate which class it is along with **the campus name**. Do **not** seal the envelopes.

On the day the survey is to be given, envelopes filled with the online survey tokens should be distributed to each survey administrator. The individuals administering the survey should have reviewed the *Survey Administration Guide and Procedure* (described above) prior to this point and should already be aware of their responsibilities.

## 3. Collecting Completed Survey Instruments

After the survey has been administered in each class, teachers will be instructed to seal all used and unused survey tokens in the envelopes. It is recommended that teachers not leave the room with the sealed envelopes. Rather, someone who did not participate in the actual survey administration should be sent around to collect the envelopes.

Once envelopes have been collected from every classroom, they should be forwarded to the location designated by the District Survey Coordinator where they will be collected and returned to PPRI in a **single mailing**.

## Texas School Survey of Drug and Alcohol Use Survey Administration Guide- Paper-Pencil Administration

Your school district has elected to participate in the *Texas School Survey of Drug and Alcohol Use*. The survey is sponsored and designed by the Texas Department of State Health Services (DSHS) and is facilitated by the Public Policy Research Institute (PPRI) at Texas A&M University. The district needs your assistance in administering the survey. PPRI has prepared this Survey Administration Guide to provide information and instructions for conducting the survey project.

- Your classroom envelope should contain blanks surveys and an attached Classroom Identification form to be filled out during survey administration. You will find this Classroom Identification Form attached as a label to your classroom envelope. It is very important that you write the campus name and number of students enrolled on this label.
- Please read all instructions carefully before administering the survey. It is important that these procedures are followed so that administration of the survey is uniform throughout the district.
- Your attitude towards this survey is extremely important. If students sense the importance of this information, they will take the survey more seriously. Please stress to the students that their responses are **entirely anonymous**.
- In addition, your physical position in the classroom may influence the way students answer questions. While students are taking the survey, do not walk around the room. Please sit or stand in a part of the room away from students.
- Try to keep students as quiet as possible throughout survey administration. If students are allowed to talk about the contents of the survey or their own experiences, peer influence could cause students to either inflate or under-report their own drug and alcohol habits.
- The survey should take approximately 30-45 minutes to administer.
- To maintain student confidentiality, do not view completed survey instruments.
- Students must use a #2 pencil. Be prepared to provide pencils to students who may not have one. Surveys completed in pen or marker will not be processed.

## Texas School Survey of Drug and Alcohol Use Survey Administration Procedure- Paper-Pencil Administration

1.) Before handing out surveys, please tell students the following:

"This survey is for the Public Policy Research Institute at Texas A&M University. Survey results will be used for evaluating and planning substance abuse programs for our school district. No student is required to participate in this survey. This survey will ask questions about your experiences with drugs and alcohol. Do not turn the pages or begin answering questions until we have gone over the survey instructions."

2.) Make sure all students have a #2 pencil to complete the survey. Hand out the surveys and please tell students the following:

"I cannot answer questions about specific substances or tell you what questions mean. If you are not familiar with a substance listed on the survey, mark the category "Never heard of it."

- 3.) Go over the instructions on the attached survey form, using either the elementary or secondary form as appropriate. Be sure to read aloud the Introduction, Directions, and Example Question.
- 4.) Please emphasize the following points to the students:
- \*\*This survey is completely voluntary.
- \*\*No one will know how individual students answer questions.
- \*\*If you do not feel comfortable answering a question or feel you cannot answer a question honestly, leave it blank.
- \*\*A drink means a can or bottle of beer or wine cooler, a 4 ounce glass of wine, a shot of liquor, or a mixed drink.
- 5.) If students refuse to take the survey, have them work quietly at their seat or send them to an area designated by your campus coordinator.
- 6.) Collect all completed surveys and place in envelope. Please do not view completed surveys instruments. **Fill out the attached Classroom Identification Form.** PLEASE DO NOT PUT UNUSED SURVEYS IN THE ENVELOPE. Seal the envelope.
- 7.) Return classroom envelope and unused surveys to your campus coordinator.

## Texas School Survey of Drug and Alcohol Use Survey Administration Guide- Online Administration

Your school district has elected to participate in the *Texas School Survey of Drug and Alcohol Use*. The survey is sponsored and designed by the Texas Department of State Health Services (DSHS) and is facilitated by the Public Policy Research Institute (PPRI) at Texas A&M University. The district needs your assistance in administering the survey. PPRI has prepared this Survey Administration Guide to provide information and instructions for conducting the survey project.

- Your classroom envelope should contain a small piece of paper for each student with a number, which is an online survey token. The survey tokens are unique numbers that each student will need to log in to the online survey.
- You will also find a label on the front of your classroom envelope titled 'Classroom Identification form'. This form will need to be filled out during survey administration. It is very important that you write the **campus name** and **number of students enrolled** on this label.
- Please read all instructions carefully before administering the survey. It is important that these procedures are followed so that administration of the survey is uniform throughout the district.
- Your attitude towards this survey is extremely important. If students sense the importance of this information, they will take the survey more seriously. Please stress to the students that their responses are **entirely anonymous**. The tokens are completely random and there is no way to know which student responded to a particular survey.
- In addition, your physical position in the classroom may influence the way students answer questions. While students are taking the survey, do not walk around the room. Please sit or stand in a part of the room away from students.
- Try to keep students as quiet as possible throughout survey administration. If students are allowed to talk about the contents of the survey or their own experiences, peer influence could cause students to either inflate or under-report their own drug and alcohol habits.
- The survey should take approximately 30-45 minutes to administer.

Thank you very much for your assistance with this important research project.

Texas School Survey of Drug and Alcohol Use

## Survey Administration Procedure- Online Administration

1.) Have all participating students enter the following address into their web browser:

For grades 4-6: <a href="http://txss.tamu.edu/4-6/">http://txss.tamu.edu/4-6/</a>
For grades 7-12: <a href="http://txss.tamu.edu/7-12/">http://txss.tamu.edu/4-6/</a>

(After this step is complete, students need to click on the button on the center of the web page to be directed to the main survey page)

2.) Before handing out the online survey tokens, please tell students the following:

"This survey is for the Public Policy Research Institute at Texas A&M University. Survey results will be used for evaluating and planning substance abuse programs for our school district. No student is required to participate in this survey. Please do not begin answering questions until we have gone over the survey instructions."

- 3.) Hand out the online survey tokens and please tell students the following:
  - This is a survey that measures tobacco, drug, and alcohol use and other information relating to tobacco, alcohol, inhalant, and drug use;
  - This survey is being given to as many as 300,000 students in the state of Texas;
  - There is no way for anyone to know how you respond to the questions;
  - You can skip any questions or not take the survey at all if you do not want to. There is no penalty if you skip questions or leave the survey blank;
  - On the survey, a drink means a can or bottle of beer or wine cooler, a 4 ounce glass of wine, a shot of liquor, or a mixed drink;
  - I cannot answer questions about specific substances or tell you what questions mean;
  - If you are not familiar with a substance listed, mark the category "Never heard of it."
  - At the end of the survey, please make sure you click the 'Submit' button

If you feel you need to talk with someone about problems with tobacco, inhalants, alcohol, or drugs, counselors from the Worker's Assistant Program of Texas are available by calling this toll free number: 1-877-9-NO-DRUG (1-877-966-3784).

This study has been reviewed and approved by the Institutional Review Board - Human Subjects in Research, Texas A&M University. For research-related questions regarding subjects' rights, the Institutional Review Board may be contacted at (979) 458-4067 or <a href="mailto:irreduced-irre

- 6.) Tell the students to type the number on their paper in the token box and click 'ENTER'. The students will then need to read the instructions and click 'NEXT' to proceed to survey questions.
- 7.) If students refuse to take the survey, have them work quietly at their seat or send them to an area designated by your campus coordinator.
- 8.) Collect all papers with the survey tokens and place in envelope. <u>Please fill out the attached Classroom Identification Form.</u> Seal the envelope and return it to your campus coordinator.

Appendix E
Computation of Sample Weights

#### Weighting the Texas School Survey of Substance Abuse

#### **District Level Weights**

Within the selected districts, students were sampled via a simple random sample of classrooms (i.e., a sample in which all classrooms in the district had an equal chance of getting in) picked directly from the district. Every student in the sampled classrooms was asked to participate in the survey.

A weight  $w_{ilk}$  is assigned to each sampled classroom k, in grade l, in district i:

$$w_{ilk} = \frac{N_{il}}{n_{il}} \tag{1}$$

where,

 $N_{il}$  = total no. of classrooms in grade l, in district i = no. of classrooms sampled in grade l, in district i

Since every student in the sampled classrooms was included in the sample, each student's probability of selection is equal to his/her classroom's probability of selection. Thus, each sampled student in these districts is assigned his/her classroom weight. Therefore the weight  $w_{ilr}$  of the  $r^{th}$  sampled student in classroom k is,

$$W_{ilr} = W_{ilk} \tag{2}$$

<u>Remark</u>: In (1), if the true  $N_{il}$  is not known for a particular district, estimate it with,

$$\hat{N}_{il} = \frac{M_{il}}{\hat{c}_{il}} \tag{3}$$

where,  $\hat{c}_{il} = m_{il} / n_{il}$  and

 $M_{il}$  = total no. of students in grade l, in district i  $m_{il}$  = no. of students sampled in grade l, in district i

Notice that in this case, the weights in (1) reduce to,  $w_{ilk} = M_{il} / m_{il}$ .

#### **District Level Estimates**

The total no. of students in grade l, in district i, giving a positive response to a question is estimated as

$$\hat{Y}_{il} = \sum_{r=1}^{m_{il}} w_{ilr} y_{ilr} \tag{4}$$

where

$$y_{ilr} = \begin{cases} 1 & \text{if the } r^{\text{th}} \text{ student's response is positive} \\ 0 & \text{otherwise} \end{cases},$$

and  $w_{ilr}$  is as defined in (2).

The proportion of students in grade l, in district i, giving a positive response to a question is estimated as,

$$\hat{\overline{Y}}_{il} = \frac{\hat{Y}_{il}}{M_{il}}$$

#### Stratum Level Weights

The districts were selected from strata that were formed on the basis of two factors, namely, how urban the districts were and the levels of enrollment. That is, once the districts were grouped by the degree to which they were urban, strata were formed by collecting together districts of similar size within these urban classes. Next, a simple random sample of districts was chosen from each stratum. A few large districts were selected to the sample with probability one. These districts were grouped into two strata based on the relative sizes of the districts.

Stratum level weights are assigned to the sampled students as follows. The weight  $w_{ij}^{s}$ 

of the  $r^{th}$  sampled student, from grade l, in district i in stratum h is,

$$w_{hilr}^{s} = \frac{1}{p_{hi}} w_{ilr} \tag{5}$$

where

$$p_{hi} = \begin{cases} 1 & \text{if district } i \text{ was sampled with probability 1} \\ \phi_{hi} & \text{otherwise} \end{cases}, \tag{6}$$

and within grade l, for each district i in stratum h,  $w_{ilr}$  is as defined in (4), and  $\phi_{hi} = n_h / N_h$  with

 $n_h$  = no. of districts sampled from stratum h

 $N_h$  = total number of districts in stratum h

Notice that within stratum h,  $\phi_{hi}$  does not depend on district i.

#### **State Level estimates**

The total no. of students in grade l, giving a positive response to a question is estimated as

$$\hat{Y}_l = \sum_{h=1}^H \hat{Y}_{hl} \tag{7}$$

with

$$\hat{Y}_{hl} = \sum_{i=1}^{n_h} \sum_{r=1}^{m_{il}} w_{hilr}^s y_{ilr}$$
(8)

where, within grade l, for each district i in stratum h,  $y_{ilr}$ , and  $m_{il}$  are as defined above.

The proportion of students in grade l, in the state, giving a positive response to a question is estimated as,

$$\hat{\bar{Y}}_{l} = \frac{1}{M_{l}} \sum_{h=1}^{H} \hat{Y}_{hl} \tag{9}$$

where,

 $M_{l}$  =total number of students in grade l in state.

Appendix F Standard Error Computation

## Standard Error Computation for the Texas School Survey of Substance Use

The standard errors of the percentages were estimated via the SUDAAN software package. The variance estimation techniques in SUDAAN are specifically designed to address the complexity of the sample design in finite population sampling. The variance estimators generally available for these complex designs do not have closed form solutions and, therefore, require iterative approaches in computations that would yield solutions as close to the true value as possible. SUDAAN software package is available for this purpose and is gaining wide popularity in survey sampling applications.

In applying SUDAAN to the current survey which consisted of a multi-stage stratified sampling design, the input SAS data set requires the following re-coding of the stratification variable for strata 21 and 22 (due to certainty stratum 21 having only one district responding to survey):

```
if strata=22 then strata=21;

(Here: strata= stratum identifier) Next, the data set needs to be sorted as follows:

proc sort data=___;
by strata distid q2 class;
run;
```

Here, distid= district identifier, q2= grade identifier, class= classroom identifier.

```
The SUDAAN code needs to incorporate the following:
proc crosstab data=" " filetype=sas design=WOR;
nest strata distid q2 class;
```

The specified design WOR uses Taylor series variance estimation methods<sup>1</sup> in computing standard errors of percentages in the multi-stage stratified sampling design of this survey.

\_

<sup>&</sup>lt;sup>1</sup> See SUDAAN User's Manual (Volume 1), page 3-4.

Appendix G
Standard Error and Confidence Interval Tables

# **Elementary Grades – 2010**

## **Standard Error and Confidence Intervals**

All Students-Overall and by Grade

Past Year Use

Lifetime Use

Male-Overall and by Grade

**Past Year Use** 

Lifetime Use

Female-Overall and by Grade

**Past Year Use** 

Lifetime Use

Whites-Overall and by Grade

**Past Year Use** 

Lifetime Use

African American-Overall and by Grade

**Past Year Use** 

Lifetime Use

Mexican American-Overall and by Grade

Past Year Use

Lifetime Use

**Border-Overall and by Grade** 

**Past Year Use** 

Lifetime Use

Non-border-Overall and by Grade

Past Year Use

Lifetime Use

Lifetime Use of Selected Substances - Overall					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	5.6%	4.9%	6.3%	0.37%	
ALCOHOL	21.5%	20.0%	22.9%	0.73%	
INHALANTS	10.4%	9.7%	11.1%	0.35%	
MARIJUANA	1.9%	1.5%	2.2%	0.17%	
CIGARETTES	4.7%	4.1%	5.3%	0.29%	
SMOKELESS TOBACCO	1.5%	1.2%	1.8%	0.15%	
BEER	14.4%	13.1%	15.7%	0.65%	
WINE COOLERS	8.8%	8.0%	9.6%	0.41%	
WINE	10.4%	9.6%	11.3%	0.44%	
LIQUOR	5.7%	5.1%	6.4%	0.33%	

Past Year Use	of Selected Su	bstances	- Overall	
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	3.1%	2.6%	3.5%	0.22%
ALCOHOL	13.7%	12.6%	14.8%	0.55%
INHALANTS	7.5%	6.9%	8.0%	0.27%
MARIJUANA	1.3%	1.0%	1.6%	0.15%
CIGARETTES	2.5%	2.1%	2.9%	0.18%
SMOKELESS TOBACCO	0.9%	0.7%	1.1%	0.09%
BEER	8.2%	7.4%	9.1%	0.43%
WINE COOLERS	5.4%	4.9%	6.0%	0.27%
WINE	6.3%	5.7%	7.0%	0.32%
LIQUOR	3.6%	3.1%	4.2%	0.26%

Lifetime Use of Selected Substances - Grade 4						
CATEGORY	Prevalence	Boundaries		Standard		
		Lower	Upper	Error		
TOBACCO	3.9%	3.1%	4.7%	0.42%		
ALCOHOL	16.0%	14.5%	17.6%	0.78%		
INHALANTS	10.5%	9.5%	11.4%	0.48%		
MARIJUANA	0.7%	0.5%	0.8%	0.07%		
CIGARETTES	2.9%	2.3%	3.5%	0.30%		
SMOKELESS TOBACCO	1.3%	0.9%	1.7%	0.20%		
BEER	10.7%	9.4%	12.1%	0.66%		
WINE COOLERS	5.1%	4.4%	5.9%	0.38%		
WINE	6.7%	5.8%	7.6%	0.44%		
LIQUOR	2.7%	2.3%	3.1%	0.19%		

Past Year Use of Selected Substances - Grade 4					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	2.3%	1.9%	2.8%	0.23%	
ALCOHOL	10.6%	9.4%	11.9%	0.63%	
INHALANTS	7.4%	6.6%	8.3%	0.43%	
MARIJUANA	0.4%	0.3%	0.5%	0.05%	
CIGARETTES	1.6%	1.3%	2.0%	0.17%	
SMOKELESS TOBACCO	0.9%	0.6%	1.2%	0.16%	
BEER	6.8%	5.8%	7.8%	0.50%	
WINE COOLERS	3.2%	2.7%	3.6%	0.21%	
WINE	4.1%	3.6%	4.7%	0.28%	
LIQUOR	1.6%	1.3%	1.9%	0.14%	

Lifetime Use o	Lifetime Use of Selected Substances - Grade 5						
CATEGORY	Prevalence	Boundaries		Standard			
		Lower	Upper	Error			
TOBACCO	4.6%	3.7%	5.5%	0.45%			
ALCOHOL	19.6%	18.0%	21.1%	0.80%			
INHALANTS	9.0%	8.2%	9.9%	0.42%			
MARIJUANA	1.2%	0.8%	1.5%	0.16%			
CIGARETTES	3.7%	3.1%	4.4%	0.32%			
SMOKELESS TOBACCO	1.3%	0.9%	1.8%	0.22%			
BEER	13.3%	11.8%	14.8%	0.77%			
WINE COOLERS	7.2%	6.4%	8.0%	0.40%			
WINE	9.1%	8.1%	10.1%	0.50%			
LIQUOR	4.3%	3.6%	5.0%	0.36%			

Past Year Use of Selected Substances - Grade 5					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	2.2%	1.7%	2.8%	0.26%	
ALCOHOL	12.2%	10.8%	13.5%	0.66%	
INHALANTS	6.2%	5.6%	6.7%	0.26%	
MARIJUANA	0.7%	0.5%	1.0%	0.14%	
CIGARETTES	1.8%	1.4%	2.2%	0.19%	
SMOKELESS TOBACCO	0.7%	0.4%	1.0%	0.15%	
BEER	7.1%	5.9%	8.3%	0.59%	
WINE COOLERS	4.4%	3.8%	4.9%	0.29%	
WINE	5.6%	4.9%	6.3%	0.34%	
LIQUOR	2.7%	2.1%	3.2%	0.29%	

Lifetime Use of Selected Substances - Grade 6					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	8.4%	7.4%	9.5%	0.53%	
ALCOHOL	29.0%	26.9%	31.0%	1.03%	
INHALANTS	11.8%	10.5%	13.1%	0.65%	
MARIJUANA	3.8%	3.0%	4.6%	0.39%	
CIGARETTES	7.5%	6.5%	8.4%	0.47%	
SMOKELESS TOBACCO	1.9%	1.5%	2.4%	0.21%	
BEER	19.3%	17.4%	21.3%	0.96%	
WINE COOLERS	14.3%	12.8%	15.8%	0.75%	
WINE	15.6%	14.2%	16.9%	0.68%	
LIQUOR	10.3%	9.0%	11.6%	0.63%	

Past Year Use of Selected Substances - Grade 6					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	4.7%	4.1%	5.4%	0.33%	
ALCOHOL	18.6%	16.8%	20.3%	0.86%	
INHALANTS	8.8%	7.6%	9.9%	0.58%	
MARIJUANA	2.9%	2.2%	3.5%	0.34%	
CIGARETTES	4.2%	3.6%	4.7%	0.28%	
SMOKELESS TOBACCO	1.1%	0.8%	1.3%	0.14%	
BEER	11.0%	9.6%	12.3%	0.69%	
WINE COOLERS	8.9%	7.8%	10.0%	0.55%	
WINE	9.4%	8.3%	10.5%	0.54%	
LIQUOR	6.7%	5.7%	7.7%	0.50%	

Lifetime Use	Lifetime Use of Selected Substances - Overall Male					
CATEGORY	Prevalence	Bour	ndaries	Standard		
		Lower	Upper	Error		
TOBACCO	7.0%	6.1%	7.9%	0.46%		
ALCOHOL	24.0%	22.3%	25.7%	0.85%		
INHALANTS	11.2%	10.4%	12.0%	0.41%		
MARIJUANA	2.4%	2.0%	2.8%	0.22%		
CIGARETTES	5.7%	4.9%	6.4%	0.38%		
SMOKELESS TOBACCO	2.3%	1.9%	2.8%	0.23%		
BEER	16.9%	15.4%	18.4%	0.75%		
WINE COOLERS	9.0%	8.2%	9.8%	0.41%		
WINE	11.0%	9.9%	12.1%	0.54%		
LIQUOR	6.2%	5.5%	7.0%	0.37%		

LIQUOR	6.2%	5.5%	7.0%	0.37%		
Lifetime Use of Selected Substances - Grade 4 Male						
CATEGORY	Prevalence	Bour	ndaries	Standard		
		Lower	Upper	Error		
TOBACCO	4.9%	3.8%	5.9%	0.53%		
ALCOHOL	18.9%	17.1%	20.8%	0.92%		
INHALANTS	12.4%	11.1%	13.6%	0.62%		
MARIJUANA	0.9%	0.7%	1.1%	0.10%		
CIGARETTES	3.6%	2.7%	4.6%	0.47%		
SMOKELESS TOBACCO	1.8%	1.3%	2.2%	0.23%		
BEER	13.3%	11.6%	15.0%	0.84%		
WINE COOLERS	5.7%	4.9%	6.5%	0.41%		
WINE	7.2%	6.0%	8.4%	0.60%		
LIQUOR	3.0%	2.5%	3.5%	0.25%		

Lifetime Use o	Lifetime Use of Selected Substances - Grade 5 Male					
CATEGORY	Prevalence	Bour	ndaries	Standard		
		Lower	Upper	Error		
TOBACCO	5.9%	4.6%	7.1%	0.61%		
ALCOHOL	22.4%	20.4%	24.5%	1.01%		
INHALANTS	9.6%	8.6%	10.5%	0.48%		
MARIJUANA	1.5%	1.0%	1.9%	0.23%		
CIGARETTES	4.5%	3.7%	5.4%	0.41%		
SMOKELESS TOBACCO	2.1%	1.3%	2.9%	0.42%		
BEER	16.0%	14.1%	17.9%	0.95%		
WINE COOLERS	7.3%	6.3%	8.3%	0.48%		
WINE	9.6%	8.3%	10.9%	0.66%		
LIQUOR	4.8%	3.9%	5.6%	0.43%		

Lifetime Use of Selected Substances - Grade 6 Male					
CATEGORY	Prevalence	Bour	Standard		
CATEGORI	Trevalence				
		Lower	Upper	Error	
TOBACCO	10.4%	8.9%	11.8%	0.73%	
ALCOHOL	31.0%	28.5%	33.4%	1.24%	
INHALANTS	11.8%	10.4%	13.1%	0.69%	
MARIJUANA	4.9%	3.9%	5.8%	0.47%	
CIGARETTES	9.0%	7.7%	10.3%	0.65%	
SMOKELESS TOBACCO	3.1%	2.4%	3.8%	0.34%	
BEER	21.6%	19.3%	23.9%	1.17%	
WINE COOLERS	14.1%	12.5%	15.7%	0.81%	
WINE	16.3%	14.7%	17.9%	0.78%	
LIQUOR	11.0%	9.6%	12.3%	0.69%	

Past Year Use of Selected Substances - Overall Male					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	3.9%	3.3%	4.5%	0.30%	
ALCOHOL	14.9%	13.7%	16.2%	0.63%	
INHALANTS	7.8%	7.1%	8.5%	0.34%	
MARIJUANA	1.7%	1.3%	2.0%	0.18%	
CIGARETTES	3.0%	2.6%	3.5%	0.23%	
SMOKELESS TOBACCO	1.4%	1.1%	1.7%	0.16%	
BEER	9.5%	8.4%	10.5%	0.52%	
WINE COOLERS	5.3%	4.8%	5.8%	0.26%	
WINE	6.4%	5.7%	7.2%	0.39%	
LIQUOR	3.7%	3.1%	4.3%	0.30%	

Past Year Use o	of Selected Sub	stances - C	Grade 4 Male	<u>,</u>
CATEGORY	Prevalence	Bour	ndaries	Standard
		Lower	Upper	Error
TOBACCO	2.9%	2.2%	3.6%	0.34%
ALCOHOL	12.2%	10.7%	13.7%	0.75%
INHALANTS	8.5%	7.4%	9.6%	0.55%
MARIJUANA	0.6%	0.5%	0.8%	0.08%
CIGARETTES	2.0%	1.5%	2.5%	0.26%
SMOKELESS TOBACCO	1.3%	0.8%	1.7%	0.24%
BEER	8.0%	6.8%	9.2%	0.60%
WINE COOLERS	3.4%	2.9%	3.9%	0.25%
WINE	4.4%	3.6%	5.2%	0.41%
LIQUOR	1.7%	1.4%	2.1%	0.18%

Past Year Use of Selected Substances - Grade 5 Male					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	3.0%	2.2%	3.7%	0.38%	
ALCOHOL	13.8%	11.9%	15.7%	0.95%	
INHALANTS	6.4%	5.7%	7.2%	0.37%	
MARIJUANA	0.9%	0.5%	1.3%	0.20%	
CIGARETTES	2.2%	1.7%	2.7%	0.24%	
SMOKELESS TOBACCO	1.2%	0.6%	1.8%	0.30%	
BEER	8.9%	7.1%	10.7%	0.89%	
WINE COOLERS	4.2%	3.5%	4.9%	0.35%	
WINE	5.7%	4.8%	6.6%	0.45%	
LIQUOR	2.9%	2.0%	3.7%	0.42%	

Past Year Use of Selected Substances - Grade 6 Male				
CATEGORY	Prevalence	Bour	idaries	Standard
		Lower	Upper	Error
TOBACCO	5.8%	4.9%	6.8%	0.48%
ALCOHOL	19.0%	17.2%	20.7%	0.88%
INHALANTS	8.4%	7.1%	9.6%	0.61%
MARIJUANA	3.5%	2.7%	4.3%	0.39%
CIGARETTES	4.9%	4.2%	5.7%	0.38%
SMOKELESS TOBACCO	1.7%	1.2%	2.2%	0.25%
BEER	11.6%	10.1%	13.0%	0.73%
WINE COOLERS	8.4%	7.3%	9.5%	0.55%
WINE	9.3%	8.1%	10.5%	0.59%
LIQUOR	6.6%	5.6%	7.7%	0.51%

Lifetime Use of	Lifetime Use of Selected Substances - Overall Female				
CATEGORY	Prevalence	Bou	ndaries	Standard	
		Lower	Upper	Error	
TOBACCO	4.3%	3.6%	4.9%	0.33%	
ALCOHOL	19.0%	17.6%	20.4%	0.71%	
INHALANTS	9.7%	8.9%	10.5%	0.41%	
MARIJUANA	1.4%	1.1%	1.7%	0.15%	
CIGARETTES	3.7%	3.2%	4.3%	0.27%	
SMOKELESS TOBACCO	0.8%	0.5%	1.0%	0.11%	
BEER	12.0%	10.8%	13.3%	0.64%	
WINE COOLERS	8.7%	7.7%	9.6%	0.46%	
WINE	9.9%	9.1%	10.7%	0.41%	
LIQUOR	5.3%	4.6%	6.0%	0.34%	

CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	2.3%	1.9%	2.7%	0.22%
ALCOHOL	12.6%	11.5%	13.7%	0.55%
INHALANTS	7.2%	6.6%	7.7%	0.29%
MARIJUANA	1.0%	0.7%	1.3%	0.13%
CIGARETTES	2.0%	1.6%	2.4%	0.21%
SMOKELESS TOBACCO	0.4%	0.3%	0.5%	0.06%
BEER	7.1%	6.3%	7.9%	0.41%
WINE COOLERS	5.6%	4.9%	6.3%	0.34%
WINE	6.3%	5.6%	7.0%	0.34%
LIQUOR	3.6%	3.0%	4.1%	0.27%
Past Year Use of Selected Substances - Grade 4 Female				

Lifetime Use of Selected Substances - Grade 4 Female				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	3.0%	2.2%	3.8%	0.41%
ALCOHOL	13.3%	11.7%	14.9%	0.80%
INHALANTS	8.7%	7.7%	9.6%	0.48%
MARIJUANA	0.4%	0.3%	0.6%	0.10%
CIGARETTES	2.2%	1.6%	2.8%	0.29%
SMOKELESS TOBACCO	0.9%	0.4%	1.3%	0.22%
BEER	8.3%	7.1%	9.5%	0.60%
WINE COOLERS	4.6%	3.7%	5.5%	0.45%
WINE	6.2%	5.3%	7.1%	0.44%
LIQUOR	2.4%	1.9%	2.8%	0.23%

Past Year Use of Selected Substances - Grade 4 Female					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	1.8%	1.3%	2.2%	0.24%	
ALCOHOL	9.2%	7.8%	10.5%	0.66%	
INHALANTS	6.4%	5.6%	7.2%	0.39%	
MARIJUANA	0.2%	0.1%	0.4%	0.07%	
CIGARETTES	1.3%	0.8%	1.8%	0.25%	
SMOKELESS TOBACCO	0.5%	0.3%	0.8%	0.12%	
BEER	5.5%	4.5%	6.5%	0.50%	
WINE COOLERS	3.0%	2.3%	3.6%	0.32%	
WINE	3.9%	3.3%	4.5%	0.32%	
LIQUOR	1.5%	1.1%	1.9%	0.19%	

Lifetime Use of Selected Substances - Grade 5 Female					
CATEGORY	Prevalence	Bou	ndaries	Standard	
		Lower	Upper	Error	
TOBACCO	3.4%	2.6%	4.1%	0.39%	
ALCOHOL	16.9%	15.1%	18.6%	0.89%	
INHALANTS	8.6%	7.3%	9.8%	0.63%	
MARIJUANA	0.9%	0.6%	1.1%	0.13%	
CIGARETTES	3.0%	2.3%	3.7%	0.36%	
SMOKELESS TOBACCO	0.6%	0.4%	0.8%	0.11%	
BEER	10.8%	9.2%	12.4%	0.80%	
WINE COOLERS	7.1%	6.1%	8.0%	0.47%	
WINE	8.7%	7.7%	9.6%	0.48%	
LIQUOR	3.9%	3.2%	4.6%	0.37%	

Past Year Use of Selected Substances - Grade 5 Female				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	1.6%	1.1%	2.0%	0.21%
ALCOHOL	10.6%	9.4%	11.8%	0.59%
INHALANTS	5.9%	5.3%	6.5%	0.32%
MARIJUANA	0.6%	0.3%	0.8%	0.11%
CIGARETTES	1.4%	1.0%	1.8%	0.20%
SMOKELESS TOBACCO	0.3%	0.2%	0.4%	0.05%
BEER	5.4%	4.5%	6.3%	0.43%
WINE COOLERS	4.5%	3.8%	5.2%	0.34%
WINE	5.5%	4.9%	6.2%	0.33%
LIQUOR	2.5%	1.9%	3.0%	0.27%

Lifetime Use of Selected Substances -Grade 6 Female					
CATEGORY	Prevalence	Bou	ndaries	Standard	
		Lower	Upper	Error	
TOBACCO	6.6%	5.6%	7.6%	0.51%	
ALCOHOL	27.1%	25.0%	29.1%	1.02%	
INHALANTS	11.9%	10.3%	13.5%	0.79%	
MARIJUANA	2.8%	2.1%	3.5%	0.37%	
CIGARETTES	6.1%	5.2%	6.9%	0.42%	
SMOKELESS TOBACCO	0.8%	0.5%	1.1%	0.15%	
BEER	17.2%	15.3%	19.1%	0.95%	
WINE COOLERS	14.5%	13.0%	16.1%	0.78%	
WINE	14.9%	13.5%	16.3%	0.70%	
LIQUOR	9.7%	8.4%	11.0%	0.66%	

Past Year Use of Selected Substances - Grade 6 Female					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	3.7%	2.9%	4.4%	0.37%	
ALCOHOL	18.2%	16.1%	20.2%	1.02%	
INHALANTS	9.2%	7.8%	10.6%	0.72%	
MARIJUANA	2.2%	1.6%	2.9%	0.33%	
CIGARETTES	3.4%	2.7%	4.1%	0.33%	
SMOKELESS TOBACCO	0.4%	0.3%	0.6%	0.09%	
BEER	10.4%	8.9%	11.8%	0.73%	
WINE COOLERS	9.4%	8.0%	10.8%	0.69%	
WINE	9.5%	8.2%	10.7%	0.63%	
LIQUOR	6.8%	5.6%	8.1%	0.62%	

Lifetime Use of Selected Substances - Overall White					
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	5.1%	3.8%	6.4%	0.67%	
ALCOHOL	19.2%	16.9%	21.6%	1.17%	
INHALANTS	9.5%	8.4%	10.6%	0.56%	
MARIJUANA	1.0%	0.6%	1.4%	0.19%	
CIGARETTES	3.6%	2.6%	4.7%	0.52%	
SMOKELESS TOBACCO	2.2%	1.6%	2.8%	0.28%	
BEER	12.8%	10.6%	14.9%	1.07%	
WINE COOLERS	7.0%	5.9%	8.1%	0.54%	
WINE	10.4%	9.4%	11.5%	0.53%	
LIQUOR	5.0%	4.0%	6.0%	0.49%	

Past Year Use of S	Selected Subst	ances - Ov	erall White	e
CATEGORY	Prevalence	Bound	laries	Standard
		Lower	Upper	Error
TOBACCO	2.8%	2.0%	3.6%	0.41%
ALCOHOL	11.7%	10.1%	13.3%	0.80%
INHALANTS	6.4%	5.6%	7.3%	0.41%
MARIJUANA	0.6%	0.3%	0.9%	0.17%
CIGARETTES	1.9%	1.3%	2.5%	0.31%
SMOKELESS TOBACCO	1.3%	0.9%	1.7%	0.20%
BEER	6.9%	5.5%	8.3%	0.70%
WINE COOLERS	4.2%	3.4%	4.9%	0.36%
WINE	6.1%	5.3%	6.8%	0.39%
LIQUOR	3.1%	2.4%	3.8%	0.35%

Lifetime Use of Se	elected Substa	nces - Gr	ade 4 Wh	ite
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	3.6%	2.5%	4.6%	0.51%
ALCOHOL	14.2%	12.2%	16.2%	1.02%
INHALANTS	10.7%	9.1%	12.4%	0.83%
MARIJUANA	0.4%	0.2%	0.5%	0.08%
CIGARETTES	2.3%	1.4%	3.1%	0.44%
SMOKELESS TOBACCO	1.6%	1.1%	2.2%	0.25%
BEER	8.8%	6.9%	10.7%	0.94%
WINE COOLERS	4.5%	3.4%	5.5%	0.52%
WINE	6.8%	5.6%	8.0%	0.60%
LIQUOR	1.9%	1.3%	2.6%	0.32%

Past Year Use of S	elected Substa	ances - Gra	de 4 Whit	е
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	2.2%	1.4%	3.1%	0.42%
ALCOHOL	8.7%	7.0%	10.5%	0.86%
INHALANTS	7.7%	6.3%	9.1%	0.71%
MARIJUANA	0.2%	0.1%	0.3%	0.07%
CIGARETTES	1.4%	0.6%	2.2%	0.39%
SMOKELESS TOBACCO	1.2%	0.7%	1.6%	0.23%
BEER	5.2%	3.7%	6.7%	0.76%
WINE COOLERS	2.8%	2.3%	3.4%	0.28%
WINE	3.6%	2.7%	4.4%	0.44%
LIQUOR	1.3%	0.9%	1.7%	0.21%

Lifetime Use of Se	elected Substa	nces - Gr	ade 5 Wh	ite
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	4.8%	2.8%	6.8%	1.01%
ALCOHOL	19.3%	16.6%	21.9%	1.32%
INHALANTS	9.0%	7.4%	10.6%	0.82%
MARIJUANA	0.9%	0.2%	1.7%	0.39%
CIGARETTES	3.3%	1.8%	4.7%	0.72%
SMOKELESS TOBACCO	2.2%	1.4%	3.0%	0.42%
BEER	13.3%	10.8%	15.8%	1.26%
WINE COOLERS	6.6%	5.5%	7.8%	0.58%
WINE	10.4%	8.8%	11.9%	0.77%
LIQUOR	4.5%	3.1%	5.8%	0.68%

Past Year Use of S	elected Substa	ances - Gra	de 5 Whit	е
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	2.3%	1.3%	3.3%	0.50%
ALCOHOL	11.8%	9.6%	14.0%	1.10%
INHALANTS	5.4%	4.7%	6.1%	0.37%
MARIJUANA	0.6%	-0.1%	1.4%	0.39%
CIGARETTES	1.5%	0.8%	2.2%	0.35%
SMOKELESS TOBACCO	1.1%	0.6%	1.6%	0.24%
BEER	6.6%	4.6%	8.7%	1.03%
WINE COOLERS	3.9%	3.0%	4.7%	0.44%
WINE	6.5%	5.4%	7.5%	0.51%
LIQUOR	2.6%	1.6%	3.7%	0.52%

Lifetime Use of Selected Substances - Grade 6 White				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	7.0%	4.9%	9.1%	1.05%
ALCOHOL	24.2%	20.6%	27.8%	1.80%
INHALANTS	8.7%	7.2%	10.3%	0.76%
MARIJUANA	1.6%	1.0%	2.3%	0.33%
CIGARETTES	5.4%	3.7%	7.2%	0.88%
SMOKELESS TOBACCO	2.8%	1.8%	3.7%	0.48%
BEER	16.2%	12.8%	19.6%	1.69%
WINE COOLERS	9.9%	8.0%	11.8%	0.96%
WINE	14.1%	12.5%	15.7%	0.80%
LIQUOR	8.6%	6.7%	10.6%	0.96%

Past Year Use of S	elected Substa	ances - Gra	de 6 Whit	е
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	3.9%	2.6%	5.1%	0.64%
ALCOHOL	14.7%	12.2%	17.1%	1.24%
INHALANTS	6.2%	5.0%	7.4%	0.59%
MARIJUANA	1.0%	0.5%	1.4%	0.24%
CIGARETTES	2.9%	1.9%	3.8%	0.47%
SMOKELESS TOBACCO	1.6%	0.8%	2.4%	0.39%
BEER	8.8%	6.6%	11.0%	1.10%
WINE COOLERS	5.8%	4.4%	7.2%	0.69%
WINE	8.1%	6.8%	9.5%	0.66%
LIQUOR	5.4%	3.8%	7.0%	0.79%

Lifetime Use of Selected Substances - Overall African/American				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	6.7%	5.5%	8.0%	0.62%
ALCOHOL	23.8%	21.8%	25.7%	0.98%
INHALANTS	10.1%	9.1%	11.2%	0.54%
MARIJUANA	1.7%	1.0%	2.4%	0.35%
CIGARETTES	5.2%	4.6%	5.8%	0.29%
SMOKELESS TOBACCO	2.0%	1.1%	3.0%	0.46%
BEER	13.2%	11.3%	15.0%	0.93%
WINE COOLERS	12.3%	11.1%	13.5%	0.61%
WINE	11.4%	9.8%	13.0%	0.80%
LIQUOR	5.9%	4.6%	7.1%	0.63%

Past Year Use of Selecto	ed Substances -	Overall A	African/A	merican
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	3.1%	2.5%	3.8%	0.31%
ALCOHOL	14.6%	12.6%	16.7%	1.03%
INHALANTS	7.7%	6.9%	8.4%	0.38%
MARIJUANA	1.2%	0.7%	1.7%	0.24%
CIGARETTES	2.3%	1.6%	3.0%	0.37%
SMOKELESS TOBACCO	1.0%	0.5%	1.6%	0.30%
BEER	7.2%	5.8%	8.6%	0.69%
WINE COOLERS	6.9%	5.5%	8.3%	0.69%
WINE	6.4%	5.2%	7.6%	0.59%
LIQUOR	3.9%	2.6%	5.1%	0.62%

Lifetime Use of Selected	l Substances - 0	Grade 4 A	frican/Ar	nerican	
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	6.0%	3.4%	8.6%	1.29%	
ALCOHOL	19.6%	16.4%	22.8%	1.61%	
INHALANTS	10.5%	7.4%	13.6%	1.56%	
MARIJUANA	0.8%	0.2%	1.5%	0.34%	
CIGARETTES	3.7%	2.6%	4.7%	0.52%	
SMOKELESS TOBACCO	2.6%	0.6%	4.6%	1.00%	
BEER	11.3%	9.5%	13.2%	0.95%	
WINE COOLERS	8.4%	5.0%	11.8%	1.70%	
WINE	8.5%	6.4%	10.6%	1.06%	
LIQUOR	4.4%	3.2%	5.6%	0.60%	

Past Year Use of Selecte	d Substances -	Grade 4	African/A	merican
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	3.2%	2.2%	4.2%	0.51%
ALCOHOL	14.5%	12.0%	17.0%	1.24%
INHALANTS	7.9%	4.9%	10.9%	1.50%
MARIJUANA	0.5%	0.1%	0.8%	0.17%
CIGARETTES	1.7%	0.7%	2.8%	0.54%
SMOKELESS TOBACCO	1.5%	0.2%	2.9%	0.69%
BEER	8.5%	6.7%	10.3%	0.91%
WINE COOLERS	4.8%	3.5%	6.2%	0.67%
WINE	5.6%	3.9%	7.3%	0.85%
LIQUOR	2.6%	1.7%	3.5%	0.45%

Lifetime Use of Selected	l Substances - 0	Grade 5 A	frican/Ar	nerican
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	5.5%	4.3%	6.8%	0.62%
ALCOHOL	21.3%	18.1%	24.5%	1.61%
INHALANTS	8.3%	5.7%	10.9%	1.30%
MARIJUANA	0.9%	0.5%	1.3%	0.20%
CIGARETTES	4.7%	3.7%	5.7%	0.52%
SMOKELESS TOBACCO	1.4%	0.5%	2.3%	0.45%
BEER	12.3%	9.6%	15.0%	1.35%
WINE COOLERS	9.8%	7.3%	12.3%	1.23%
WINE	10.0%	7.9%	12.0%	1.03%
LIQUOR	3.2%	1.8%	4.5%	0.68%

Past Year Use of Selecte	Past Year Use of Selected Substances - Grade 5 African/American				
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	2.5%	1.2%	3.8%	0.64%	
ALCOHOL	12.3%	9.1%	15.4%	1.56%	
INHALANTS	5.8%	4.0%	7.6%	0.92%	
MARIJUANA	0.6%	0.3%	1.0%	0.17%	
CIGARETTES	2.1%	0.9%	3.3%	0.59%	
SMOKELESS TOBACCO	0.9%	0.1%	1.7%	0.41%	
BEER	5.6%	3.7%	7.5%	0.95%	
WINE COOLERS	6.1%	4.1%	8.1%	0.98%	
WINE	5.2%	3.8%	6.7%	0.74%	
LIQUOR	2.2%	1.2%	3.3%	0.52%	

Lifetime Use of Selected Substances - Grade 6 African/American				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	8.7%	6.8%	10.5%	0.94%
ALCOHOL	30.4%	26.3%	34.4%	2.03%
INHALANTS	11.6%	8.5%	14.7%	1.55%
MARIJUANA	3.4%	2.1%	4.7%	0.66%
CIGARETTES	7.3%	6.0%	8.5%	0.63%
SMOKELESS TOBACCO	2.2%	1.1%	3.3%	0.56%
BEER	15.9%	12.6%	19.3%	1.68%
WINE COOLERS	18.5%	15.4%	21.7%	1.56%
WINE	15.5%	11.6%	19.4%	1.95%
LIQUOR	10.0%	7.0%	13.1%	1.54%

Past Year Use of Selecte	Past Year Use of Selected Substances - Grade 6 African/American			
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	3.7%	2.7%	4.7%	0.48%
ALCOHOL	17.1%	12.8%	21.4%	2.15%
INHALANTS	9.2%	6.5%	11.9%	1.34%
MARIJUANA	2.6%	1.5%	3.6%	0.54%
CIGARETTES	3.1%	2.3%	3.8%	0.39%
SMOKELESS TOBACCO	0.7%	0.1%	1.3%	0.30%
BEER	7.5%	4.9%	10.1%	1.29%
WINE COOLERS	9.6%	6.5%	12.7%	1.56%
WINE	8.3%	5.7%	11.0%	1.34%
LIQUOR	6.7%	3.8%	9.6%	1.46%

Lifetime Use of Selected	l Substances - C	verall Me	exican/An	nerican
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	5.8%	5.1%	6.4%	0.31%
ALCOHOL	22.7%	21.3%	24.2%	0.72%
INHALANTS	11.0%	10.1%	11.9%	0.44%
MARIJUANA	2.5%	2.1%	2.9%	0.20%
CIGARETTES	5.3%	4.7%	5.9%	0.29%
SMOKELESS TOBACCO	1.0%	0.8%	1.1%	0.09%
BEER	16.2%	14.9%	17.5%	0.64%
WINE COOLERS	9.3%	8.5%	10.2%	0.42%
WINE	10.3%	9.3%	11.3%	0.51%
LIQUOR	6.4%	5.6%	7.1%	0.38%

Past Year Use of Selecte	ed Substances -	Overall N	/lexican/A	merican
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	3.4%	2.9%	3.8%	0.21%
ALCOHOL	15.2%	14.0%	16.3%	0.58%
INHALANTS	8.0%	7.2%	8.8%	0.38%
MARIJUANA	1.9%	1.5%	2.3%	0.19%
CIGARETTES	3.0%	2.6%	3.4%	0.20%
SMOKELESS TOBACCO	0.6%	0.5%	0.7%	0.06%
BEER	9.7%	8.7%	10.6%	0.48%
WINE COOLERS	6.1%	5.5%	6.8%	0.33%
WINE	6.6%	5.8%	7.4%	0.39%
LIQUOR	4.1%	3.5%	4.7%	0.29%

Lifetime Use of Selected	Lifetime Use of Selected Substances - Grade 4 Mexican/American				
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	3.5%	2.9%	4.2%	0.33%	
ALCOHOL	16.5%	14.7%	18.4%	0.92%	
INHALANTS	9.9%	8.8%	11.0%	0.53%	
MARIJUANA	0.8%	0.6%	1.0%	0.10%	
CIGARETTES	3.0%	2.4%	3.7%	0.32%	
SMOKELESS TOBACCO	0.8%	0.6%	1.0%	0.09%	
BEER	12.0%	10.3%	13.7%	0.85%	
WINE COOLERS	4.7%	4.1%	5.4%	0.32%	
WINE	6.2%	5.0%	7.4%	0.59%	
LIQUOR	2.8%	2.3%	3.3%	0.25%	

Past Year Use of Selecte	d Substances -	Grade 4 N	∕lexican/A	merican
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	2.2%	1.8%	2.5%	0.18%
ALCOHOL	10.9%	9.7%	12.2%	0.63%
INHALANTS	7.0%	6.1%	7.8%	0.44%
MARIJUANA	0.6%	0.4%	0.7%	0.06%
CIGARETTES	1.8%	1.5%	2.1%	0.16%
SMOKELESS TOBACCO	0.6%	0.4%	0.8%	0.10%
BEER	7.4%	6.3%	8.5%	0.55%
WINE COOLERS	3.0%	2.5%	3.5%	0.24%
WINE	4.1%	3.4%	4.9%	0.37%
LIQUOR	1.6%	1.2%	2.0%	0.20%

Lifetime Use of Selected Substances - Grade 5 Mexican/American				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	4.3%	3.8%	4.8%	0.26%
ALCOHOL	19.8%	18.1%	21.6%	0.86%
INHALANTS	9.2%	8.3%	10.0%	0.43%
MARIJUANA	1.4%	1.1%	1.7%	0.14%
CIGARETTES	3.9%	3.5%	4.4%	0.22%
SMOKELESS TOBACCO	0.8%	0.5%	1.0%	0.14%
BEER	14.0%	12.4%	15.5%	0.79%
WINE COOLERS	7.1%	6.3%	7.8%	0.37%
WINE	8.2%	7.1%	9.4%	0.58%
LIQUOR	4.7%	3.9%	5.6%	0.40%

Past Year Use of Selecte	d Substances -	Grade 5 N	/lexican/A	merican
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	2.2%	1.9%	2.6%	0.18%
ALCOHOL	12.8%	11.5%	14.1%	0.66%
INHALANTS	6.7%	5.9%	7.5%	0.40%
MARIJUANA	0.8%	0.6%	1.0%	0.10%
CIGARETTES	1.9%	1.6%	2.3%	0.16%
SMOKELESS TOBACCO	0.4%	0.2%	0.7%	0.11%
BEER	8.1%	7.1%	9.1%	0.51%
WINE COOLERS	4.3%	3.7%	5.0%	0.31%
WINE	5.2%	4.4%	6.0%	0.41%
LIQUOR	3.0%	2.3%	3.6%	0.33%

Lifetime Use of Selected Substances - Grade 6 Mexican/American				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	9.6%	8.8%	10.5%	0.42%
ALCOHOL	32.4%	30.6%	34.1%	0.88%
INHALANTS	14.1%	12.7%	15.5%	0.70%
MARIJUANA	5.6%	4.7%	6.4%	0.44%
CIGARETTES	9.2%	8.3%	10.0%	0.40%
SMOKELESS TOBACCO	1.3%	1.1%	1.6%	0.14%
BEER	22.9%	21.2%	24.7%	0.88%
WINE COOLERS	16.5%	15.0%	18.1%	0.76%
WINE	16.8%	15.5%	18.1%	0.63%
LIQUOR	11.9%	10.7%	13.0%	0.56%

Past Year Use of Selected Substances - Grade 6 Mexican/American				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	5.7%	5.1%	6.4%	0.31%
ALCOHOL	22.1%	20.6%	23.5%	0.72%
INHALANTS	10.5%	9.2%	11.8%	0.64%
MARIJUANA	4.3%	3.5%	5.2%	0.41%
CIGARETTES	5.5%	4.9%	6.1%	0.30%
SMOKELESS TOBACCO	0.8%	0.6%	1.0%	0.10%
BEER	13.7%	12.4%	15.1%	0.67%
WINE COOLERS	11.2%	10.0%	12.4%	0.61%
WINE	10.6%	9.4%	11.8%	0.60%
LIQUOR	8.0%	7.1%	8.9%	0.45%

Lifetime Use of Selected Substances - Overall Border					
CATEGORY	Prevalence	Boun	Standard		
		Lower	Upper	Error	
TOBACCO	4.8%	4.3%	5.3%	0.24%	
ALCOHOL	17.9%	16.2%	19.6%	0.86%	
INHALANTS	10.3%	9.1%	11.4%	0.57%	
MARIJUANA	2.0%	1.6%	2.4%	0.20%	
CIGARETTES	4.5%	4.0%	4.9%	0.23%	
SMOKELESS TOBACCO	0.8%	0.6%	1.0%	0.09%	
BEER	12.1%	10.9%	13.2%	0.58%	
WINE COOLERS	7.3%	6.5%	8.0%	0.39%	
WINE	7.2%	6.1%	8.4%	0.56%	
LIQUOR	4.1%	3.3%	4.9%	0.40%	

Lifetime Use of	Selected Subst	ances - Gr	ade 4 Bord	er
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	2.2%	1.9%	2.5%	0.15%
ALCOHOL	12.1%	10.4%	13.8%	0.85%
INHALANTS	10.2%	8.3%	12.1%	0.94%
MARIJUANA	0.6%	0.4%	0.7%	0.08%
CIGARETTES	1.8%	1.5%	2.2%	0.16%
SMOKELESS TOBACCO	0.5%	0.3%	0.6%	0.08%
BEER	8.3%	7.0%	9.5%	0.61%
WINE COOLERS	3.9%	3.1%	4.7%	0.40%
WINE	4.1%	2.9%	5.3%	0.59%
LIQUOR	1.7%	1.2%	2.1%	0.23%

Lifetime Use of	Selected Subst	ances - Gr	ade 5 Bord	er
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	3.7%	3.2%	4.1%	0.24%
ALCOHOL	15.5%	13.8%	17.2%	0.86%
INHALANTS	7.9%	6.8%	9.0%	0.54%
MARIJUANA	1.1%	0.8%	1.5%	0.18%
CIGARETTES	3.4%	3.0%	3.9%	0.23%
SMOKELESS TOBACCO	0.5%	0.4%	0.7%	0.08%
BEER	10.5%	9.3%	11.7%	0.60%
WINE COOLERS	5.8%	5.1%	6.6%	0.38%
WINE	5.2%	4.0%	6.4%	0.60%
LIQUOR	2.8%	2.1%	3.4%	0.32%

Lifetime Use of Selected Substances - Grade 6 Border					
Lifetiffie Ose of	Jeiecteu Jubst	ances - Gi	aue o boru	C1	
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	9.3%	8.1%	10.5%	0.58%	
ALCOHOL	27.6%	24.7%	30.4%	1.42%	
INHALANTS	13.1%	11.4%	14.8%	0.85%	
MARIJUANA	4.6%	3.6%	5.6%	0.50%	
CIGARETTES	8.8%	7.7%	9.9%	0.54%	
SMOKELESS TOBACCO	1.5%	0.9%	2.1%	0.28%	
BEER	18.5%	15.9%	21.0%	1.26%	
WINE COOLERS	12.8%	10.9%	14.8%	0.97%	
WINE	13.3%	12.0%	14.6%	0.64%	
LIQUOR	8.6%	7.0%	10.1%	0.77%	

Past Year Use of Selected Substances - Overall Border				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	2.8%	2.4%	3.2%	0.18%
ALCOHOL	11.6%	10.4%	12.8%	0.61%
INHALANTS	7.7%	6.7%	8.7%	0.51%
MARIJUANA	1.5%	1.1%	1.8%	0.16%
CIGARETTES	2.6%	2.3%	2.9%	0.17%
SMOKELESS TOBACCO	0.4%	0.3%	0.6%	0.09%
BEER	7.1%	6.5%	7.8%	0.34%
WINE COOLERS	4.6%	4.2%	5.1%	0.24%
WINE	4.5%	3.7%	5.2%	0.37%
LIQUOR	2.6%	2.1%	3.1%	0.24%

Past Year Use of Selected Substances - Grade 4 Border					
Past Year Use o	of Selected Sub	stances - Gr	ade 4 Borde	r	
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	1.4%	1.1%	1.7%	0.14%	
ALCOHOL	7.9%	6.6%	9.2%	0.66%	
INHALANTS	7.9%	6.1%	9.6%	0.89%	
MARIJUANA	0.4%	0.2%	0.5%	0.07%	
CIGARETTES	1.2%	0.9%	1.5%	0.16%	
SMOKELESS TOBACCO	0.2%	0.1%	0.4%	0.07%	
BEER	4.8%	4.2%	5.5%	0.34%	
WINE COOLERS	2.5%	1.9%	3.1%	0.29%	
WINE	2.6%	1.7%	3.5%	0.44%	
LIQUOR	1.0%	0.7%	1.2%	0.12%	

Past Year Use of Selected Substances - Grade 5 Border					
CATEGORY	Prevalence	Boun	Standard		
		Lower	Upper	Error	
TOBACCO	1.9%	1.5%	2.3%	0.21%	
ALCOHOL	9.3%	7.7%	10.8%	0.77%	
INHALANTS	5.4%	4.5%	6.3%	0.45%	
MARIJUANA	0.7%	0.4%	1.0%	0.16%	
CIGARETTES	1.8%	1.5%	2.2%	0.18%	
SMOKELESS TOBACCO	0.2%	0.1%	0.3%	0.07%	
BEER	5.8%	5.0%	6.6%	0.41%	
WINE COOLERS	3.5%	2.9%	4.2%	0.31%	
WINE	3.3%	2.5%	4.0%	0.38%	
LIQUOR	1.8%	1.2%	2.3%	0.27%	

Past Year Use o	f Selected Sub	stances - Gr	ade 6 Borde	r
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	5.5%	4.5%	6.5%	0.48%
ALCOHOL	18.8%	17.3%	20.2%	0.74%
INHALANTS	10.1%	9.0%	11.3%	0.58%
MARIJUANA	3.6%	2.8%	4.4%	0.39%
CIGARETTES	5.2%	4.3%	6.0%	0.44%
SMOKELESS TOBACCO	1.0%	0.6%	1.4%	0.22%
BEER	11.4%	10.1%	12.7%	0.65%
WINE COOLERS	8.5%	7.2%	9.7%	0.63%
WINE	8.1%	7.3%	9.0%	0.42%
LIQUOR	5.6%	4.7%	6.4%	0.41%

Lifetime Use of Selec	Lifetime Use of Selected Substances - Overall Non-Border				
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	5.8%	4.9%	6.7%	0.46%	
ALCOHOL	22.3%	20.5%	24.1%	0.91%	
INHALANTS	10.5%	9.6%	11.3%	0.41%	
MARIJUANA	1.8%	1.4%	2.3%	0.21%	
CIGARETTES	4.7%	4.0%	5.5%	0.36%	
SMOKELESS TOBACCO	1.7%	1.3%	2.1%	0.18%	
BEER	15.0%	13.4%	16.6%	0.81%	
WINE COOLERS	9.2%	8.2%	10.2%	0.51%	
WINE	11.2%	10.1%	12.3%	0.54%	
LIQUOR	6.1%	5.3%	7.0%	0.41%	

Lifetime Use of Selec	ted Substance	s - Overal	l Non-Boi	rder
CATEGORY	Prevalence	Boun	Boundaries	
		Lower	Upper	Error
TOBACCO	3.2%	2.6%	3.7%	0.28%
ALCOHOL	14.3%	12.9%	15.6%	0.68%
INHALANTS	7.3%	6.7%	8.0%	0.31%
MARIJUANA	1.3%	0.9%	1.6%	0.18%
CIGARETTES	2.5%	2.0%	2.9%	0.22%
SMOKELESS TOBACCO	1.0%	0.8%	1.2%	0.11%
BEER	8.5%	7.4%	9.6%	0.54%
WINE COOLERS	5.6%	5.0%	6.3%	0.34%
WINE	6.8%	6.0%	7.6%	0.40%
LIQUOR	3.9%	3.2%	4.6%	0.33%

Lifetime Use of Select	ted Substances	- Grade 4	Non-Bor	der
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	4.3%	3.3%	5.4%	0.52%
ALCOHOL	17.1%	15.2%	19.0%	0.97%
INHALANTS	10.5%	9.4%	11.6%	0.56%
MARIJUANA	0.7%	0.5%	0.9%	0.09%
CIGARETTES	3.2%	2.4%	3.9%	0.37%
SMOKELESS TOBACCO	1.5%	1.0%	2.0%	0.25%
BEER	11.4%	9.7%	13.1%	0.83%
WINE COOLERS	5.4%	4.5%	6.4%	0.47%
WINE	7.4%	6.3%	8.5%	0.55%
LIQUOR	2.9%	2.5%	3.4%	0.23%

Lifetime Use of Selec	ted Substances	- Grade	4 Non-Bo	rder
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	2.6%	2.0%	3.1%	0.28%
ALCOHOL	11.4%	9.8%	12.9%	0.78%
INHALANTS	7.2%	6.3%	8.2%	0.47%
MARIJUANA	0.4%	0.3%	0.5%	0.06%
CIGARETTES	1.8%	1.4%	2.2%	0.20%
SMOKELESS TOBACCO	1.1%	0.7%	1.5%	0.19%
BEER	7.3%	6.0%	8.5%	0.63%
WINE COOLERS	3.3%	2.8%	3.8%	0.25%
WINE	4.5%	3.8%	5.2%	0.35%
LIQUOR	1.8%	1.4%	2.1%	0.18%

Lifetime Use of Selected Substances - Grade 5 Non-Border				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	4.8%	3.7%	5.9%	0.56%
ALCOHOL	20.6%	18.6%	22.5%	0.96%
INHALANTS	9.3%	8.3%	10.4%	0.52%
MARIJUANA	1.2%	0.8%	1.6%	0.20%
CIGARETTES	3.8%	3.0%	4.6%	0.40%
SMOKELESS TOBACCO	1.5%	1.0%	2.1%	0.28%
BEER	14.0%	12.1%	15.9%	0.95%
WINE COOLERS	7.5%	6.5%	8.5%	0.49%
WINE	10.1%	9.0%	11.3%	0.57%
LIQUOR	4.7%	3.8%	5.6%	0.44%

Lifetime Use of Selected Substances - Grade 5 Non-Border				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	2.3%	1.7%	3.0%	0.32%
ALCOHOL	12.9%	11.3%	14.5%	0.80%
INHALANTS	6.3%	5.7%	6.9%	0.30%
MARIJUANA	0.7%	0.4%	1.1%	0.17%
CIGARETTES	1.8%	1.3%	2.2%	0.24%
SMOKELESS TOBACCO	0.9%	0.5%	1.3%	0.19%
BEER	7.4%	6.0%	8.9%	0.73%
WINE COOLERS	4.6%	3.8%	5.3%	0.36%
WINE	6.2%	5.4%	7.0%	0.39%
LIQUOR	2.9%	2.2%	3.6%	0.35%

Lifetime Use of Selected Substances - Grade 6 Non-Border				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	8.2%	7.0%	9.5%	0.64%
ALCOHOL	29.3%	26.8%	31.7%	1.22%
INHALANTS	11.5%	10.0%	13.1%	0.76%
MARIJUANA	3.6%	2.7%	4.6%	0.46%
CIGARETTES	7.2%	6.1%	8.3%	0.55%
SMOKELESS TOBACCO	2.0%	1.6%	2.5%	0.25%
BEER	19.5%	17.2%	21.8%	1.15%
WINE COOLERS	14.6%	12.9%	16.4%	0.89%
WINE	16.1%	14.4%	17.7%	0.82%
LIQUOR	10.7%	9.2%	12.2%	0.75%

Lifetime Use of Selected Substances - Grade 6 Non-Border				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	4.5%	3.8%	5.3%	0.39%
ALCOHOL	18.5%	16.4%	20.6%	1.04%
INHALANTS	8.5%	7.1%	9.8%	0.68%
MARIJUANA	2.7%	1.9%	3.5%	0.40%
CIGARETTES	3.9%	3.3%	4.6%	0.33%
SMOKELESS TOBACCO	1.1%	0.7%	1.4%	0.17%
BEER	10.8%	9.2%	12.5%	0.82%
WINE COOLERS	9.0%	7.7%	10.3%	0.66%
WINE	9.6%	8.3%	11.0%	0.67%
LIQUOR	7.0%	5.8%	8.2%	0.61%

## Secondary Grades – 2010

## **Standard Error and Confidence Intervals**

All Students-Overall and by Grade

Past 30 Day Use

Lifetime Use

Male-Overall and by Grade

Past 30 Day Use

Lifetime Use

Female-Overall and by Grade

Past 30 Day Use

Lifetime Use

Whites-Overall and by Grade

Past 30 Day Use

Lifetime Use

African American-Overall and by Grade

Past 30 Day Use

Lifetime Use

**Mexican American-Overall and by Grade** 

Past 30 Day Use

Lifetime Use

**Border-Overall and by Grade** 

Past 30 Day Use

Lifetime Use

Non-border-Overall and by Grade

Past 30 Day Use

**Lifetime Use** 

Lifetime Use of Selected Substances - Overall				
CATEGORY	Prevalence	Bound	aries	Standard
		Lower	Upper	Error
TOBACCO	30.5%	28.5%	32.4%	0.97%
CIGARETTES	28.7%	26.8%	30.5%	0.93%
SMOKELESS TOBACCO	9.5%	8.4%	10.5%	0.54%
ALCOHOL	61.8%	59.6%	64.0%	1.09%
INHALANTS	17.2%	16.3%	18.0%	0.43%
ANY ILLICIT DRUG	27.9%	25.9%	29.9%	1.01%
MARIJUANA	26.2%	24.1%	28.4%	1.08%
MARIJUANA ONLY	13.9%	12.7%	15.2%	0.63%
COCAINE OR CRACK	5.4%	4.8%	6.0%	0.30%
COCAINE	5.0%	4.4%	5.6%	0.30%
CRACK	1.6%	1.5%	1.8%	0.08%
HALLUCINOGENS	4.6%	4.0%	5.2%	0.31%
UPPERS	5.3%	4.5%	6.0%	0.36%
DOWNERS	5.7%	4.9%	6.5%	0.41%
ROHYPNOL	2.6%	2.2%	3.1%	0.24%
STEROIDS	1.4%	1.3%	1.5%	0.07%
ECSTASY	6.8%	6.1%	7.5%	0.34%
HEROIN	1.4%	1.3%	1.6%	0.07%
METHAMPHETAMINE	3.0%	2.8%	3.2%	0.10%
CODEINE	12.3%	11.3%	13.3%	0.51%
OXYCONTIN ETC	3.1%	2.7%	3.5%	0.22%
VICODIN ETC	5.3%	4.2%	6.4%	0.55%
VALIUM ETC	2.4%	2.0%	2.7%	0.18%
XANAX ETC	4.5%	3.6%	5.4%	0.44%
DXM ETC	5.4%	5.0%	5.8%	0.20%

Lifetime Use of Selected Substances - Overall Grade 7				
CATEGORY	Prevalence	Bound	aries	Standard
		Lower	Upper	Error
TOBACCO	15.1%	13.7%	16.5%	0.69%
CIGARETTES	14.1%	12.8%	15.4%	0.66%
SMOKELESS TOBACCO	3.9%	3.4%	4.5%	0.28%
ALCOHOL	42.1%	39.4%	44.7%	1.33%
INHALANTS	19.3%	18.0%	20.6%	0.64%
ANY ILLICIT DRUG	11.5%	10.1%	13.0%	0.72%
MARIJUANA	9.5%	8.1%	10.8%	0.68%
MARIJUANA ONLY	5.1%	4.3%	6.0%	0.44%
COCAINE OR CRACK	2.5%	2.0%	2.9%	0.21%
COCAINE	1.8%	1.5%	2.2%	0.16%

Past 30-day Use of Selected Substances - Overall				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	12.5%	11.4%	13.6%	0.54%
CIGARETTES	11.2%	10.2%	12.2%	0.50%
SMOKELESS TOBACCO	3.7%	3.2%	4.1%	0.23%
ALCOHOL	29.0%	27.5%	30.6%	0.77%
INHALANTS	6.0%	5.7%	6.4%	0.19%
ANY ILLICIT DRUG	13.1%	12.0%	14.3%	0.56%
MARIJUANA	11.4%	10.3%	12.5%	0.56%
MARIJUANA ONLY	4.7%	4.1%	5.3%	0.29%
COCAINE OR CRACK	1.7%	1.5%	1.9%	0.10%
COCAINE	1.5%	1.3%	1.7%	0.10%
CRACK	0.5%	0.4%	0.5%	0.03%
HALLUCINOGENS	1.5%	1.4%	1.7%	0.07%
UPPERS	2.2%	1.9%	2.5%	0.14%
DOWNERS	2.4%	2.0%	2.7%	0.18%
ROHYPNOL	1.0%	0.9%	1.2%	0.09%
STEROIDS	0.5%	0.4%	0.5%	0.04%
ECSTASY	2.5%	2.2%	2.9%	0.18%
HEROIN	0.5%	0.5%	0.6%	0.03%
METHAMPHETAMINE	0.8%	0.7%	0.9%	0.04%
CODEINE	4.8%	4.3%	5.3%	0.25%
OXYCONTIN ETC	1.0%	0.8%	1.1%	0.07%
VICODIN ETC	1.8%	1.4%	2.3%	0.22%
VALIUM ETC	0.7%	0.6%	0.9%	0.06%
XANAX ETC	1.7%	1.3%	2.2%	0.23%
DXM ETC	1.9%	1.7%	2.1%	0.09%

Past 30-day Use of Selected Substances - Overall Grade 7				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	4.8%	4.2%	5.3%	0.27%
CIGARETTES	4.3%	3.8%	4.8%	0.25%
SMOKELESS TOBACCO	1.3%	1.0%	1.6%	0.14%
ALCOHOL	15.0%	13.5%	16.6%	0.76%
INHALANTS	8.4%	7.8%	9.1%	0.32%
ANY ILLICIT DRUG	5.2%	4.4%	6.0%	0.40%
MARIJUANA	3.9%	3.2%	4.5%	0.34%
MARIJUANA ONLY	1.7%	1.4%	2.0%	0.15%
COCAINE OR CRACK	0.9%	0.7%	1.1%	0.09%
COCAINE	0.6%	0.5%	0.8%	0.07%

CRACK	1.5%	1.2%	1.8%	0.15%
HALLUCINOGENS	1.5%	1.3%	1.7%	0.12%
UPPERS	1.2%	1.0%	1.5%	0.13%
DOWNERS	1.2%	1.0%	1.5%	0.12%
ROHYPNOL	1.4%	1.1%	1.7%	0.15%
STEROIDS	1.3%	1.1%	1.4%	0.09%
ECSTASY	2.3%	1.9%	2.6%	0.18%
HEROIN	1.4%	1.2%	1.7%	0.13%
METHAMPHETAMINE	2.1%	1.8%	2.4%	0.14%
CODEINE	7.6%	6.9%	8.4%	0.36%
OXYCONTIN ETC	0.9%	0.8%	1.1%	0.08%
VICODIN ETC	1.1%	0.8%	1.3%	0.13%
VALIUM ETC	0.9%	0.8%	1.1%	0.08%
XANAX ETC	1.3%	1.0%	1.5%	0.11%
DXM ETC	3.9%	3.5%	4.3%	0.21%

CRACK	0.5%	0.4%	0.7%	0.06%
HALLUCINOGENS	0.6%	0.5%	0.8%	0.08%
UPPERS	0.5%	0.4%	0.6%	0.06%
DOWNERS	0.5%	0.4%	0.6%	0.06%
ROHYPNOL	0.5%	0.3%	0.6%	0.07%
STEROIDS	0.4%	0.3%	0.5%	0.05%
ECSTASY	0.8%	0.6%	1.0%	0.10%
HEROIN	0.5%	0.4%	0.6%	0.05%
METHAMPHETAMINE	0.6%	0.5%	0.8%	0.07%
CODEINE	3.2%	2.8%	3.7%	0.22%
OXYCONTIN ETC	0.4%	0.3%	0.5%	0.05%
VICODIN ETC	0.4%	0.3%	0.6%	0.08%
VALIUM ETC	0.4%	0.2%	0.5%	0.06%
XANAX ETC	0.6%	0.4%	0.7%	0.07%
DXM ETC	1.6%	1.3%	1.9%	0.15%

Lifetime Use of Selected Substances - Overall Grade 8				
CATEGORY	Prevalence	Bound	aries	Standard
		Lower	Upper	Error
TOBACCO	23.2%	21.2%	25.1%	0.98%
CIGARETTES	21.7%	19.8%	23.5%	0.92%
SMOKELESS TOBACCO	6.3%	5.3%	7.2%	0.48%
ALCOHOL	55.1%	52.5%	57.7%	1.31%
INHALANTS	21.1%	19.9%	22.3%	0.61%
ANY ILLICIT DRUG	18.9%	16.8%	21.1%	1.08%
MARIJUANA	16.5%	14.3%	18.6%	1.07%
MARIJUANA ONLY	9.0%	7.7%	10.2%	0.63%
COCAINE OR CRACK	3.3%	2.7%	3.9%	0.30%
COCAINE	2.7%	2.2%	3.2%	0.26%
CRACK	1.8%	1.5%	2.0%	0.14%
HALLUCINOGENS	3.1%	2.6%	3.7%	0.27%
UPPERS	2.6%	2.1%	3.1%	0.25%
DOWNERS	2.8%	2.2%	3.3%	0.27%
ROHYPNOL	2.6%	2.0%	3.2%	0.29%
STEROIDS	1.5%	1.3%	1.8%	0.11%
ECSTASY	4.5%	3.7%	5.3%	0.40%
HEROIN	1.8%	1.4%	2.1%	0.18%
METHAMPHETAMINE	2.6%	2.2%	3.0%	0.20%
CODEINE	9.5%	8.5%	10.6%	0.53%
OXYCONTIN ETC	1.8%	1.4%	2.1%	0.18%
VICODIN ETC	2.1%	1.6%	2.6%	0.25%
VALIUM ETC	1.3%	0.9%	1.7%	0.18%

Past 30-day Use of Selected Substances - Overall Grade 8					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	7.5%	6.6%	8.4%	0.47%	
CIGARETTES	6.5%	5.8%	7.3%	0.39%	
SMOKELESS TOBACCO	2.2%	1.8%	2.6%	0.19%	
ALCOHOL	21.7%	19.6%	23.9%	1.08%	
INHALANTS	8.4%	7.7%	9.2%	0.37%	
ANY ILLICIT DRUG	8.9%	7.6%	10.2%	0.64%	
MARIJUANA	6.8%	5.8%	7.9%	0.53%	
MARIJUANA ONLY	2.9%	2.4%	3.5%	0.27%	
COCAINE OR CRACK	1.3%	1.1%	1.5%	0.12%	
COCAINE	1.0%	0.8%	1.2%	0.10%	
CRACK	0.6%	0.5%	0.8%	0.07%	
HALLUCINOGENS	1.1%	0.8%	1.3%	0.14%	
UPPERS	1.2%	0.9%	1.5%	0.17%	
DOWNERS	1.3%	0.9%	1.7%	0.19%	
ROHYPNOL	1.2%	0.8%	1.5%	0.18%	
STEROIDS	0.4%	0.3%	0.5%	0.05%	
ECSTASY	1.8%	1.3%	2.3%	0.25%	
HEROIN	0.7%	0.5%	1.0%	0.10%	
METHAMPHETAMINE	0.9%	0.7%	1.2%	0.13%	
CODEINE	4.4%	3.9%	4.9%	0.24%	
OXYCONTIN ETC	0.6%	0.4%	0.7%	0.08%	
VICODIN ETC	0.7%	0.5%	0.9%	0.11%	
VALIUM ETC	0.6%	0.4%	0.8%	0.10%	

XANAX ETC	2.2%	1.5%	2.9%	0.37%
DXM ETC	4.9%	4.1%	5.6%	0.37%

Lifetime Use of Selected Substances - Overall Grade 9				
CATEGORY	Prevalence	Bound	aries	Standard
		Lower	Upper	Error
TOBACCO	28.9%	27.0%	30.9%	0.97%
CIGARETTES	27.3%	25.5%	29.2%	0.92%
SMOKELESS TOBACCO	8.2%	7.2%	9.3%	0.54%
ALCOHOL	62.8%	60.6%	65.0%	1.08%
INHALANTS	17.7%	16.1%	19.3%	0.81%
ANY ILLICIT DRUG	26.7%	24.8%	28.6%	0.96%
MARIJUANA	25.2%	23.3%	27.0%	0.92%
MARIJUANA ONLY	13.3%	12.0%	14.6%	0.64%
COCAINE OR CRACK	4.7%	4.1%	5.4%	0.34%
COCAINE	4.3%	3.7%	4.9%	0.32%
CRACK	1.6%	1.4%	1.9%	0.13%
HALLUCINOGENS	4.6%	3.8%	5.3%	0.36%
UPPERS	4.7%	3.9%	5.5%	0.40%
DOWNERS	5.1%	4.3%	5.9%	0.41%
ROHYPNOL	3.0%	2.4%	3.5%	0.28%
STEROIDS	1.3%	1.0%	1.6%	0.13%
ECSTASY	6.7%	5.7%	7.6%	0.47%
HEROIN	1.5%	1.2%	1.8%	0.15%
METHAMPHETAMINE	2.9%	2.6%	3.3%	0.16%
CODEINE	12.3%	11.4%	13.3%	0.47%
OXYCONTIN ETC	2.9%	2.4%	3.3%	0.23%
VICODIN ETC	4.2%	3.3%	5.0%	0.43%
VALIUM ETC	2.1%	1.7%	2.5%	0.20%
XANAX ETC	4.0%	3.0%	5.0%	0.49%
DXM ETC	5.7%	5.1%	6.4%	0.33%

Lifetime Use of Selected Substances - Overall Grade 10					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	34.9%	32.8%	37.0%	1.05%	
CIGARETTES	32.8%	30.8%	34.8%	1.00%	
SMOKELESS TOBACCO	11.3%	9.7%	12.8%	0.76%	
ALCOHOL	69.3%	67.3%	71.3%	1.00%	
INHALANTS	16.5%	15.0%	18.0%	0.75%	
ANY ILLICIT DRUG	33.6%	31.0%	36.3%	1.35%	
MARIJUANA	32.2%	29.4%	35.0%	1.40%	

XANAX ETC	1.0%	0.6%	1.4%	0.21%
DXM ETC	2.1%	1.7%	2.6%	0.22%

Past 30-day Use of Selected Substances - Overall Grade 9				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	11.0%	9.8%	12.1%	0.57%
CIGARETTES	9.7%	8.6%	10.8%	0.56%
SMOKELESS TOBACCO	3.2%	2.7%	3.7%	0.25%
ALCOHOL	28.7%	27.0%	30.5%	0.88%
INHALANTS	6.4%	5.7%	7.1%	0.36%
ANY ILLICIT DRUG	12.6%	11.4%	13.7%	0.60%
MARIJUANA	11.1%	10.0%	12.3%	0.57%
MARIJUANA ONLY	4.6%	3.8%	5.3%	0.36%
COCAINE OR CRACK	1.7%	1.4%	2.0%	0.14%
COCAINE	1.6%	1.3%	1.9%	0.15%
CRACK	0.4%	0.3%	0.5%	0.04%
HALLUCINOGENS	1.8%	1.5%	2.0%	0.12%
UPPERS	2.2%	1.9%	2.5%	0.15%
DOWNERS	2.4%	2.0%	2.8%	0.20%
ROHYPNOL	1.1%	0.9%	1.4%	0.12%
STEROIDS	0.5%	0.3%	0.6%	0.06%
ECSTASY	2.6%	2.2%	3.1%	0.22%
HEROIN	0.5%	0.3%	0.6%	0.06%
METHAMPHETAMINE	0.8%	0.6%	1.0%	0.08%
CODEINE	5.0%	4.4%	5.6%	0.32%
OXYCONTIN ETC	1.1%	0.9%	1.3%	0.09%
VICODIN ETC	1.6%	1.3%	2.0%	0.17%
VALIUM ETC	0.7%	0.6%	0.8%	0.07%
XANAX ETC	1.6%	1.1%	2.2%	0.28%
DXM ETC	2.2%	1.8%	2.5%	0.17%

Past 30-day Use of Selected Substances - Overall Grade 10					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	13.9%	12.8%	15.0%	0.55%	
CIGARETTES	12.3%	11.3%	13.4%	0.53%	
SMOKELESS TOBACCO	4.5%	3.8%	5.2%	0.33%	
ALCOHOL	33.0%	31.6%	34.4%	0.70%	
INHALANTS	4.8%	4.3%	5.2%	0.23%	
ANY ILLICIT DRUG	16.1%	14.6%	17.6%	0.75%	
MARIJUANA	14.2%	12.7%	15.7%	0.74%	

MARIJUANA ONLY	17.2%	15.6%	18.9%	0.82%
COCAINE OR CRACK	6.3%	5.4%	7.3%	0.46%
COCAINE	5.9%	5.0%	6.8%	0.46%
CRACK	1.6%	1.4%	1.9%	0.13%
HALLUCINOGENS	5.6%	4.7%	6.4%	0.42%
UPPERS	7.0%	5.8%	8.3%	0.61%
DOWNERS	7.1%	6.0%	8.1%	0.54%
ROHYPNOL	3.1%	2.6%	3.6%	0.26%
STEROIDS	1.3%	1.0%	1.6%	0.14%
ECSTASY	8.0%	7.0%	8.9%	0.49%
HEROIN	1.4%	1.1%	1.7%	0.15%
METHAMPHETAMINE	3.1%	2.7%	3.4%	0.18%
CODEINE	14.6%	13.0%	16.3%	0.83%
OXYCONTIN ETC	4.0%	3.2%	4.8%	0.39%
VICODIN ETC	6.9%	5.1%	8.7%	0.90%
VALIUM ETC	2.9%	2.3%	3.4%	0.27%
XANAX ETC	6.0%	4.4%	7.6%	0.80%
DXM ETC	6.4%	5.7%	7.1%	0.33%

	•			
MARIJUANA ONLY	6.2%	5.3%	7.2%	0.47%
COCAINE OR CRACK	1.8%	1.5%	2.2%	0.15%
COCAINE	1.7%	1.4%	2.0%	0.15%
CRACK	0.4%	0.2%	0.5%	0.07%
HALLUCINOGENS	1.7%	1.2%	2.1%	0.23%
UPPERS	2.9%	2.3%	3.5%	0.29%
DOWNERS	3.0%	2.4%	3.7%	0.33%
ROHYPNOL	1.3%	1.0%	1.5%	0.13%
STEROIDS	0.4%	0.3%	0.6%	0.07%
ECSTASY	2.9%	2.2%	3.6%	0.34%
HEROIN	0.6%	0.5%	0.8%	0.08%
METHAMPHETAMINE	0.8%	0.6%	0.9%	0.07%
CODEINE	5.8%	4.5%	7.1%	0.64%
OXYCONTIN ETC	1.2%	0.9%	1.5%	0.15%
VICODIN ETC	2.4%	1.6%	3.2%	0.40%
VALIUM ETC	0.8%	0.6%	0.9%	0.09%
XANAX ETC	2.4%	1.3%	3.5%	0.55%
DXM ETC	2.1%	1.8%	2.4%	0.15%

Lifetime Use of Selected Substances - Overall Grade 11				
CATEGORY	Prevalence	Bound	aries	Standard
		Lower	Upper	Error
TOBACCO	39.7%	37.9%	41.5%	0.89%
CIGARETTES	37.2%	35.4%	39.0%	0.88%
SMOKELESS TOBACCO	13.7%	12.3%	15.2%	0.72%
ALCOHOL	70.2%	68.3%	72.1%	0.95%
INHALANTS	14.8%	13.6%	16.0%	0.61%
ANY ILLICIT DRUG	38.2%	36.4%	40.1%	0.95%
MARIJUANA	36.8%	34.6%	39.1%	1.14%
MARIJUANA ONLY	19.3%	17.7%	21.0%	0.82%
COCAINE OR CRACK	7.3%	6.5%	8.2%	0.43%
COCAINE	7.2%	6.3%	8.0%	0.43%
CRACK	1.6%	1.3%	1.9%	0.16%
HALLUCINOGENS	6.5%	5.7%	7.2%	0.38%
UPPERS	8.1%	7.2%	9.0%	0.44%
DOWNERS	9.1%	7.9%	10.2%	0.58%
ROHYPNOL	3.0%	2.2%	3.8%	0.38%
STEROIDS	1.5%	1.1%	1.8%	0.17%
ECSTASY	9.4%	8.3%	10.4%	0.52%
HEROIN	1.2%	0.9%	1.5%	0.15%
METHAMPHETAMINE	3.6%	3.1%	4.1%	0.25%
CODEINE	15.1%	13.9%	16.4%	0.63%

Past 30-day Use of Selected Substances - Overall Grade 11					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	17.8%	16.4%	19.2%	0.70%	
CIGARETTES	16.1%	14.9%	17.4%	0.62%	
SMOKELESS TOBACCO	5.3%	4.5%	6.1%	0.41%	
ALCOHOL	35.8%	34.1%	37.5%	0.86%	
INHALANTS	4.2%	3.6%	4.7%	0.27%	
ANY ILLICIT DRUG	17.7%	16.5%	18.8%	0.57%	
MARIJUANA	15.9%	14.6%	17.2%	0.65%	
MARIJUANA ONLY	6.2%	5.4%	7.1%	0.42%	
COCAINE OR CRACK	2.2%	1.8%	2.5%	0.17%	
COCAINE	2.1%	1.7%	2.4%	0.17%	
CRACK	0.5%	0.4%	0.6%	0.05%	
HALLUCINOGENS	2.0%	1.7%	2.3%	0.16%	
UPPERS	3.2%	2.8%	3.6%	0.21%	
DOWNERS	3.8%	3.2%	4.3%	0.27%	
ROHYPNOL	1.3%	0.9%	1.6%	0.16%	
STEROIDS	0.5%	0.3%	0.7%	0.09%	
ECSTASY	3.5%	2.9%	4.0%	0.27%	
HEROIN	0.5%	0.3%	0.6%	0.08%	
METHAMPHETAMINE	0.9%	0.7%	1.0%	0.09%	
CODEINE	5.4%	4.8%	6.0%	0.31%	

OXYCONTIN ETC	4.5%	3.9%	5.1%	0.29%
VICODIN ETC	8.8%	7.3%	10.2%	0.73%
VALIUM ETC	3.3%	2.7%	3.9%	0.29%
XANAX ETC	7.0%	6.0%	7.9%	0.49%
DXM ETC	5.7%	5.1%	6.3%	0.30%

OXYCONTIN ETC	1.4%	1.1%	1.6%	0.14%
VICODIN ETC	3.0%	2.3%	3.7%	0.34%
VALIUM ETC	1.0%	0.7%	1.3%	0.16%
XANAX ETC	2.6%	2.0%	3.2%	0.29%
DXM ETC	1.8%	1.5%	2.1%	0.14%

Lifetime Use of Selected Substances - Overall Grade 12				
CATEGORY	Prevalence	Bound	aries	Standard
		Lower	Upper	Error
TOBACCO	45.1%	42.7%	47.4%	1.15%
CIGARETTES	42.8%	40.7%	45.0%	1.06%
SMOKELESS TOBACCO	15.1%	13.6%	16.6%	0.77%
ALCOHOL	75.0%	73.2%	76.9%	0.93%
INHALANTS	12.5%	11.3%	13.8%	0.63%
ANY ILLICIT DRUG	42.4%	40.3%	44.6%	1.08%
MARIJUANA	41.5%	39.2%	43.8%	1.15%
MARIJUANA ONLY	21.8%	20.0%	23.7%	0.92%
COCAINE OR CRACK	9.1%	8.0%	10.1%	0.53%
COCAINE	8.9%	7.8%	10.0%	0.54%
CRACK	1.6%	1.4%	1.9%	0.13%
HALLUCINOGENS	7.3%	6.3%	8.2%	0.46%
UPPERS	9.0%	7.8%	10.2%	0.60%
DOWNERS	10.1%	8.8%	11.5%	0.68%
ROHYPNOL	2.9%	2.1%	3.7%	0.40%
STEROIDS	1.7%	1.4%	2.0%	0.15%
ECSTASY	11.2%	9.7%	12.7%	0.74%
HEROIN	1.1%	0.9%	1.4%	0.10%
METHAMPHETAMINE	3.8%	3.4%	4.2%	0.21%
CODEINE	15.6%	14.2%	17.0%	0.70%
OXYCONTIN ETC	5.0%	4.4%	5.7%	0.32%
VICODIN ETC	10.0%	8.6%	11.4%	0.70%
VALIUM ETC	4.1%	3.5%	4.8%	0.34%
XANAX ETC	7.6%	6.4%	8.7%	0.58%
DXM ETC	5.7%	4.9%	6.5%	0.40%

Past 30-day Use of Selected Substances - Overall Grade 12					
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	22.7%	20.9%	24.5%	0.90%	
CIGARETTES	20.9%	19.2%	22.5%	0.83%	
SMOKELESS TOBACCO	6.2%	5.4%	7.1%	0.43%	
ALCOHOL	43.4%	41.5%	45.2%	0.93%	
INHALANTS	3.2%	2.8%	3.7%	0.23%	
ANY ILLICIT DRUG	20.3%	18.8%	21.8%	0.73%	
MARIJUANA	18.4%	16.8%	19.9%	0.77%	
MARIJUANA ONLY	7.4%	6.6%	8.3%	0.42%	
COCAINE OR CRACK	2.2%	1.8%	2.6%	0.20%	
COCAINE	2.1%	1.7%	2.5%	0.20%	
CRACK	0.4%	0.2%	0.5%	0.07%	
HALLUCINOGENS	2.1%	1.7%	2.4%	0.19%	
UPPERS	3.5%	2.9%	4.0%	0.27%	
DOWNERS	3.9%	3.3%	4.5%	0.30%	
ROHYPNOL	1.0%	0.7%	1.3%	0.14%	
STEROIDS	0.7%	0.5%	0.9%	0.09%	
ECSTASY	3.9%	3.1%	4.8%	0.42%	
HEROIN	0.4%	0.3%	0.5%	0.06%	
METHAMPHETAMINE	0.7%	0.6%	0.9%	0.09%	
CODEINE	5.4%	4.6%	6.1%	0.37%	
OXYCONTIN ETC	1.4%	1.1%	1.6%	0.13%	
VICODIN ETC	3.1%	2.6%	3.7%	0.30%	
VALIUM ETC	1.2%	0.8%	1.6%	0.18%	
XANAX ETC	2.4%	1.9%	2.8%	0.22%	
DXM ETC	1.6%	1.3%	1.9%	0.15%	

Lifetime Use of Selected Substances - Male Overall				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	33.7%	31.7%	35.8%	1.03%
CIGARETTES	30.9%	28.9%	32.8%	0.97%
SMOKELESS TOBACCO	13.9%	12.3%	15.6%	0.82%
ALCOHOL	60.9%	58.9%	62.9%	1.00%
INHALANTS	16.7%	15.9%	17.6%	0.43%
ANY ILLICIT DRUG	30.8%	28.7%	32.9%	1.07%
MARIJUANA	29.6%	27.3%	31.8%	1.12%
MARIJUANA ONLY	15.8%	14.4%	17.1%	0.67%
COCAINE OR CRACK	6.0%	5.3%	6.7%	0.34%
COCAINE	5.7%	5.0%	6.4%	0.34%
CRACK	1.6%	1.4%	1.8%	0.08%
HALLUCINOGENS	6.0%	5.1%	6.9%	0.43%
UPPERS	5.6%	4.9%	6.3%	0.36%
DOWNERS	6.3%	5.4%	7.1%	0.43%
ROHYPNOL	2.7%	2.3%	3.1%	0.22%
STEROIDS	1.8%	1.6%	2.1%	0.13%
ECSTASY	7.2%	6.5%	7.8%	0.32%
HEROIN	1.4%	1.3%	1.6%	0.09%
METHAMPHETAMINE	3.1%	2.8%	3.3%	0.11%
CODEINE	13.2%	12.1%	14.2%	0.53%
OXYCONTIN ETC	3.6%	3.1%	4.1%	0.25%
VICODIN ETC	5.7%	4.6%	6.9%	0.58%
VALIUM ETC	2.6%	2.2%	3.0%	0.21%
XANAX ETC	4.8%	4.0%	5.6%	0.41%
DXM ETC	5.7%	5.2%	6.2%	0.25%

Lifetime Use of Selected Substances - Male Grade 7					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	16.7%	15.3%	18.2%	0.74%	
CIGARETTES	15.1%	13.8%	16.5%	0.67%	
SMOKELESS TOBACCO	5.3%	4.4%	6.3%	0.48%	
ALCOHOL	41.7%	38.9%	44.6%	1.45%	
INHALANTS	18.1%	16.9%	19.4%	0.63%	
ANY ILLICIT DRUG	12.4%	10.8%	14.1%	0.83%	
MARIJUANA	10.8%	9.1%	12.5%	0.83%	
MARIJUANA ONLY	6.3%	5.1%	7.5%	0.60%	

Past 30-day Use of Selected Substances - Male Overall				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	15.2%	14.0%	16.4%	0.61%
CIGARETTES	13.0%	12.0%	14.1%	0.52%
SMOKELESS TOBACCO	5.7%	4.9%	6.6%	0.40%
ALCOHOL	29.1%	27.8%	30.5%	0.69%
INHALANTS	5.9%	5.5%	6.4%	0.23%
ANY ILLICIT DRUG	14.9%	13.7%	16.0%	0.58%
MARIJUANA	13.2%	12.0%	14.3%	0.57%
MARIJUANA ONLY	5.4%	4.7%	6.0%	0.32%
COCAINE OR CRACK	1.9%	1.7%	2.1%	0.11%
COCAINE	1.7%	1.5%	2.0%	0.12%
CRACK	0.4%	0.4%	0.5%	0.04%
HALLUCINOGENS	2.0%	1.8%	2.3%	0.12%
UPPERS	2.5%	2.1%	2.8%	0.18%
DOWNERS	2.7%	2.4%	3.1%	0.19%
ROHYPNOL	1.1%	0.9%	1.3%	0.10%
STEROIDS	0.7%	0.5%	0.8%	0.07%
ECSTASY	2.8%	2.5%	3.1%	0.17%
HEROIN	0.5%	0.4%	0.6%	0.04%
METHAMPHETAMINE	0.9%	0.7%	1.0%	0.07%
CODEINE	5.5%	4.9%	6.2%	0.30%
OXYCONTIN ETC	1.2%	1.0%	1.4%	0.10%
VICODIN ETC	2.1%	1.6%	2.7%	0.28%
VALIUM ETC	0.9%	0.7%	1.0%	0.07%
XANAX ETC	1.9%	1.4%	2.5%	0.26%
DXM ETC	2.2%	1.9%	2.4%	0.12%

Past 30-day Use of Selected Substances - Male Grade 7					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	5.4%	4.7%	6.1%	0.36%	
CIGARETTES	4.7%	4.0%	5.3%	0.33%	
SMOKELESS TOBACCO	1.8%	1.4%	2.2%	0.20%	
ALCOHOL	14.3%	13.1%	15.5%	0.60%	
INHALANTS	7.7%	6.9%	8.6%	0.43%	
ANY ILLICIT DRUG	5.3%	4.4%	6.2%	0.45%	
MARIJUANA	4.0%	3.3%	4.8%	0.40%	
MARIJUANA ONLY	1.9%	1.4%	2.5%	0.25%	

COCAINE OR CRACK	2.2%	1.7%	2.7%	0.25%
COCAINE	1.6%	1.2%	2.0%	0.20%
CRACK	1.0%	0.8%	1.3%	0.13%
HALLUCINOGENS	1.5%	1.2%	1.9%	0.19%
UPPERS	1.0%	0.6%	1.4%	0.18%
DOWNERS	1.0%	0.7%	1.4%	0.16%
ROHYPNOL	1.4%	1.1%	1.7%	0.16%
STEROIDS	1.3%	1.0%	1.5%	0.14%
ECSTASY	2.3%	1.8%	2.9%	0.27%
HEROIN	1.3%	1.0%	1.6%	0.15%
METHAMPHETAMINE	1.7%	1.4%	2.1%	0.16%
CODEINE	6.4%	5.6%	7.2%	0.42%
OXYCONTIN ETC	0.9%	0.6%	1.1%	0.12%
VICODIN ETC	0.9%	0.6%	1.2%	0.13%
VALIUM ETC	0.8%	0.6%	1.1%	0.12%
XANAX ETC	1.1%	0.8%	1.4%	0.15%
DXM ETC	3.3%	2.7%	3.9%	0.29%

COCAINE OR CRACK	0.8%	0.6%	1.0%	0.11%
COCAINE	0.5%	0.4%	0.7%	0.08%
CRACK	0.4%	0.3%	0.6%	0.08%
HALLUCINOGENS	0.7%	0.4%	0.9%	0.12%
UPPERS	0.4%	0.3%	0.6%	0.08%
DOWNERS	0.5%	0.3%	0.6%	0.10%
ROHYPNOL	0.5%	0.3%	0.6%	0.08%
STEROIDS	0.3%	0.2%	0.4%	0.06%
ECSTASY	0.7%	0.4%	0.9%	0.13%
HEROIN	0.4%	0.2%	0.5%	0.07%
METHAMPHETAMINE	0.6%	0.4%	0.8%	0.10%
CODEINE	3.0%	2.5%	3.4%	0.22%
OXYCONTIN ETC	0.4%	0.2%	0.5%	0.08%
VICODIN ETC	0.3%	0.2%	0.5%	0.08%
VALIUM ETC	0.4%	0.2%	0.6%	0.10%
XANAX ETC	0.5%	0.3%	0.6%	0.09%
DXM ETC	1.3%	1.0%	1.7%	0.16%

Lifetime Use of Selected Substances - Male Grade 8				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	25.3%	22.9%	27.6%	1.16%
CIGARETTES	22.8%	20.9%	24.8%	1.00%
SMOKELESS TOBACCO	8.7%	7.0%	10.4%	0.84%
ALCOHOL	55.0%	52.5%	57.4%	1.23%
INHALANTS	19.1%	17.2%	21.0%	0.95%
ANY ILLICIT DRUG	20.9%	18.5%	23.2%	1.16%
MARIJUANA	19.1%	16.7%	21.4%	1.17%
MARIJUANA ONLY	10.9%	9.2%	12.5%	0.80%
COCAINE OR CRACK	3.3%	2.6%	3.9%	0.35%
COCAINE	2.9%	2.2%	3.5%	0.31%
CRACK	1.5%	1.1%	1.8%	0.18%
HALLUCINOGENS	3.9%	3.0%	4.7%	0.42%
UPPERS	2.8%	2.1%	3.4%	0.32%
DOWNERS	2.7%	1.9%	3.6%	0.42%
ROHYPNOL	2.8%	2.1%	3.4%	0.33%
STEROIDS	1.9%	1.4%	2.3%	0.22%
ECSTASY	4.6%	3.7%	5.5%	0.45%
HEROIN	1.4%	1.0%	1.9%	0.21%
METHAMPHETAMINE	2.2%	1.7%	2.8%	0.27%

Past 30-day Use of Selected Substances - Male Grade 8				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	8.5%	7.5%	9.6%	0.53%
CIGARETTES	7.1%	6.2%	8.0%	0.44%
SMOKELESS TOBACCO	2.9%	2.3%	3.6%	0.32%
ALCOHOL	21.5%	19.5%	23.5%	1.00%
INHALANTS	7.3%	6.3%	8.3%	0.50%
ANY ILLICIT DRUG	9.3%	7.9%	10.8%	0.72%
MARIJUANA	7.7%	6.5%	8.9%	0.58%
MARIJUANA ONLY	3.6%	2.8%	4.4%	0.39%
COCAINE OR CRACK	1.2%	0.9%	1.4%	0.11%
COCAINE	1.0%	0.8%	1.2%	0.10%
CRACK	0.5%	0.3%	0.6%	0.08%
HALLUCINOGENS	1.1%	0.7%	1.5%	0.19%
UPPERS	1.4%	0.8%	1.9%	0.26%
DOWNERS	1.3%	0.7%	2.0%	0.34%
ROHYPNOL	1.2%	0.8%	1.7%	0.24%
STEROIDS	0.5%	0.3%	0.7%	0.09%
ECSTASY	1.7%	1.2%	2.2%	0.24%
HEROIN	0.5%	0.3%	0.7%	0.10%
METHAMPHETAMINE	0.9%	0.6%	1.2%	0.17%

CODEINE	8.9%	7.8%	9.9%	0.53%
OXYCONTIN ETC	1.7%	1.2%	2.2%	0.24%
VICODIN ETC	2.2%	1.6%	2.8%	0.30%
VALIUM ETC	1.0%	0.6%	1.3%	0.18%
XANAX ETC	2.1%	1.4%	2.8%	0.35%
DXM ETC	4.1%	3.4%	4.8%	0.35%

Lifetime Use of Selected Substances - Male Grade 9				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	31.4%	29.3%	33.4%	1.02%
CIGARETTES	28.9%	27.1%	30.7%	0.90%
SMOKELESS TOBACCO	11.5%	10.1%	12.9%	0.70%
ALCOHOL	61.0%	58.9%	63.0%	1.03%
INHALANTS	16.4%	14.8%	18.0%	0.78%
ANY ILLICIT DRUG	29.3%	27.3%	31.3%	1.00%
MARIJUANA	28.1%	26.1%	30.0%	0.97%
MARIJUANA ONLY	15.2%	13.9%	16.6%	0.67%
COCAINE OR CRACK	5.3%	4.6%	6.1%	0.39%
COCAINE	4.9%	4.2%	5.7%	0.37%
CRACK	1.6%	1.2%	2.0%	0.20%
HALLUCINOGENS	5.7%	4.8%	6.6%	0.44%
UPPERS	5.0%	4.1%	6.0%	0.45%
DOWNERS	5.5%	4.6%	6.5%	0.47%
ROHYPNOL	3.1%	2.5%	3.8%	0.33%
STEROIDS	1.7%	1.2%	2.2%	0.24%
ECSTASY	6.6%	5.8%	7.4%	0.41%
HEROIN	1.5%	1.1%	1.8%	0.20%
METHAMPHETAMINE	2.8%	2.4%	3.1%	0.18%
CODEINE	12.4%	11.3%	13.5%	0.55%
OXYCONTIN ETC	3.1%	2.5%	3.6%	0.28%
VICODIN ETC	4.2%	3.2%	5.2%	0.50%
VALIUM ETC	2.3%	1.8%	2.7%	0.22%
XANAX ETC	4.1%	3.2%	5.0%	0.44%
DXM ETC	6.1%	5.1%	7.0%	0.48%

Lifetime Use of Selected Substances - Male Grade 10					
CATEGORY	Prevalence	Bour	Standard		
		Lower	Upper	Error	
TOBACCO	38.69	36.0%	41.1%	1.29%	

CODEINE	4.2%	3.5%	4.8%	0.31%
OXYCONTIN ETC	0.6%	0.4%	0.8%	0.09%
VICODIN ETC	0.7%	0.5%	1.0%	0.13%
VALIUM ETC	0.4%	0.2%	0.6%	0.11%
XANAX ETC	0.8%	0.4%	1.3%	0.21%
DXM ETC	1.9%	1.5%	2.4%	0.22%

Past 30-day Use o	Past 30-day Use of Selected Substances - Male Grade 9				
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	13.3%	11.9%	14.7%	0.71%	
CIGARETTES	11.1%	9.7%	12.6%	0.71%	
SMOKELESS TOBACCO	4.8%	4.0%	5.6%	0.39%	
ALCOHOL	28.2%	26.3%	30.1%	0.97%	
INHALANTS	6.4%	5.6%	7.3%	0.42%	
ANY ILLICIT DRUG	14.1%	12.9%	15.3%	0.62%	
MARIJUANA	12.8%	11.6%	13.9%	0.58%	
MARIJUANA ONLY	5.4%	4.5%	6.3%	0.45%	
COCAINE OR CRACK	2.0%	1.6%	2.5%	0.23%	
COCAINE	1.9%	1.4%	2.3%	0.23%	
CRACK	0.3%	0.2%	0.5%	0.07%	
HALLUCINOGENS	2.4%	2.0%	2.8%	0.22%	
UPPERS	2.3%	1.9%	2.7%	0.20%	
DOWNERS	2.5%	2.1%	2.9%	0.19%	
ROHYPNOL	1.3%	1.0%	1.6%	0.16%	
STEROIDS	0.6%	0.4%	0.8%	0.12%	
ECSTASY	2.9%	2.4%	3.4%	0.24%	
HEROIN	0.5%	0.3%	0.6%	0.08%	
METHAMPHETAMINE	0.7%	0.5%	1.0%	0.13%	
CODEINE	5.2%	4.5%	5.9%	0.36%	
OXYCONTIN ETC	1.1%	0.8%	1.4%	0.16%	
VICODIN ETC	1.8%	1.3%	2.3%	0.26%	
VALIUM ETC	0.7%	0.5%	0.9%	0.09%	
XANAX ETC	1.7%	1.2%	2.3%	0.27%	
DXM ETC	2.4%	2.0%	2.9%	0.23%	

Past 30-day Use of Selected Substances - Male Grade 10					
CATEGORY		Prevalence	Boundaries		Standard
			Lower	Upper	Error
TOBACCO		16.9%	15.5%	18.3%	0.68%

CICADETTEC	25.50/	22.00/	20.40/	1 200/
CIGARETTES	35.5%	32.9%	38.1%	1.28%
SMOKELESS TOBACCO	15.7%	13.7%	17.7%	0.99%
ALCOHOL	67.8%	65.4%	70.1%	1.18%
INHALANTS	16.7%	15.2%	18.1%	0.73%
ANY ILLICIT DRUG	37.7%	35.1%	40.3%	1.29%
MARIJUANA	36.9%	34.1%	39.7%	1.41%
MARIJUANA ONLY	20.1%	18.1%	22.0%	0.98%
COCAINE OR CRACK	7.0%	5.8%	8.2%	0.60%
COCAINE	6.7%	5.6%	7.9%	0.58%
CRACK	1.5%	1.1%	1.9%	0.20%
HALLUCINOGENS	7.5%	6.4%	8.6%	0.55%
UPPERS	7.6%	6.5%	8.7%	0.56%
DOWNERS	7.9%	6.9%	8.9%	0.49%
ROHYPNOL	2.7%	2.3%	3.2%	0.24%
STEROIDS	1.7%	1.3%	2.2%	0.22%
ECSTASY	8.1%	7.1%	9.2%	0.53%
HEROIN	1.5%	1.1%	1.9%	0.21%
METHAMPHETAMINE	3.2%	2.7%	3.7%	0.25%
CODEINE	15.7%	14.3%	17.1%	0.71%
OXYCONTIN ETC	4.7%	3.9%	5.4%	0.37%
VICODIN ETC	7.6%	5.7%	9.4%	0.92%
VALIUM ETC	3.4%	2.7%	4.2%	0.39%
XANAX ETC	6.4%	5.2%	7.6%	0.60%
DXM ETC	7.1%	6.2%	8.1%	0.48%

	-			_
CIGARETTES	14.3%	13.1%	15.4%	0.57%
SMOKELESS TOBACCO	7.0%	5.8%	8.1%	0.57%
ALCOHOL	32.4%	30.8%	33.9%	0.78%
INHALANTS	4.6%	4.1%	5.1%	0.25%
ANY ILLICIT DRUG	18.4%	16.8%	19.9%	0.76%
MARIJUANA	16.7%	14.9%	18.4%	0.87%
MARIJUANA ONLY	6.9%	5.9%	7.9%	0.50%
COCAINE OR CRACK	2.2%	1.8%	2.7%	0.23%
COCAINE	2.1%	1.6%	2.6%	0.23%
CRACK	0.3%	0.2%	0.5%	0.07%
HALLUCINOGENS	2.1%	1.6%	2.7%	0.28%
UPPERS	3.4%	2.6%	4.1%	0.39%
DOWNERS	3.6%	3.0%	4.2%	0.29%
ROHYPNOL	1.3%	0.9%	1.6%	0.16%
STEROIDS	0.7%	0.5%	1.0%	0.12%
ECSTASY	3.2%	2.5%	3.8%	0.32%
HEROIN	0.6%	0.3%	0.8%	0.11%
METHAMPHETAMINE	0.8%	0.5%	1.1%	0.15%
CODEINE	6.7%	5.5%	7.9%	0.61%
OXYCONTIN ETC	1.4%	1.0%	1.7%	0.18%
VICODIN ETC	2.8%	1.7%	3.9%	0.56%
VALIUM ETC	0.9%	0.7%	1.2%	0.14%
XANAX ETC	2.8%	1.9%	3.7%	0.45%
DXM ETC	2.3%	1.9%	2.7%	0.19%

Lifetime Use of Selected Substances - Male Grade 11					
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	45.1%	42.7%	47.6%	1.23%	
CIGARETTES	40.9%	38.8%	43.1%	1.07%	
SMOKELESS TOBACCO	21.7%	18.9%	24.5%	1.41%	
ALCOHOL	69.5%	67.7%	71.3%	0.89%	
INHALANTS	15.7%	13.8%	17.5%	0.92%	
ANY ILLICIT DRUG	42.8%	40.4%	45.1%	1.16%	
MARIJUANA	41.6%	39.1%	44.1%	1.24%	
MARIJUANA ONLY	21.4%	19.3%	23.5%	1.05%	
COCAINE OR CRACK	8.6%	7.5%	9.7%	0.54%	
COCAINE	8.6%	7.5%	9.7%	0.54%	
CRACK	1.8%	1.3%	2.3%	0.25%	
HALLUCINOGENS	8.5%	7.2%	9.8%	0.67%	

Past 30-day Use of Selected Substances - Male Grade 11					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	22.0%	20.4%	23.7%	0.81%	
CIGARETTES	19.0%	17.8%	20.2%	0.62%	
SMOKELESS TOBACCO	9.0%	7.3%	10.6%	0.82%	
ALCOHOL	37.5%	35.7%	39.2%	0.86%	
INHALANTS	5.1%	4.2%	5.9%	0.44%	
ANY ILLICIT DRUG	20.4%	19.2%	21.7%	0.63%	
MARIJUANA	18.1%	16.6%	19.5%	0.73%	
MARIJUANA ONLY	7.0%	5.8%	8.2%	0.62%	
COCAINE OR CRACK	2.4%	2.0%	2.9%	0.22%	
COCAINE	2.3%	1.9%	2.7%	0.22%	
CRACK	0.6%	0.4%	0.8%	0.11%	
HALLUCINOGENS	2.9%	2.2%	3.5%	0.32%	

UPPERS	8.9%	7.8%	10.1%	0.57%
DOWNERS	10.2%	8.7%	11.7%	0.77%
ROHYPNOL	3.0%	2.3%	3.8%	0.38%
STEROIDS	2.1%	1.6%	2.7%	0.26%
ECSTASY	10.4%	8.9%	12.0%	0.77%
HEROIN	1.4%	1.0%	1.7%	0.18%
METHAMPHETAMINE	4.1%	3.3%	4.9%	0.41%
CODEINE	18.0%	16.4%	19.6%	0.79%
OXYCONTIN ETC	5.8%	5.0%	6.6%	0.41%
VICODIN ETC	10.1%	8.4%	11.8%	0.84%
VALIUM ETC	3.9%	3.3%	4.6%	0.31%
XANAX ETC	7.3%	6.1%	8.5%	0.58%
DXM ETC	7.3%	6.4%	8.2%	0.45%

UPPERS	3.8%	3.1%	4.4%	0.31%
DOWNERS	4.3%	3.6%	5.1%	0.38%
ROHYPNOL	1.2%	0.9%	1.5%	0.17%
STEROIDS	0.8%	0.5%	1.1%	0.16%
ECSTASY	3.9%	3.1%	4.6%	0.38%
HEROIN	0.5%	0.3%	0.7%	0.09%
METHAMPHETAMINE	1.2%	0.8%	1.6%	0.19%
CODEINE	7.1%	6.2%	7.9%	0.42%
OXYCONTIN ETC	1.8%	1.4%	2.3%	0.20%
VICODIN ETC	3.7%	2.6%	4.8%	0.55%
VALIUM ETC	1.2%	0.7%	1.7%	0.23%
XANAX ETC	3.1%	2.0%	4.1%	0.53%
DXM ETC	2.7%	2.2%	3.3%	0.28%

Lifetime Use of Selected Substances - Male Grade 12					
CATEGORY	Prevalence		daries	Standard	
		Lower	Upper	Error	
TOBACCO	49.8%	47.6%	52.1%	1.13%	
CIGARETTES	46.0%	43.9%	48.1%	1.06%	
SMOKELESS TOBACCO	23.4%	20.7%	26.2%	1.36%	
ALCOHOL	74.1%	72.2%	76.0%	0.94%	
INHALANTS	13.8%	12.3%	15.3%	0.75%	
ANY ILLICIT DRUG	46.1%	43.8%	48.3%	1.12%	
MARIJUANA	45.4%	43.0%	47.7%	1.18%	
MARIJUANA ONLY	22.9%	20.8%	25.0%	1.04%	
COCAINE OR CRACK	10.9%	9.5%	12.3%	0.70%	
COCAINE	10.7%	9.2%	12.2%	0.73%	
CRACK	2.2%	1.8%	2.6%	0.21%	
HALLUCINOGENS	9.8%	8.6%	11.1%	0.64%	
UPPERS	9.5%	8.2%	10.7%	0.64%	
DOWNERS	11.7%	10.2%	13.2%	0.74%	
ROHYPNOL	3.3%	2.2%	4.3%	0.51%	
STEROIDS	2.6%	2.1%	3.1%	0.25%	
ECSTASY	12.1%	10.8%	13.5%	0.68%	
HEROIN	1.6%	1.2%	2.0%	0.19%	
METHAMPHETAMINE	4.7%	4.2%	5.2%	0.26%	
CODEINE	19.2%	17.5%	21.0%	0.88%	
OXYCONTIN ETC	6.3%	5.5%	7.0%	0.40%	
VICODIN ETC	10.8%	9.2%	12.4%	0.79%	
VALIUM ETC	4.8%	4.1%	5.5%	0.35%	

Past 30-day Use of Selected Substances - Male Grade 12					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	27.8%	25.4%	30.2%	1.19%	
CIGARETTES	24.7%	22.5%	26.8%	1.07%	
SMOKELESS TOBACCO	10.3%	8.7%	12.0%	0.83%	
ALCOHOL	44.6%	42.7%	46.6%	0.97%	
INHALANTS	3.8%	3.1%	4.5%	0.36%	
ANY ILLICIT DRUG	24.0%	22.1%	25.9%	0.95%	
MARIJUANA	21.8%	19.8%	23.7%	0.96%	
MARIJUANA ONLY	8.0%	6.8%	9.1%	0.56%	
COCAINE OR CRACK	3.0%	2.5%	3.6%	0.28%	
COCAINE	2.9%	2.4%	3.5%	0.29%	
CRACK	0.5%	0.3%	0.6%	0.08%	
HALLUCINOGENS	3.1%	2.5%	3.7%	0.30%	
UPPERS	4.1%	3.5%	4.8%	0.30%	
DOWNERS	5.0%	4.2%	5.7%	0.37%	
ROHYPNOL	1.3%	0.9%	1.7%	0.22%	
STEROIDS	1.1%	0.8%	1.4%	0.14%	
ECSTASY	5.0%	4.0%	6.0%	0.48%	
HEROIN	0.7%	0.5%	1.0%	0.11%	
METHAMPHETAMINE	1.1%	0.8%	1.4%	0.14%	
CODEINE	7.7%	6.5%	9.0%	0.63%	
OXYCONTIN ETC	2.0%	1.5%	2.4%	0.24%	
VICODIN ETC	3.9%	3.2%	4.6%	0.34%	
VALIUM ETC	1.7%	1.2%	2.1%	0.21%	

XANAX ETC	8.5%	7.4%	9.6%	0.55%	
DXM ETC	6.7%	5.7%	7.7%	0.48%	

XANAX ETC	3.0%	2.4%	3.5%	0.28%
DXM FTC	2.2%	1.7%	2.7%	0.25%

Lifetime Use of Selected Substances - Female Overall				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	27.4%	25.5%	29.4%	0.96%
CIGARETTES	26.7%	24.8%	28.5%	0.93%
SMOKELESS TOBACCO	5.4%	4.6%	6.2%	0.40%
ALCOHOL	62.7%	60.2%	65.2%	1.25%
INHALANTS	17.6%	16.6%	18.7%	0.53%
ANY ILLICIT DRUG	25.2%	23.3%	27.2%	0.97%
MARIJUANA	23.2%	21.1%	25.3%	1.05%
MARIJUANA ONLY	12.2%	11.0%	13.5%	0.61%
COCAINE OR CRACK	4.8%	4.2%	5.4%	0.29%
COCAINE	4.3%	3.7%	4.9%	0.29%
CRACK	1.6%	1.5%	1.8%	0.09%
HALLUCINOGENS	3.4%	2.9%	3.8%	0.22%
UPPERS	4.9%	4.2%	5.7%	0.39%
DOWNERS	5.1%	4.3%	6.0%	0.41%
ROHYPNOL	2.6%	2.0%	3.1%	0.28%
STEROIDS	1.0%	0.9%	1.1%	0.06%
ECSTASY	6.5%	5.7%	7.3%	0.40%
HEROIN	1.4%	1.2%	1.6%	0.08%
METHAMPHETAMINE	2.9%	2.7%	3.2%	0.13%
CODEINE	11.5%	10.5%	12.6%	0.54%
OXYCONTIN ETC	2.6%	2.2%	3.1%	0.22%
VICODIN ETC	4.8%	3.8%	5.9%	0.52%
VALIUM ETC	2.1%	1.8%	2.4%	0.16%
XANAX ETC	4.3%	3.4%	5.3%	0.48%
DXM ETC	5.0%	4.6%	5.5%	0.23%

Lifetime Use of Selected Substances - Female Grade 7					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	13.6%	12.0%	15.2%	0.79%	
CIGARETTES	13.2%	11.7%	14.7%	0.77%	
SMOKELESS TOBACCO	2.6%	2.2%	3.1%	0.23%	
ALCOHOL	42.5%	39.5%	45.5%	1.50%	
INHALANTS	20.5%	18.8%	22.1%	0.84%	
ANY ILLICIT DRUG	10.7%	9.2%	12.2%	0.75%	
MARIJUANA	8.3%	6.9%	9.6%	0.67%	
MARIJUANA ONLY	4.1%	3.2%	4.9%	0.42%	
COCAINE OR CRACK	2.7%	2.2%	3.3%	0.28%	
COCAINE	2.0%	1.6%	2.4%	0.20%	

Past 30-day Use of Selected Substances - Female Overall					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	10.0%	9.0%	11.1%	0.52%	
CIGARETTES	9.6%	8.5%	10.6%	0.52%	
SMOKELESS TOBACCO	1.8%	1.5%	2.0%	0.14%	
ALCOHOL	28.9%	27.1%	30.8%	0.91%	
INHALANTS	6.1%	5.7%	6.6%	0.21%	
ANY ILLICIT DRUG	11.5%	10.4%	12.6%	0.55%	
MARIJUANA	9.8%	8.6%	10.9%	0.57%	
MARIJUANA ONLY	4.1%	3.5%	4.7%	0.29%	
COCAINE OR CRACK	1.4%	1.2%	1.6%	0.10%	
COCAINE	1.3%	1.1%	1.4%	0.10%	
CRACK	0.5%	0.4%	0.6%	0.04%	
HALLUCINOGENS	1.0%	0.9%	1.2%	0.08%	
UPPERS	1.9%	1.6%	2.2%	0.15%	
DOWNERS	2.1%	1.7%	2.4%	0.19%	
ROHYPNOL	1.0%	0.7%	1.2%	0.12%	
STEROIDS	0.3%	0.2%	0.4%	0.04%	
ECSTASY	2.3%	1.9%	2.7%	0.21%	
HEROIN	0.5%	0.5%	0.6%	0.04%	
METHAMPHETAMINE	0.7%	0.6%	0.8%	0.04%	
CODEINE	4.2%	3.7%	4.6%	0.22%	
OXYCONTIN ETC	0.8%	0.7%	0.9%	0.07%	
VICODIN ETC	1.5%	1.2%	1.9%	0.17%	
VALIUM ETC	0.6%	0.5%	0.8%	0.07%	
XANAX ETC	1.5%	1.1%	1.9%	0.21%	
DXM ETC	1.7%	1.5%	1.9%	0.10%	

Past 30-day Use of Selected Substances - Female Grade 7					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	4.2%	3.5%	4.8%	0.32%	
CIGARETTES	3.9%	3.3%	4.5%	0.30%	
SMOKELESS TOBACCO	0.9%	0.5%	1.2%	0.16%	
ALCOHOL	15.7%	13.7%	17.7%	1.01%	
INHALANTS	9.1%	8.3%	9.9%	0.40%	
ANY ILLICIT DRUG	5.0%	4.2%	5.9%	0.42%	
MARIJUANA	3.7%	2.9%	4.4%	0.38%	
MARIJUANA ONLY	1.4%	1.0%	1.8%	0.18%	
COCAINE OR CRACK	1.0%	0.8%	1.3%	0.12%	
COCAINE	0.7%	0.5%	0.9%	0.10%	

CRACK	1.9%	1.5%	2.4%	0.22%
HALLUCINOGENS	1.5%	1.2%	1.7%	0.12%
UPPERS	1.5%	1.1%	1.8%	0.17%
DOWNERS	1.4%	1.1%	1.8%	0.17%
ROHYPNOL	1.4%	1.0%	1.8%	0.20%
STEROIDS	1.3%	1.0%	1.5%	0.14%
ECSTASY	2.2%	1.8%	2.6%	0.21%
HEROIN	1.6%	1.2%	1.9%	0.17%
METHAMPHETAMINE	2.4%	2.0%	2.8%	0.20%
CODEINE	8.7%	7.6%	9.8%	0.55%
OXYCONTIN ETC	1.0%	0.8%	1.2%	0.11%
VICODIN ETC	1.2%	0.9%	1.6%	0.19%
VALIUM ETC	1.0%	0.7%	1.2%	0.13%
XANAX ETC	1.4%	1.0%	1.7%	0.18%
DXM ETC	4.4%	3.8%	5.1%	0.33%

		_		_
CRACK	0.7%	0.5%	0.8%	0.08%
HALLUCINOGENS	0.6%	0.4%	0.8%	0.09%
UPPERS	0.6%	0.4%	0.8%	0.09%
DOWNERS	0.5%	0.4%	0.7%	0.09%
ROHYPNOL	0.5%	0.3%	0.7%	0.10%
STEROIDS	0.4%	0.3%	0.6%	0.07%
ECSTASY	1.0%	0.7%	1.2%	0.13%
HEROIN	0.6%	0.4%	0.7%	0.08%
METHAMPHETAMINE	0.7%	0.5%	0.8%	0.09%
CODEINE	3.4%	2.8%	4.0%	0.30%
OXYCONTIN ETC	0.4%	0.3%	0.5%	0.06%
VICODIN ETC	0.5%	0.2%	0.8%	0.14%
VALIUM ETC	0.4%	0.2%	0.5%	0.08%
XANAX ETC	0.7%	0.4%	0.9%	0.12%
DXM ETC	1.9%	1.4%	2.3%	0.23%

Lifetime Use of Selected Substances - Female Grade 8					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	21.3%	19.1%	23.4%	1.10%	
CIGARETTES	20.6%	18.5%	22.7%	1.05%	
SMOKELESS TOBACCO	4.0%	3.1%	4.9%	0.44%	
ALCOHOL	55.1%	51.7%	58.5%	1.71%	
INHALANTS	23.0%	21.2%	24.8%	0.90%	
ANY ILLICIT DRUG	17.1%	14.6%	19.6%	1.26%	
MARIJUANA	14.0%	11.6%	16.4%	1.20%	
MARIJUANA ONLY	7.2%	6.0%	8.4%	0.61%	
COCAINE OR CRACK	3.4%	2.7%	4.1%	0.34%	
COCAINE	2.6%	2.0%	3.2%	0.30%	
CRACK	2.0%	1.6%	2.4%	0.20%	
HALLUCINOGENS	2.5%	1.9%	3.0%	0.27%	
UPPERS	2.5%	2.0%	3.0%	0.26%	
DOWNERS	2.8%	2.2%	3.5%	0.32%	
ROHYPNOL	2.4%	1.8%	3.1%	0.32%	
STEROIDS	1.3%	0.9%	1.6%	0.16%	
ECSTASY	4.4%	3.5%	5.4%	0.49%	
HEROIN	2.0%	1.6%	2.5%	0.23%	
METHAMPHETAMINE	3.0%	2.3%	3.7%	0.34%	
CODEINE	10.1%	8.8%	11.4%	0.65%	
OXYCONTIN ETC	1.9%	1.4%	2.3%	0.23%	
VICODIN ETC	2.0%	1.5%	2.5%	0.25%	
VALIUM ETC	1.6%	1.2%	2.0%	0.21%	

Past 30-day Use of Selected Substances - Female Grade 8					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	6.5%	5.4%	7.6%	0.53%	
CIGARETTES	6.0%	5.1%	6.9%	0.46%	
SMOKELESS TOBACCO	1.6%	1.2%	1.9%	0.19%	
ALCOHOL	21.9%	19.3%	24.5%	1.28%	
INHALANTS	9.5%	8.5%	10.5%	0.49%	
ANY ILLICIT DRUG	8.5%	7.0%	9.9%	0.73%	
MARIJUANA	6.0%	4.7%	7.2%	0.62%	
MARIJUANA ONLY	2.3%	1.9%	2.8%	0.23%	
COCAINE OR CRACK	1.4%	1.0%	1.7%	0.17%	
COCAINE	1.1%	0.7%	1.4%	0.16%	
CRACK	0.7%	0.5%	1.0%	0.10%	
HALLUCINOGENS	1.0%	0.7%	1.3%	0.16%	
UPPERS	1.0%	0.7%	1.4%	0.16%	
DOWNERS	1.2%	0.8%	1.7%	0.24%	
ROHYPNOL	1.1%	0.7%	1.5%	0.19%	
STEROIDS	0.4%	0.2%	0.5%	0.08%	
ECSTASY	1.9%	1.3%	2.5%	0.29%	
HEROIN	1.0%	0.7%	1.3%	0.15%	
METHAMPHETAMINE	1.0%	0.5%	1.4%	0.21%	
CODEINE	4.6%	4.0%	5.2%	0.30%	
OXYCONTIN ETC	0.6%	0.3%	0.8%	0.12%	
VICODIN ETC	0.7%	0.4%	1.0%	0.14%	
VALIUM ETC	0.7%	0.5%	0.9%	0.12%	

XANAX ETC	2.3%	1.3%	3.2%	0.48%
DXM ETC	5.6%	4.4%	6.8%	0.59%

Lifetime Use of Selected Substances - Female Grade 9				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	26.7%	24.4%	29.0%	1.16%
CIGARETTES	25.9%	23.6%	28.1%	1.12%
SMOKELESS TOBACCO	5.3%	4.2%	6.5%	0.57%
ALCOHOL	64.6%	62.0%	67.2%	1.30%
INHALANTS	18.9%	16.9%	20.9%	1.00%
ANY ILLICIT DRUG	24.3%	22.2%	26.5%	1.07%
MARIJUANA	22.6%	20.5%	24.7%	1.05%
MARIJUANA ONLY	11.5%	10.0%	13.0%	0.74%
COCAINE OR CRACK	4.2%	3.4%	5.0%	0.41%
COCAINE	3.7%	3.0%	4.5%	0.38%
CRACK	1.6%	1.3%	2.0%	0.17%
HALLUCINOGENS	3.5%	2.8%	4.2%	0.36%
UPPERS	4.4%	3.6%	5.3%	0.42%
DOWNERS	4.8%	3.9%	5.6%	0.41%
ROHYPNOL	2.8%	2.2%	3.5%	0.30%
STEROIDS	1.0%	0.7%	1.2%	0.10%
ECSTASY	6.7%	5.5%	7.9%	0.61%
HEROIN	1.6%	1.2%	2.0%	0.18%
METHAMPHETAMINE	3.1%	2.5%	3.6%	0.27%
CODEINE	12.2%	11.2%	13.2%	0.50%
OXYCONTIN ETC	2.7%	2.1%	3.3%	0.28%
VICODIN ETC	4.1%	3.2%	5.0%	0.46%
VALIUM ETC	1.9%	1.4%	2.4%	0.24%
XANAX ETC	4.0%	2.8%	5.1%	0.59%
DXM ETC	5.4%	4.5%	6.2%	0.41%

Lifetime Use of Selected Substances - Female Grade 10					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	31.8%	29.4%	34.1%	1.17%	
CIGARETTES	30.5%	28.4%	32.6%	1.05%	
SMOKELESS TOBACCO	7.5%	5.6%	9.4%	0.94%	
ALCOHOL	70.7%	68.6%	72.8%	1.04%	
INHALANTS	16.4%	14.4%	18.4%	0.98%	
ANY ILLICIT DRUG	30.2%	27.1%	33.4%	1.58%	
MARIJUANA	28.2%	25.1%	31.4%	1.57%	

XANAX ETC	1.1%	0.6%	1.6%	0.25%
DXM ETC	2.3%	1.7%	2.9%	0.31%

Past 30-day Use of Selected Substances - Female Grade 9				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	8.9%	7.4%	10.3%	0.72%
CIGARETTES	8.4%	7.1%	9.8%	0.66%
SMOKELESS TOBACCO	1.7%	1.2%	2.2%	0.24%
ALCOHOL	29.2%	27.2%	31.3%	1.01%
INHALANTS	6.4%	5.5%	7.2%	0.40%
ANY ILLICIT DRUG	11.2%	9.8%	12.5%	0.67%
MARIJUANA	9.7%	8.3%	11.1%	0.69%
MARIJUANA ONLY	3.8%	3.1%	4.4%	0.33%
COCAINE OR CRACK	1.4%	1.1%	1.8%	0.17%
COCAINE	1.3%	0.9%	1.6%	0.17%
CRACK	0.5%	0.3%	0.7%	0.08%
HALLUCINOGENS	1.2%	1.0%	1.4%	0.11%
UPPERS	2.0%	1.7%	2.4%	0.19%
DOWNERS	2.3%	1.8%	2.8%	0.23%
ROHYPNOL	1.0%	0.7%	1.2%	0.12%
STEROIDS	0.3%	0.2%	0.5%	0.06%
ECSTASY	2.4%	1.9%	2.9%	0.26%
HEROIN	0.5%	0.3%	0.6%	0.08%
METHAMPHETAMINE	0.9%	0.6%	1.1%	0.12%
CODEINE	4.8%	4.1%	5.6%	0.37%
OXYCONTIN ETC	1.0%	0.8%	1.2%	0.10%
VICODIN ETC	1.4%	1.1%	1.8%	0.19%
VALIUM ETC	0.7%	0.5%	0.8%	0.08%
XANAX ETC	1.5%	0.9%	2.2%	0.32%
DXM ETC	1.9%	1.6%	2.3%	0.18%

Past 30-day Use of Selected Substances - Female Grade 10					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	11.3%	10.0%	12.7%	0.68%	
CIGARETTES	10.7%	9.4%	12.0%	0.66%	
SMOKELESS TOBACCO	2.4%	1.7%	3.2%	0.37%	
ALCOHOL	33.7%	31.7%	35.7%	1.00%	
INHALANTS	4.9%	4.3%	5.5%	0.31%	
ANY ILLICIT DRUG	14.2%	12.3%	16.1%	0.95%	
MARIJUANA	12.1%	10.5%	13.7%	0.81%	

MARIJUANA ONLY	14.8%	13.2%	16.5%	0.82%
COCAINE OR CRACK	5.7%	4.7%	6.7%	0.49%
COCAINE	5.2%	4.3%	6.1%	0.47%
CRACK	1.7%	1.4%	2.0%	0.16%
HALLUCINOGENS	4.0%	3.2%	4.8%	0.38%
UPPERS	6.6%	4.9%	8.2%	0.81%
DOWNERS	6.4%	4.8%	8.0%	0.79%
ROHYPNOL	3.4%	2.5%	4.2%	0.43%
STEROIDS	0.9%	0.6%	1.2%	0.16%
ECSTASY	7.8%	6.7%	9.0%	0.59%
HEROIN	1.4%	1.0%	1.8%	0.20%
METHAMPHETAMINE	3.0%	2.6%	3.4%	0.20%
CODEINE	13.8%	11.7%	15.9%	1.07%
OXYCONTIN ETC	3.4%	2.3%	4.5%	0.55%
VICODIN ETC	6.3%	4.4%	8.2%	0.93%
VALIUM ETC	2.4%	1.9%	2.9%	0.24%
XANAX ETC	5.7%	3.6%	7.8%	1.05%
DXM ETC	5.8%	4.8%	6.8%	0.50%

MARIJUANA ONLY	5.7%	4.4%	6.9%	0.61%
COCAINE OR CRACK	1.5%	1.1%	1.9%	0.19%
COCAINE	1.4%	1.0%	1.8%	0.20%
CRACK	0.4%	0.2%	0.6%	0.08%
HALLUCINOGENS	1.3%	0.8%	1.7%	0.23%
UPPERS	2.5%	1.9%	3.0%	0.27%
DOWNERS	2.6%	1.8%	3.4%	0.40%
ROHYPNOL	1.2%	0.8%	1.7%	0.22%
STEROIDS	0.1%	0.1%	0.2%	0.05%
ECSTASY	2.7%	1.9%	3.6%	0.43%
HEROIN	0.7%	0.4%	0.9%	0.13%
METHAMPHETAMINE	0.7%	0.5%	0.9%	0.09%
CODEINE	5.0%	3.6%	6.4%	0.71%
OXYCONTIN ETC	1.1%	0.8%	1.5%	0.18%
VICODIN ETC	2.1%	1.6%	2.7%	0.29%
VALIUM ETC	0.6%	0.4%	0.8%	0.11%
XANAX ETC	2.1%	0.8%	3.5%	0.66%
DXM ETC	2.0%	1.5%	2.4%	0.22%

Lifetime Use of Selected Substances - Female Grade 11					
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	34.6%	32.8%	36.5%	0.93%	
CIGARETTES	33.7%	31.8%	35.7%	0.98%	
SMOKELESS TOBACCO	6.4%	5.7%	7.1%	0.35%	
ALCOHOL	70.8%	68.1%	73.5%	1.35%	
INHALANTS	14.0%	12.2%	15.8%	0.90%	
ANY ILLICIT DRUG	34.0%	32.0%	35.9%	0.98%	
MARIJUANA	32.3%	29.9%	34.7%	1.22%	
MARIJUANA ONLY	17.3%	15.7%	18.9%	0.80%	
COCAINE OR CRACK	6.2%	5.3%	7.1%	0.46%	
COCAINE	5.9%	5.0%	6.8%	0.46%	
CRACK	1.4%	1.1%	1.6%	0.15%	
HALLUCINOGENS	4.6%	3.9%	5.3%	0.36%	
UPPERS	7.2%	6.2%	8.3%	0.53%	
DOWNERS	8.0%	6.8%	9.2%	0.60%	
ROHYPNOL	3.0%	1.9%	4.0%	0.52%	
STEROIDS	0.9%	0.6%	1.1%	0.13%	
ECSTASY	8.4%	7.1%	9.8%	0.69%	
HEROIN	1.0%	0.7%	1.3%	0.16%	
METHAMPHETAMINE	3.2%	2.8%	3.6%	0.21%	
CODEINE	12.5%	10.7%	14.3%	0.90%	

Past 30-day Use of Selected Substances - Female Grade 11				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	13.8%	12.4%	15.3%	0.73%
CIGARETTES	13.5%	12.0%	15.0%	0.76%
SMOKELESS TOBACCO	1.9%	1.7%	2.2%	0.15%
ALCOHOL	34.3%	32.1%	36.4%	1.07%
INHALANTS	3.3%	2.5%	4.2%	0.42%
ANY ILLICIT DRUG	15.0%	13.5%	16.6%	0.79%
MARIJUANA	13.8%	12.1%	15.4%	0.83%
MARIJUANA ONLY	5.5%	4.7%	6.3%	0.40%
COCAINE OR CRACK	1.9%	1.5%	2.4%	0.21%
COCAINE	1.8%	1.4%	2.3%	0.21%
CRACK	0.4%	0.2%	0.5%	0.07%
HALLUCINOGENS	1.2%	0.8%	1.5%	0.17%
UPPERS	2.7%	2.0%	3.3%	0.33%
DOWNERS	3.2%	2.4%	4.0%	0.39%
ROHYPNOL	1.3%	0.7%	1.9%	0.28%
STEROIDS	0.2%	0.1%	0.3%	0.05%
ECSTASY	3.2%	2.5%	3.9%	0.34%
HEROIN	0.4%	0.2%	0.6%	0.10%
METHAMPHETAMINE	0.6%	0.4%	0.8%	0.11%
CODEINE	3.8%	3.2%	4.4%	0.30%

OXYCONTIN ETC	3.3%	2.8%	3.8%	0.26%
VICODIN ETC	7.5%	6.1%	8.9%	0.68%
VALIUM ETC	2.7%	2.1%	3.4%	0.33%
XANAX ETC	6.7%	5.7%	7.7%	0.50%
DXM ETC	4.2%	3.5%	4.9%	0.36%

Lifetime Use of Selected Substances - Female Grade 12					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	40.2%	37.4%	43.0%	1.40%	
CIGARETTES	39.6%	36.9%	42.4%	1.37%	
SMOKELESS TOBACCO	6.9%	5.8%	8.0%	0.53%	
ALCOHOL	75.9%	73.5%	78.3%	1.20%	
INHALANTS	11.1%	9.6%	12.6%	0.76%	
ANY ILLICIT DRUG	38.8%	36.3%	41.3%	1.26%	
MARIJUANA	37.6%	34.9%	40.2%	1.32%	
MARIJUANA ONLY	20.8%	18.8%	22.9%	1.02%	
COCAINE OR CRACK	7.2%	6.2%	8.3%	0.53%	
COCAINE	7.1%	6.1%	8.2%	0.52%	
CRACK	1.1%	0.7%	1.4%	0.17%	
HALLUCINOGENS	4.6%	3.7%	5.5%	0.46%	
UPPERS	8.5%	7.2%	9.9%	0.70%	
DOWNERS	8.6%	7.1%	10.0%	0.72%	
ROHYPNOL	2.5%	1.8%	3.2%	0.35%	
STEROIDS	0.8%	0.6%	1.1%	0.12%	
ECSTASY	10.2%	8.4%	12.0%	0.89%	
HEROIN	0.6%	0.3%	0.8%	0.13%	
METHAMPHETAMINE	2.9%	2.2%	3.6%	0.34%	
CODEINE	12.0%	10.3%	13.7%	0.84%	
OXYCONTIN ETC	3.8%	3.0%	4.5%	0.38%	
VICODIN ETC	9.1%	7.5%	10.6%	0.77%	
VALIUM ETC	3.5%	2.6%	4.3%	0.43%	
XANAX ETC	6.7%	5.3%	8.1%	0.69%	
DXM ETC	4.7%	3.8%	5.5%	0.44%	

OXYCONTIN ETC	0.9%	0.6%	1.3%	0.17%
VICODIN ETC	2.3%	1.8%	2.8%	0.24%
VALIUM ETC	0.8%	0.4%	1.2%	0.18%
XANAX ETC	2.1%	1.6%	2.7%	0.27%
DXM ETC	1.0%	0.7%	1.3%	0.16%

Past 30-day Use of Selected Substances - Female Grade 12					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	17.5%	15.7%	19.3%	0.89%	
CIGARETTES	17.0%	15.2%	18.9%	0.93%	
SMOKELESS TOBACCO	2.1%	1.8%	2.5%	0.19%	
ALCOHOL	42.0%	39.5%	44.5%	1.27%	
INHALANTS	2.6%	2.2%	3.0%	0.22%	
ANY ILLICIT DRUG	16.5%	15.1%	18.0%	0.71%	
MARIJUANA	15.0%	13.4%	16.6%	0.80%	
MARIJUANA ONLY	6.9%	6.0%	7.8%	0.46%	
COCAINE OR CRACK	1.4%	1.0%	1.8%	0.20%	
COCAINE	1.3%	0.9%	1.7%	0.19%	
CRACK	0.3%	0.1%	0.4%	0.09%	
HALLUCINOGENS	0.9%	0.6%	1.3%	0.16%	
UPPERS	2.7%	2.1%	3.4%	0.34%	
DOWNERS	2.7%	2.0%	3.5%	0.39%	
ROHYPNOL	0.7%	0.4%	0.9%	0.12%	
STEROIDS	0.3%	0.1%	0.5%	0.10%	
ECSTASY	2.8%	1.9%	3.6%	0.43%	
HEROIN	0.1%	0.0%	0.1%	0.03%	
METHAMPHETAMINE	0.4%	0.2%	0.5%	0.07%	
CODEINE	3.1%	2.6%	3.6%	0.25%	
OXYCONTIN ETC	0.8%	0.6%	0.9%	0.09%	
VICODIN ETC	2.4%	1.8%	3.0%	0.29%	
VALIUM ETC	0.7%	0.3%	1.2%	0.22%	
XANAX ETC	1.8%	1.3%	2.2%	0.22%	
DXM ETC	1.0%	0.7%	1.3%	0.14%	

Lifetime Use of Selected Substances - White Overall				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	30.3%	27.8%	32.7%	1.21%
CIGARETTES	27.3%	24.9%	29.7%	1.18%
SMOKELESS TOBACCO	13.0%	11.6%	14.3%	0.68%
ALCOHOL	58.5%	55.3%	61.7%	1.59%
INHALANTS	16.2%	15.1%	17.3%	0.55%
ANY ILLICIT DRUG	23.6%	21.8%	25.5%	0.92%
MARIJUANA	22.4%	20.3%	24.5%	1.07%
MARIJUANA ONLY	11.5%	10.6%	12.4%	0.46%
COCAINE OR CRACK	3.6%	2.9%	4.2%	0.33%
COCAINE	3.2%	2.5%	3.9%	0.33%
CRACK	1.2%	0.9%	1.4%	0.11%
HALLUCINOGENS	5.5%	4.7%	6.2%	0.37%
UPPERS	6.2%	5.3%	7.1%	0.44%
DOWNERS	6.1%	5.2%	7.0%	0.45%
ROHYPNOL	1.6%	1.2%	1.9%	0.17%
STEROIDS	1.7%	1.4%	1.9%	0.13%
ECSTASY	5.8%	5.0%	6.5%	0.36%
HEROIN	1.2%	1.0%	1.4%	0.10%
METHAMPHETAMINE	3.1%	2.6%	3.5%	0.21%
CODEINE	11.3%	10.1%	12.5%	0.60%
OXYCONTIN ETC	4.0%	3.4%	4.7%	0.31%
VICODIN ETC	7.5%	6.3%	8.7%	0.60%
VALIUM ETC	3.2%	2.7%	3.8%	0.27%
XANAX ETC	5.5%	4.8%	6.1%	0.33%
DXM ETC	5.2%	4.7%	5.6%	0.23%

Lifetime Use of Selected Substances - White Grade 7						
CATEGORY	Prevalence	Boundaries		Standard		
		Lower	Upper	Error		
TOBACCO	12.0%	9.6%	14.4%	1.19%		
CIGARETTES	10.6%	8.5%	12.7%	1.04%		
SMOKELESS TOBACCO	4.0%	3.0%	4.9%	0.47%		
ALCOHOL	34.0%	29.9%	38.1%	2.05%		
INHALANTS	15.6%	13.9%	17.3%	0.87%		
ANY ILLICIT DRUG	6.6%	5.4%	7.7%	0.57%		
MARIJUANA	5.2%	4.1%	6.2%	0.51%		
MARIJUANA ONLY	2.7%	2.0%	3.4%	0.35%		
COCAINE OR CRACK	1.1%	0.6%	1.7%	0.26%		
COCAINE	0.5%	0.3%	0.8%	0.14%		

				1
Past 30-day Use of Selected Substances - White Overall				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	14.1%	12.9%	15.4%	0.64%
CIGARETTES	12.1%	11.0%	13.2%	0.56%
SMOKELESS TOBACCO	5.3%	4.6%	5.9%	0.32%
ALCOHOL	26.4%	24.4%	28.3%	0.96%
INHALANTS	5.2%	4.7%	5.8%	0.28%
ANY ILLICIT DRUG	11.0%	9.6%	12.5%	0.71%
MARIJUANA	9.9%	8.4%	11.4%	0.74%
MARIJUANA ONLY	3.7%	3.0%	4.5%	0.36%
COCAINE OR CRACK	0.9%	0.7%	1.1%	0.10%
COCAINE	0.8%	0.6%	1.0%	0.09%
CRACK	0.2%	0.2%	0.3%	0.05%
HALLUCINOGENS	1.7%	1.4%	2.0%	0.13%
UPPERS	2.5%	2.1%	2.9%	0.19%
DOWNERS	2.5%	2.1%	3.0%	0.22%
ROHYPNOL	0.5%	0.4%	0.7%	0.08%
STEROIDS	0.5%	0.4%	0.7%	0.07%
ECSTASY	1.9%	1.5%	2.2%	0.18%
HEROIN	0.4%	0.3%	0.5%	0.06%
METHAMPHETAMINE	0.7%	0.5%	0.9%	0.08%
CODEINE	3.7%	3.2%	4.2%	0.25%
OXYCONTIN ETC	1.1%	0.8%	1.3%	0.12%
VICODIN ETC	2.4%	2.0%	2.8%	0.21%
VALIUM ETC	0.8%	0.6%	1.0%	0.09%
XANAX ETC	1.9%	1.6%	2.1%	0.13%
DXM ETC	1.5%	1.3%	1.7%	0.10%

Past 30-day Use of Selected Substances - White Grade 7					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	3.7%	2.7%	4.6%	0.48%	
CIGARETTES	3.1%	2.4%	3.8%	0.36%	
SMOKELESS TOBACCO	1.1%	0.5%	1.6%	0.28%	
ALCOHOL	9.5%	8.3%	10.8%	0.63%	
INHALANTS	6.3%	5.5%	7.1%	0.39%	
ANY ILLICIT DRUG	2.2%	1.6%	2.8%	0.32%	
MARIJUANA	1.6%	1.1%	2.1%	0.25%	
MARIJUANA ONLY	0.6%	0.3%	0.9%	0.16%	
COCAINE OR CRACK	0.3%	0.1%	0.5%	0.10%	
COCAINE	0.0%	0.0%	0.1%	0.02%	

CRACK	0.9%	0.4%	1.3%	0.22%
HALLUCINOGENS	1.3%	0.9%	1.7%	0.18%
UPPERS	0.7%	0.5%	1.0%	0.13%
DOWNERS	0.8%	0.5%	1.1%	0.14%
ROHYPNOL	0.7%	0.4%	0.9%	0.12%
STEROIDS	1.4%	1.0%	1.9%	0.22%
ECSTASY	1.3%	0.9%	1.8%	0.22%
HEROIN	1.0%	0.6%	1.4%	0.20%
METHAMPHETAMINE	1.5%	1.1%	1.9%	0.20%
CODEINE	7.4%	6.3%	8.5%	0.54%
OXYCONTIN ETC	1.0%	0.6%	1.3%	0.19%
VICODIN ETC	1.2%	0.7%	1.7%	0.24%
VALIUM ETC	0.9%	0.6%	1.2%	0.14%
XANAX ETC	1.4%	1.0%	1.9%	0.20%
DXM ETC	3.1%	2.5%	3.7%	0.32%

CDACK	0.20/	0.10/	0.50/	0.110/
CRACK	0.3%	0.1%	0.5%	0.11%
HALLUCINOGENS	0.4%	0.2%	0.6%	0.10%
UPPERS	0.3%	0.1%	0.4%	0.09%
DOWNERS	0.3%	0.1%	0.4%	0.08%
ROHYPNOL	0.2%	0.1%	0.3%	0.05%
STEROIDS	0.3%	0.1%	0.5%	0.09%
ECSTASY	0.4%	0.2%	0.6%	0.10%
HEROIN	0.3%	0.1%	0.5%	0.09%
METHAMPHETAMINE	0.3%	0.2%	0.5%	0.09%
CODEINE	2.8%	2.2%	3.4%	0.30%
OXYCONTIN ETC	0.2%	0.1%	0.4%	0.06%
VICODIN ETC	0.4%	0.2%	0.7%	0.13%
VALIUM ETC	0.3%	0.1%	0.5%	0.10%
XANAX ETC	0.5%	0.3%	0.7%	0.10%
DXM ETC	0.9%	0.6%	1.2%	0.14%

Lifetime Use of Selected Substances - White Grade 8				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	19.9%	16.8%	23.0%	1.55%
CIGARETTES	17.5%	15.0%	20.1%	1.29%
SMOKELESS TOBACCO	7.6%	5.6%	9.5%	0.97%
ALCOHOL	48.1%	44.6%	51.6%	1.74%
INHALANTS	18.2%	16.8%	19.6%	0.69%
ANY ILLICIT DRUG	12.2%	10.3%	14.0%	0.92%
MARIJUANA	10.0%	8.1%	11.9%	0.95%
MARIJUANA ONLY	5.7%	4.5%	6.8%	0.57%
COCAINE OR CRACK	1.6%	1.1%	2.1%	0.27%
COCAINE	1.2%	0.7%	1.6%	0.23%
CRACK	0.9%	0.6%	1.2%	0.15%
HALLUCINOGENS	3.0%	2.1%	3.8%	0.41%
UPPERS	2.2%	1.7%	2.7%	0.24%
DOWNERS	2.0%	1.4%	2.6%	0.30%
ROHYPNOL	1.1%	0.8%	1.4%	0.15%
STEROIDS	1.9%	1.3%	2.4%	0.28%
ECSTASY	2.8%	2.0%	3.5%	0.38%
HEROIN	1.1%	0.7%	1.6%	0.22%
METHAMPHETAMINE	2.0%	1.5%	2.4%	0.22%
CODEINE	7.7%	6.4%	9.0%	0.67%
OXYCONTIN ETC	1.6%	1.1%	2.2%	0.27%
VICODIN ETC	2.4%	1.6%	3.2%	0.40%
VALIUM ETC	1.2%	0.7%	1.6%	0.23%

Past 30-day Use of Selected Substances - White Grade 8					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	6.3%	4.9%	7.7%	0.68%	
CIGARETTES	5.2%	4.2%	6.2%	0.51%	
SMOKELESS TOBACCO	2.2%	1.5%	3.0%	0.37%	
ALCOHOL	14.9%	12.5%	17.4%	1.24%	
INHALANTS	7.1%	5.9%	8.3%	0.60%	
ANY ILLICIT DRUG	5.0%	4.1%	5.9%	0.44%	
MARIJUANA	3.8%	2.9%	4.6%	0.42%	
MARIJUANA ONLY	1.6%	1.1%	2.0%	0.22%	
COCAINE OR CRACK	0.4%	0.2%	0.6%	0.11%	
COCAINE	0.3%	0.1%	0.5%	0.09%	
CRACK	0.2%	0.1%	0.3%	0.05%	
HALLUCINOGENS	0.7%	0.4%	1.0%	0.15%	
UPPERS	0.7%	0.5%	1.0%	0.12%	
DOWNERS	0.8%	0.4%	1.1%	0.18%	
ROHYPNOL	0.3%	0.1%	0.5%	0.09%	
STEROIDS	0.4%	0.2%	0.5%	0.08%	
ECSTASY	1.0%	0.5%	1.4%	0.24%	
HEROIN	0.2%	0.1%	0.4%	0.06%	
METHAMPHETAMINE	0.4%	0.2%	0.6%	0.12%	
CODEINE	3.2%	2.6%	3.8%	0.30%	
OXYCONTIN ETC	0.3%	0.1%	0.5%	0.09%	
VICODIN ETC	1.0%	0.6%	1.4%	0.20%	
VALIUM ETC	0.4%	0.2%	0.6%	0.11%	

XANAX ETC	1.8%	1.1%	2.4%	0.34%
DXM ETC	3.2%	2.5%	3.9%	0.34%

Lifetime Use of Selected Substances - White Grade 9					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	26.6%	23.6%	29.6%	1.50%	
CIGARETTES	23.9%	21.2%	26.7%	1.37%	
SMOKELESS TOBACCO	10.6%	9.1%	12.0%	0.72%	
ALCOHOL	57.9%	54.2%	61.5%	1.85%	
INHALANTS	16.8%	15.0%	18.6%	0.90%	
ANY ILLICIT DRUG	20.3%	18.1%	22.4%	1.08%	
MARIJUANA	19.4%	17.1%	21.7%	1.15%	
MARIJUANA ONLY	10.1%	8.7%	11.5%	0.72%	
COCAINE OR CRACK	2.3%	1.8%	2.9%	0.26%	
COCAINE	2.1%	1.6%	2.6%	0.25%	
CRACK	0.9%	0.5%	1.2%	0.17%	
HALLUCINOGENS	4.7%	3.8%	5.7%	0.47%	
UPPERS	4.8%	4.0%	5.6%	0.41%	
DOWNERS	4.6%	3.7%	5.5%	0.45%	
ROHYPNOL	1.4%	1.1%	1.8%	0.18%	
STEROIDS	1.3%	0.9%	1.7%	0.21%	
ECSTASY	4.5%	3.6%	5.5%	0.48%	
HEROIN	1.2%	0.7%	1.6%	0.23%	
METHAMPHETAMINE	2.8%	2.0%	3.6%	0.40%	
CODEINE	10.4%	8.9%	12.0%	0.76%	
OXYCONTIN ETC	3.6%	2.9%	4.3%	0.37%	
VICODIN ETC	5.9%	4.8%	7.0%	0.53%	
VALIUM ETC	2.3%	1.7%	3.0%	0.32%	
XANAX ETC	4.4%	3.5%	5.3%	0.43%	
DXM ETC	5.1%	4.5%	5.7%	0.31%	

Lifetime Use of Selected Substances - White Grade 10						
CATEGORY	Prevalence	Boundaries		Standard		
		Lower	Upper	Error		
TOBACCO	35.8%	33.2%	38.4%	1.30%		
CIGARETTES	32.3%	29.5%	35.1%	1.39%		
SMOKELESS TOBACCO	15.9%	14.1%	17.6%	0.88%		
ALCOHOL	67.8%	65.7%	69.9%	1.07%		
INHALANTS	17.2%	15.3%	19.0%	0.91%		
ANY ILLICIT DRUG	28.9%	26.8%	31.0%	1.05%		
MARIJUANA	27.7%	25.5%	30.0%	1.13%		

XANAX ETC	0.8%	0.4%	1.2%	0.21%
DXM ETC	1.4%	1.0%	1.8%	0.20%

Past 30-day Use of Selected Substances - White Grade 9				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	11.7%	9.9%	13.4%	0.87%
CIGARETTES	9.6%	7.9%	11.3%	0.84%
SMOKELESS TOBACCO	4.2%	3.5%	4.9%	0.37%
ALCOHOL	24.2%	21.7%	26.8%	1.29%
INHALANTS	5.9%	4.9%	6.9%	0.49%
ANY ILLICIT DRUG	9.7%	8.1%	11.3%	0.79%
MARIJUANA	9.1%	7.4%	10.7%	0.80%
MARIJUANA ONLY	3.5%	2.7%	4.3%	0.38%
COCAINE OR CRACK	0.9%	0.6%	1.2%	0.15%
COCAINE	0.8%	0.5%	1.1%	0.16%
CRACK	0.3%	0.2%	0.5%	0.08%
HALLUCINOGENS	2.0%	1.3%	2.6%	0.31%
UPPERS	2.5%	2.0%	3.0%	0.25%
DOWNERS	2.3%	1.7%	2.9%	0.31%
ROHYPNOL	0.4%	0.2%	0.7%	0.12%
STEROIDS	0.5%	0.2%	0.8%	0.15%
ECSTASY	1.6%	1.1%	2.0%	0.25%
HEROIN	0.3%	0.1%	0.5%	0.09%
METHAMPHETAMINE	0.7%	0.4%	1.1%	0.18%
CODEINE	3.7%	3.1%	4.4%	0.32%
OXYCONTIN ETC	1.2%	0.8%	1.5%	0.18%
VICODIN ETC	2.2%	1.7%	2.7%	0.23%
VALIUM ETC	0.6%	0.4%	0.9%	0.12%
XANAX ETC	1.7%	1.3%	2.1%	0.20%
DXM ETC	1.5%	1.1%	1.9%	0.20%

Past 30-day Use of Selected Substances - White Grade 10					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	16.5%	14.7%	18.2%	0.88%	
CIGARETTES	13.6%	11.8%	15.3%	0.87%	
SMOKELESS TOBACCO	7.1%	5.9%	8.3%	0.59%	
ALCOHOL	31.9%	29.9%	33.9%	1.00%	
INHALANTS	4.4%	3.5%	5.2%	0.43%	
ANY ILLICIT DRUG	14.4%	12.5%	16.3%	0.96%	
MARIJUANA	13.1%	11.0%	15.1%	1.01%	

MARIJUANA ONLY	13.6%	12.1%	15.0%	0.70%
COCAINE OR CRACK	4.6%	3.2%	6.0%	0.69%
COCAINE	4.1%	2.7%	5.5%	0.69%
CRACK	1.6%	1.0%	2.2%	0.30%
HALLUCINOGENS	7.4%	6.3%	8.4%	0.51%
UPPERS	8.6%	7.3%	9.9%	0.66%
DOWNERS	7.6%	6.3%	8.8%	0.61%
ROHYPNOL	2.2%	1.7%	2.8%	0.29%
STEROIDS	1.7%	1.3%	2.1%	0.20%
ECSTASY	7.0%	5.9%	8.2%	0.58%
HEROIN	1.3%	0.8%	1.8%	0.26%
METHAMPHETAMINE	3.6%	3.0%	4.3%	0.34%
CODEINE	13.0%	11.4%	14.6%	0.80%
OXYCONTIN ETC	5.3%	4.5%	6.1%	0.41%
VICODIN ETC	9.9%	7.9%	11.9%	1.01%
VALIUM ETC	4.3%	3.3%	5.3%	0.49%
XANAX ETC	6.8%	5.8%	7.9%	0.53%
DXM ETC	6.5%	5.7%	7.3%	0.40%

MARIJUANA ONLY	4.9%	3.7%	6.1%	0.60%
COCAINE OR CRACK	1.1%	0.6%	1.7%	0.27%
COCAINE	1.0%	0.5%	1.6%	0.28%
CRACK	0.2%	0.1%	0.3%	0.05%
HALLUCINOGENS	2.4%	1.7%	3.1%	0.34%
UPPERS	3.7%	2.9%	4.5%	0.40%
DOWNERS	3.4%	2.6%	4.2%	0.39%
ROHYPNOL	0.7%	0.4%	1.1%	0.17%
STEROIDS	0.7%	0.4%	1.0%	0.14%
ECSTASY	2.4%	1.6%	3.1%	0.37%
HEROIN	0.4%	0.1%	0.6%	0.11%
METHAMPHETAMINE	1.0%	0.6%	1.4%	0.20%
CODEINE	4.5%	3.6%	5.3%	0.41%
OXYCONTIN ETC	1.6%	1.1%	2.2%	0.27%
VICODIN ETC	3.5%	2.5%	4.5%	0.49%
VALIUM ETC	1.0%	0.7%	1.4%	0.17%
XANAX ETC	2.5%	1.8%	3.1%	0.32%
DXM ETC	1.8%	1.4%	2.2%	0.19%

Lifetime Use of Selected Substances - White Grade 11					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	41.8%	39.0%	44.7%	1.42%	
CIGARETTES	37.5%	34.7%	40.2%	1.37%	
SMOKELESS TOBACCO	20.1%	17.8%	22.3%	1.13%	
ALCOHOL	68.9%	65.7%	72.1%	1.61%	
INHALANTS	16.0%	14.0%	18.1%	1.04%	
ANY ILLICIT DRUG	35.1%	32.9%	37.4%	1.11%	
MARIJUANA	34.2%	31.3%	37.1%	1.43%	
MARIJUANA ONLY	17.2%	15.5%	18.9%	0.85%	
COCAINE OR CRACK	5.3%	4.1%	6.5%	0.60%	
COCAINE	5.1%	3.9%	6.3%	0.60%	
CRACK	1.4%	1.0%	1.7%	0.16%	
HALLUCINOGENS	7.7%	6.7%	8.7%	0.51%	
UPPERS	10.2%	8.7%	11.7%	0.74%	
DOWNERS	10.3%	8.9%	11.8%	0.73%	
ROHYPNOL	2.0%	1.3%	2.6%	0.34%	
STEROIDS	2.0%	1.5%	2.6%	0.30%	
ECSTASY	8.8%	7.1%	10.5%	0.85%	
HEROIN	1.2%	0.9%	1.6%	0.17%	
METHAMPHETAMINE	4.6%	3.5%	5.8%	0.58%	
CODEINE	14.6%	12.9%	16.4%	0.89%	

Past 30-day Use of Selected Substances - White Grade 11					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	21.9%	19.6%	24.2%	1.13%	
CIGARETTES	19.2%	17.2%	21.1%	0.98%	
SMOKELESS TOBACCO	8.3%	6.8%	9.9%	0.78%	
ALCOHOL	34.9%	32.9%	36.9%	1.00%	
INHALANTS	4.3%	3.5%	5.2%	0.43%	
ANY ILLICIT DRUG	16.4%	14.3%	18.5%	1.05%	
MARIJUANA	14.8%	12.4%	17.3%	1.23%	
MARIJUANA ONLY	5.3%	3.7%	6.9%	0.80%	
COCAINE OR CRACK	1.3%	0.8%	1.7%	0.22%	
COCAINE	1.1%	0.7%	1.6%	0.21%	
CRACK	0.4%	0.2%	0.5%	0.07%	
HALLUCINOGENS	2.2%	1.7%	2.8%	0.29%	
UPPERS	3.8%	3.2%	4.4%	0.31%	
DOWNERS	4.2%	3.3%	5.1%	0.43%	
ROHYPNOL	0.9%	0.6%	1.3%	0.18%	
STEROIDS	0.5%	0.3%	0.7%	0.11%	
ECSTASY	3.0%	2.1%	3.9%	0.43%	
HEROIN	0.5%	0.3%	0.8%	0.13%	
METHAMPHETAMINE	1.1%	0.7%	1.5%	0.19%	
CODEINE	4.2%	3.1%	5.2%	0.52%	

OXYCONTIN ETC	5.9%	4.8%	6.9%	0.54%
VICODIN ETC	12.0%	10.5%	13.5%	0.76%
VALIUM ETC	4.6%	3.7%	5.5%	0.44%
XANAX ETC	8.7%	7.5%	9.8%	0.60%
DXM ETC	6.4%	5.3%	7.5%	0.55%

Lifetime Use of Selected Substances - White Grade 12					
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	47.6%	44.8%	50.5%	1.42%	
CIGARETTES	44.0%	41.4%	46.6%	1.30%	
SMOKELESS TOBACCO	21.2%	19.0%	23.4%	1.10%	
ALCOHOL	76.2%	73.3%	79.0%	1.43%	
INHALANTS	13.4%	11.4%	15.4%	1.00%	
ANY ILLICIT DRUG	40.2%	37.6%	42.7%	1.28%	
MARIJUANA	39.8%	36.9%	42.7%	1.46%	
MARIJUANA ONLY	20.7%	19.3%	22.1%	0.71%	
COCAINE OR CRACK	6.9%	5.5%	8.3%	0.72%	
COCAINE	6.7%	5.2%	8.3%	0.76%	
CRACK	1.4%	0.8%	1.9%	0.27%	
HALLUCINOGENS	9.3%	7.8%	10.7%	0.74%	
UPPERS	11.4%	9.5%	13.2%	0.91%	
DOWNERS	12.4%	10.4%	14.4%	1.00%	
ROHYPNOL	2.0%	1.1%	2.9%	0.43%	
STEROIDS	1.7%	1.2%	2.2%	0.25%	
ECSTASY	10.7%	8.9%	12.4%	0.87%	
HEROIN	1.6%	1.1%	2.0%	0.24%	
METHAMPHETAMINE	4.0%	3.2%	4.9%	0.42%	
CODEINE	15.2%	13.3%	17.1%	0.95%	
OXYCONTIN ETC	7.1%	6.1%	8.1%	0.50%	
VICODIN ETC	14.1%	12.5%	15.7%	0.81%	
VALIUM ETC	6.1%	5.2%	7.0%	0.45%	
XANAX ETC	10.0%	8.6%	11.3%	0.68%	
DXM ETC	7.0%	5.8%	8.2%	0.59%	

OXYCONTIN ETC	1.6%	1.1%	2.0%	0.22%
VICODIN ETC	3.6%	2.8%	4.3%	0.37%
VALIUM ETC	1.2%	0.7%	1.7%	0.25%
XANAX ETC	3.2%	2.5%	3.8%	0.32%
DXM ETC	1.6%	1.2%	2.1%	0.23%

Past 30-day Use of Selected Substances - White Grade 12					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	26.2%	23.7%	28.8%	1.28%	
CIGARETTES	23.6%	21.4%	25.8%	1.11%	
SMOKELESS TOBACCO	9.2%	7.9%	10.6%	0.66%	
ALCOHOL	44.7%	41.5%	47.9%	1.59%	
INHALANTS	3.1%	2.4%	3.8%	0.36%	
ANY ILLICIT DRUG	19.3%	17.0%	21.7%	1.19%	
MARIJUANA	17.8%	15.1%	20.5%	1.35%	
MARIJUANA ONLY	6.8%	5.6%	7.9%	0.56%	
COCAINE OR CRACK	1.7%	1.1%	2.2%	0.29%	
COCAINE	1.6%	1.0%	2.2%	0.30%	
CRACK	0.2%	0.1%	0.4%	0.08%	
HALLUCINOGENS	2.5%	2.0%	3.0%	0.26%	
UPPERS	4.1%	3.2%	5.1%	0.47%	
DOWNERS	4.6%	3.7%	5.6%	0.47%	
ROHYPNOL	0.8%	0.3%	1.2%	0.21%	
STEROIDS	0.7%	0.4%	1.1%	0.17%	
ECSTASY	3.2%	2.2%	4.3%	0.52%	
HEROIN	0.5%	0.2%	0.8%	0.15%	
METHAMPHETAMINE	0.6%	0.4%	0.8%	0.11%	
CODEINE	4.0%	3.0%	4.9%	0.48%	
OXYCONTIN ETC	1.6%	1.1%	2.1%	0.26%	
VICODIN ETC	3.8%	3.0%	4.6%	0.41%	
VALIUM ETC	1.2%	0.7%	1.6%	0.24%	
XANAX ETC	2.5%	2.0%	3.0%	0.26%	
DXM ETC	1.6%	1.1%	2.2%	0.29%	

Lifetime Use of Selected Substances - African American Overall				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	23.8%	20.8%	26.7%	1.47%
CIGARETTES	22.6%	19.7%	25.5%	1.45%
SMOKELESS TOBACCO	5.3%	4.4%	6.2%	0.47%
ALCOHOL	59.5%	56.9%	62.1%	1.32%
INHALANTS	13.3%	11.7%	14.9%	0.79%
ANY ILLICIT DRUG	30.7%	27.5%	33.9%	1.61%
MARIJUANA	29.8%	26.5%	33.2%	1.68%
MARIJUANA ONLY	21.7%	19.2%	24.2%	1.26%
COCAINE OR CRACK	1.7%	1.3%	2.2%	0.23%
COCAINE	1.6%	1.2%	2.1%	0.23%
CRACK	0.7%	0.5%	1.0%	0.13%
HALLUCINOGENS	2.2%	1.5%	2.8%	0.32%
UPPERS	2.6%	1.8%	3.4%	0.40%
DOWNERS	3.1%	2.3%	3.8%	0.38%
ROHYPNOL	1.1%	0.7%	1.5%	0.19%
STEROIDS	1.5%	1.2%	1.8%	0.17%
ECSTASY	4.6%	3.6%	5.7%	0.53%
HEROIN	0.9%	0.7%	1.2%	0.12%
METHAMPHETAMINE	1.8%	1.3%	2.3%	0.24%
CODEINE	14.2%	12.9%	15.4%	0.63%
OXYCONTIN ETC	1.8%	1.2%	2.4%	0.30%
VICODIN ETC	3.4%	2.5%	4.3%	0.45%
VALIUM ETC	1.3%	0.8%	1.7%	0.20%
XANAX ETC	2.5%	1.7%	3.4%	0.41%
DXM ETC	4.3%	3.5%	5.1%	0.42%

Lifetime Use of Selected Substances - African American Grade 7					
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	15.2%	12.9%	17.6%	1.17%	
CIGARETTES	14.7%	12.5%	16.9%	1.11%	
SMOKELESS TOBACCO	3.4%	2.0%	4.7%	0.67%	
ALCOHOL	44.3%	40.6%	47.9%	1.83%	
INHALANTS	18.0%	16.2%	19.9%	0.92%	
ANY ILLICIT DRUG	12.2%	9.9%	14.5%	1.15%	
MARIJUANA	10.6%	8.2%	13.0%	1.19%	
MARIJUANA ONLY	6.7%	5.6%	7.9%	0.57%	
COCAINE OR CRACK	2.1%	0.9%	3.3%	0.61%	
COCAINE	2.1%	0.9%	3.3%	0.61%	

Past 30-day Use of Selec	cted Substances - A	African Am	erican Ove	rall
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	6.9%	6.0%	7.8%	0.46%
CIGARETTES	6.2%	5.3%	7.0%	0.44%
SMOKELESS TOBACCO	1.6%	1.2%	1.9%	0.18%
ALCOHOL	25.3%	24.0%	26.5%	0.65%
INHALANTS	5.5%	4.7%	6.3%	0.40%
ANY ILLICIT DRUG	13.1%	11.3%	14.8%	0.88%
MARIJUANA	11.7%	9.8%	13.5%	0.93%
MARIJUANA ONLY	7.3%	6.1%	8.5%	0.60%
COCAINE OR CRACK	0.6%	0.4%	0.8%	0.10%
COCAINE	0.6%	0.4%	0.7%	0.09%
CRACK	0.3%	0.2%	0.5%	0.07%
HALLUCINOGENS	0.8%	0.5%	1.1%	0.15%
UPPERS	1.2%	0.7%	1.6%	0.22%
DOWNERS	1.5%	1.0%	1.9%	0.23%
ROHYPNOL	0.5%	0.3%	0.7%	0.11%
STEROIDS	0.5%	0.4%	0.7%	0.09%
ECSTASY	1.7%	1.2%	2.3%	0.28%
HEROIN	0.4%	0.3%	0.6%	0.08%
METHAMPHETAMINE	0.7%	0.5%	1.0%	0.12%
CODEINE	5.8%	5.0%	6.6%	0.40%
OXYCONTIN ETC	0.7%	0.5%	0.9%	0.11%
VICODIN ETC	1.5%	1.0%	2.0%	0.25%
VALIUM ETC	0.5%	0.3%	0.7%	0.11%
XANAX ETC	1.1%	0.7%	1.6%	0.22%
DXM ETC	1.7%	1.3%	2.1%	0.22%

Past 30-day Use of Selected Substances - African American Grade 7					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	4.8%	3.5%	6.1%	0.66%	
CIGARETTES	4.7%	3.3%	6.0%	0.68%	
SMOKELESS TOBACCO	1.4%	0.6%	2.1%	0.39%	
ALCOHOL	16.5%	13.8%	19.2%	1.36%	
INHALANTS	9.3%	7.5%	11.1%	0.89%	
ANY ILLICIT DRUG	6.0%	4.6%	7.4%	0.70%	
MARIJUANA	4.7%	3.4%	6.0%	0.64%	
MARIJUANA ONLY	2.5%	1.8%	3.1%	0.35%	
COCAINE OR CRACK	0.6%	0.1%	1.0%	0.21%	
COCAINE	0.5%	0.1%	0.9%	0.19%	

CRACK	0.6%	0.1%	1.2%	0.27%
HALLUCINOGENS	1.4%	0.6%	2.2%	0.41%
UPPERS	1.6%	0.4%	2.8%	0.60%
DOWNERS	1.2%	0.5%	1.9%	0.33%
ROHYPNOL	0.7%	0.2%	1.1%	0.24%
STEROIDS	1.4%	0.7%	2.1%	0.35%
ECSTASY	2.4%	1.2%	3.7%	0.62%
HEROIN	0.9%	0.5%	1.3%	0.20%
METHAMPHETAMINE	1.3%	0.5%	2.2%	0.41%
CODEINE	9.2%	7.7%	10.6%	0.74%
OXYCONTIN ETC	0.4%	0.1%	0.7%	0.14%
VICODIN ETC	1.1%	0.6%	1.6%	0.24%
VALIUM ETC	0.8%	0.5%	1.1%	0.17%
XANAX ETC	1.0%	0.5%	1.5%	0.25%
DXM ETC	4.6%	2.8%	6.4%	0.90%

•	•			•
CRACK	0.3%	0.0%	0.7%	0.18%
HALLUCINOGENS	0.9%	0.2%	1.7%	0.38%
UPPERS	0.7%	0.2%	1.2%	0.25%
DOWNERS	0.7%	0.2%	1.2%	0.25%
ROHYPNOL	0.3%	0.1%	0.5%	0.11%
STEROIDS	0.7%	0.2%	1.2%	0.25%
ECSTASY	0.8%	0.4%	1.2%	0.20%
HEROIN	0.3%	0.1%	0.6%	0.13%
METHAMPHETAMINE	0.5%	0.1%	1.0%	0.23%
CODEINE	4.2%	3.1%	5.2%	0.52%
OXYCONTIN ETC	0.3%	0.1%	0.6%	0.12%
VICODIN ETC	0.8%	0.3%	1.2%	0.22%
VALIUM ETC	0.5%	0.3%	0.7%	0.11%
XANAX ETC	0.7%	0.3%	1.1%	0.20%
DXM ETC	2.2%	1.0%	3.3%	0.58%

Lifetime Use of Selected Substances - African American Grade 8					
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	20.4%	16.5%	24.3%	1.95%	
CIGARETTES	19.4%	15.2%	23.6%	2.11%	
SMOKELESS TOBACCO	3.8%	2.6%	4.9%	0.59%	
ALCOHOL	55.7%	52.2%	59.1%	1.71%	
INHALANTS	17.1%	14.5%	19.7%	1.30%	
ANY ILLICIT DRUG	23.0%	18.5%	27.6%	2.27%	
MARIJUANA	21.0%	16.4%	25.6%	2.29%	
MARIJUANA ONLY	15.9%	11.6%	20.2%	2.16%	
COCAINE OR CRACK	1.6%	1.0%	2.2%	0.28%	
COCAINE	1.6%	1.0%	2.1%	0.27%	
CRACK	1.1%	0.6%	1.7%	0.28%	
HALLUCINOGENS	1.8%	0.5%	3.0%	0.62%	
UPPERS	2.0%	0.6%	3.4%	0.70%	
DOWNERS	2.2%	1.0%	3.5%	0.64%	
ROHYPNOL	1.4%	0.3%	2.6%	0.58%	
STEROIDS	1.5%	0.7%	2.2%	0.36%	
ECSTASY	3.8%	2.3%	5.4%	0.79%	
HEROIN	1.5%	0.9%	2.1%	0.29%	
METHAMPHETAMINE	1.5%	0.7%	2.3%	0.39%	
CODEINE	11.4%	9.3%	13.5%	1.05%	
OXYCONTIN ETC	1.3%	0.1%	2.4%	0.57%	
VICODIN ETC	1.6%	0.6%	2.5%	0.47%	
VALIUM ETC	0.7%	0.2%	1.2%	0.24%	

Past 30-day Use of Selected Substances - African American Grade 8					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	5.1%	3.8%	6.3%	0.63%	
CIGARETTES	4.6%	3.3%	5.9%	0.65%	
SMOKELESS TOBACCO	1.3%	0.5%	2.0%	0.35%	
ALCOHOL	20.8%	17.5%	24.2%	1.67%	
INHALANTS	7.0%	5.6%	8.4%	0.72%	
ANY ILLICIT DRUG	9.5%	7.5%	11.6%	1.02%	
MARIJUANA	7.6%	5.7%	9.6%	0.97%	
MARIJUANA ONLY	4.9%	3.4%	6.4%	0.75%	
COCAINE OR CRACK	0.7%	0.4%	1.1%	0.18%	
COCAINE	0.6%	0.2%	1.0%	0.20%	
CRACK	0.6%	0.2%	1.0%	0.19%	
HALLUCINOGENS	1.2%	0.0%	2.3%	0.57%	
UPPERS	1.2%	0.1%	2.2%	0.53%	
DOWNERS	1.3%	0.2%	2.4%	0.56%	
ROHYPNOL	0.7%	0.0%	1.4%	0.34%	
STEROIDS	0.5%	0.2%	0.8%	0.17%	
ECSTASY	1.3%	0.6%	1.9%	0.33%	
HEROIN	0.6%	0.1%	1.0%	0.24%	
METHAMPHETAMINE	0.8%	0.3%	1.3%	0.25%	
CODEINE	3.9%	2.8%	5.0%	0.55%	
OXYCONTIN ETC	0.5%	0.0%	0.9%	0.22%	
VICODIN ETC	0.7%	0.3%	1.1%	0.21%	
VALIUM ETC	0.5%	0.0%	0.9%	0.22%	

XANAX ETC	1.7%	0.5%	3.0%	0.62%
DXM ETC	4.2%	2.9%	5.5%	0.64%

Lifetime Use of Selected Substances - African American Grade 9				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	22.6%	20.3%	24.9%	1.15%
CIGARETTES	21.3%	19.3%	23.3%	0.99%
SMOKELESS TOBACCO	5.0%	2.8%	7.1%	1.06%
ALCOHOL	60.4%	58.5%	62.3%	0.94%
INHALANTS	12.3%	8.4%	16.2%	1.95%
ANY ILLICIT DRUG	28.4%	23.9%	32.8%	2.22%
MARIJUANA	27.6%	23.3%	31.9%	2.15%
MARIJUANA ONLY	20.2%	17.6%	22.8%	1.32%
COCAINE OR CRACK	1.5%	0.6%	2.4%	0.44%
COCAINE	1.4%	0.5%	2.3%	0.44%
CRACK	0.6%	0.1%	1.1%	0.25%
HALLUCINOGENS	1.6%	0.6%	2.6%	0.50%
UPPERS	1.8%	0.9%	2.8%	0.47%
DOWNERS	2.1%	1.0%	3.1%	0.52%
ROHYPNOL	1.0%	0.4%	1.7%	0.31%
STEROIDS	1.4%	0.7%	2.1%	0.36%
ECSTASY	4.4%	2.3%	6.5%	1.03%
HEROIN	0.8%	0.2%	1.3%	0.28%
METHAMPHETAMINE	1.6%	0.7%	2.6%	0.48%
CODEINE	14.6%	12.8%	16.4%	0.90%
OXYCONTIN ETC	1.3%	0.5%	2.1%	0.40%
VICODIN ETC	2.1%	1.0%	3.2%	0.57%
VALIUM ETC	1.7%	0.7%	2.6%	0.48%
XANAX ETC	2.1%	1.0%	3.3%	0.60%
DXM ETC	4.9%	3.4%	6.3%	0.73%

Lifetime Use of Selected Substances - African American Grade 10					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	28.2%	24.0%	32.3%	2.08%	
CIGARETTES	26.9%	22.9%	30.9%	2.02%	
SMOKELESS TOBACCO	6.5%	4.4%	8.6%	1.04%	
ALCOHOL	65.6%	60.2%	71.0%	2.71%	
INHALANTS	12.5%	9.5%	15.5%	1.52%	
ANY ILLICIT DRUG	36.6%	32.6%	40.5%	1.99%	
MARIJUANA	36.3%	31.6%	40.9%	2.32%	

XANAX ETC	0.8%	0.2%	1.4%	0.29%
DXM ETC	2.1%	1.3%	3.0%	0.43%

Past 30-day Use of Selected Substances - African American Grade 9					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	7.4%	6.4%	8.4%	0.48%	
CIGARETTES	6.1%	5.2%	6.9%	0.44%	
SMOKELESS TOBACCO	2.1%	1.2%	2.9%	0.43%	
ALCOHOL	25.3%	23.0%	27.5%	1.13%	
INHALANTS	4.7%	2.7%	6.7%	0.99%	
ANY ILLICIT DRUG	10.9%	8.9%	12.9%	1.00%	
MARIJUANA	9.9%	7.8%	12.0%	1.06%	
MARIJUANA ONLY	6.2%	4.0%	8.4%	1.12%	
COCAINE OR CRACK	0.4%	0.0%	0.8%	0.20%	
COCAINE	0.4%	0.0%	0.8%	0.20%	
CRACK	0.1%	0.0%	0.2%	0.07%	
HALLUCINOGENS	0.7%	0.2%	1.1%	0.22%	
UPPERS	0.8%	0.3%	1.3%	0.24%	
DOWNERS	0.9%	0.5%	1.3%	0.22%	
ROHYPNOL	0.5%	0.1%	0.8%	0.19%	
STEROIDS	0.4%	0.1%	0.7%	0.15%	
ECSTASY	1.3%	0.6%	1.9%	0.32%	
HEROIN	0.3%	-0.1%	0.6%	0.16%	
METHAMPHETAMINE	0.7%	0.1%	1.3%	0.29%	
CODEINE	6.3%	5.0%	7.5%	0.62%	
OXYCONTIN ETC	0.8%	0.2%	1.3%	0.29%	
VICODIN ETC	0.7%	0.1%	1.2%	0.27%	
VALIUM ETC	0.7%	0.3%	1.1%	0.22%	
XANAX ETC	0.7%	0.2%	1.2%	0.26%	
DXM ETC	1.6%	0.5%	2.7%	0.55%	

Past 30-day Use of Selected Substances - African American Grade 10					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	6.9%	4.5%	9.4%	1.23%	
CIGARETTES	6.1%	3.9%	8.3%	1.09%	
SMOKELESS TOBACCO	1.5%	0.7%	2.3%	0.40%	
ALCOHOL	26.9%	24.0%	29.8%	1.45%	
INHALANTS	4.9%	3.0%	6.7%	0.93%	
ANY ILLICIT DRUG	15.1%	13.0%	17.2%	1.06%	
MARIJUANA	13.2%	10.2%	16.1%	1.47%	

MARIJUANA ONLY	26.0%	23.1%	28.9%	1.47%
COCAINE OR CRACK	1.2%	0.6%	1.8%	0.29%
COCAINE	1.1%	0.6%	1.7%	0.28%
CRACK	0.4%	0.1%	0.6%	0.13%
HALLUCINOGENS	2.6%	1.7%	3.6%	0.48%
UPPERS	3.3%	1.7%	4.9%	0.79%
DOWNERS	3.6%	2.0%	5.1%	0.76%
ROHYPNOL	1.3%	0.6%	2.1%	0.37%
STEROIDS	1.3%	0.7%	2.0%	0.33%
ECSTASY	4.9%	3.3%	6.6%	0.81%
HEROIN	1.0%	0.5%	1.6%	0.28%
METHAMPHETAMINE	1.8%	1.0%	2.6%	0.39%
CODEINE	17.5%	14.3%	20.6%	1.57%
OXYCONTIN ETC	2.1%	1.0%	3.2%	0.56%
VICODIN ETC	3.5%	1.8%	5.2%	0.84%
VALIUM ETC	1.1%	0.4%	1.8%	0.36%
XANAX ETC	3.3%	1.6%	5.0%	0.85%
DXM ETC	3.4%	1.6%	5.3%	0.90%

MARIJUANA ONLY	8.7%	6.0%	11.5%	1.37%
COCAINE OR CRACK	0.7%	0.2%	1.2%	0.24%
COCAINE	0.7%	0.2%	1.1%	0.24%
CRACK	0.2%	0.0%	0.4%	0.09%
HALLUCINOGENS	0.7%	0.2%	1.2%	0.26%
UPPERS	1.5%	0.7%	2.3%	0.40%
DOWNERS	1.6%	0.8%	2.4%	0.39%
ROHYPNOL	0.8%	0.3%	1.3%	0.25%
STEROIDS	0.2%	0.1%	0.4%	0.08%
ECSTASY	1.7%	0.5%	2.9%	0.58%
HEROIN	0.6%	0.2%	1.1%	0.24%
METHAMPHETAMINE	1.1%	0.3%	1.8%	0.37%
CODEINE	6.4%	4.9%	7.9%	0.75%
OXYCONTIN ETC	0.4%	0.1%	0.7%	0.15%
VICODIN ETC	1.3%	0.5%	2.1%	0.41%
VALIUM ETC	0.2%	0.0%	0.5%	0.11%
XANAX ETC	1.6%	0.0%	3.1%	0.77%
DXM ETC	1.2%	0.6%	1.9%	0.32%

Lifetime Use of Select	Lifetime Use of Selected Substances - African American Grade 11					
CATEGORY	Prevalence	Boun	daries	Standard		
		Lower	Upper	Error		
TOBACCO	28.3%	22.7%	33.8%	2.77%		
CIGARETTES	26.2%	20.5%	31.9%	2.85%		
SMOKELESS TOBACCO	7.4%	5.3%	9.5%	1.03%		
ALCOHOL	64.9%	61.8%	68.0%	1.55%		
INHALANTS	10.8%	8.2%	13.5%	1.32%		
ANY ILLICIT DRUG	41.2%	37.0%	45.3%	2.07%		
MARIJUANA	40.4%	35.9%	44.9%	2.25%		
MARIJUANA ONLY	30.4%	26.2%	34.5%	2.07%		
COCAINE OR CRACK	1.9%	1.1%	2.6%	0.37%		
COCAINE	1.6%	1.0%	2.3%	0.34%		
CRACK	0.8%	0.3%	1.3%	0.24%		
HALLUCINOGENS	2.4%	1.3%	3.4%	0.52%		
UPPERS	2.6%	1.6%	3.5%	0.49%		
DOWNERS	4.5%	2.5%	6.4%	0.99%		
ROHYPNOL	1.2%	0.6%	1.8%	0.29%		
STEROIDS	1.4%	0.5%	2.2%	0.42%		
ECSTASY	5.6%	4.1%	7.1%	0.75%		
HEROIN	0.9%	0.5%	1.3%	0.20%		
METHAMPHETAMINE	1.7%	1.0%	2.4%	0.36%		
CODEINE	14.9%	13.1%	16.8%	0.91%		

Past 30-day Use of Selected Substances - African American Grade 11				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	8.3%	6.2%	10.3%	1.04%
CIGARETTES	7.3%	5.6%	9.1%	0.88%
SMOKELESS TOBACCO	1.8%	0.9%	2.6%	0.43%
ALCOHOL	30.4%	26.5%	34.3%	1.96%
INHALANTS	4.1%	2.2%	5.9%	0.94%
ANY ILLICIT DRUG	17.2%	14.0%	20.4%	1.62%
MARIJUANA	16.4%	13.1%	19.7%	1.65%
MARIJUANA ONLY	10.0%	7.9%	12.1%	1.05%
COCAINE OR CRACK	0.7%	0.3%	1.1%	0.18%
COCAINE	0.6%	0.2%	0.9%	0.17%
CRACK	0.3%	0.0%	0.5%	0.11%
HALLUCINOGENS	0.8%	0.3%	1.3%	0.25%
UPPERS	1.2%	0.6%	1.8%	0.31%
DOWNERS	2.2%	1.1%	3.3%	0.57%
ROHYPNOL	0.6%	0.3%	0.9%	0.16%
STEROIDS	0.7%	0.2%	1.3%	0.27%
ECSTASY	2.0%	1.1%	3.0%	0.49%
HEROIN	0.5%	0.2%	0.8%	0.16%
METHAMPHETAMINE	0.3%	0.1%	0.6%	0.12%
CODEINE	6.4%	5.2%	7.7%	0.63%

OXYCONTIN ETC	2.8%	1.8%	3.8%	0.51%
VICODIN ETC	5.9%	4.0%	7.9%	0.97%
VALIUM ETC	1.8%	0.5%	3.1%	0.66%
XANAX ETC	3.2%	2.0%	4.3%	0.57%
DXM ETC	4.5%	3.2%	5.8%	0.64%

Lifetime Use of Selected Substances - African American Grade 12				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	29.3%	22.7%	35.9%	3.29%
CIGARETTES	28.2%	21.7%	34.7%	3.26%
SMOKELESS TOBACCO	6.1%	4.8%	7.5%	0.67%
ALCOHOL	67.9%	64.0%	71.9%	1.98%
INHALANTS	8.2%	5.8%	10.6%	1.20%
ANY ILLICIT DRUG	46.8%	41.3%	52.3%	2.76%
MARIJUANA	46.7%	41.2%	52.3%	2.77%
MARIJUANA ONLY	33.9%	28.0%	39.9%	2.98%
COCAINE OR CRACK	2.2%	1.1%	3.2%	0.51%
COCAINE	2.0%	1.1%	3.0%	0.48%
CRACK	0.8%	0.2%	1.4%	0.30%
HALLUCINOGENS	3.7%	2.6%	4.8%	0.56%
UPPERS	4.4%	2.9%	6.0%	0.77%
DOWNERS	5.6%	4.4%	6.8%	0.59%
ROHYPNOL	0.8%	0.1%	1.4%	0.30%
STEROIDS	2.0%	0.8%	3.2%	0.59%
ECSTASY	6.8%	4.9%	8.8%	0.99%
HEROIN	0.5%	0.0%	0.9%	0.23%
METHAMPHETAMINE	3.1%	1.9%	4.3%	0.61%
CODEINE	18.0%	14.5%	21.5%	1.75%
OXYCONTIN ETC	3.3%	1.8%	4.9%	0.77%
VICODIN ETC	7.1%	5.0%	9.2%	1.07%
VALIUM ETC	1.4%	0.7%	2.2%	0.39%
XANAX ETC	4.2%	2.8%	5.5%	0.69%
DXM ETC	4.2%	2.6%	5.8%	0.80%

OXYCONTIN ETC	0.9%	0.3%	1.5%	0.32%
VICODIN ETC	2.9%	1.4%	4.4%	0.76%
VALIUM ETC	0.7%	0.1%	1.3%	0.29%
XANAX ETC	1.4%	0.8%	2.1%	0.35%
DXM ETC	1.3%	0.6%	2.1%	0.39%

Past 30-day Use of Selected Substances - African American Grade 12				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	9.2%	7.5%	10.9%	0.86%
CIGARETTES	8.7%	6.9%	10.4%	0.88%
SMOKELESS TOBACCO	1.5%	0.9%	2.1%	0.30%
ALCOHOL	33.3%	29.4%	37.2%	1.94%
INHALANTS	2.4%	1.0%	3.9%	0.74%
ANY ILLICIT DRUG	21.8%	18.8%	24.8%	1.49%
MARIJUANA	20.2%	17.1%	23.2%	1.52%
MARIJUANA ONLY	12.7%	10.7%	14.7%	1.02%
COCAINE OR CRACK	0.8%	0.1%	1.5%	0.35%
COCAINE	0.6%	0.0%	1.3%	0.33%
CRACK	0.5%	0.0%	1.0%	0.26%
HALLUCINOGENS	0.5%	0.1%	0.9%	0.20%
UPPERS	1.9%	0.9%	2.9%	0.51%
DOWNERS	2.4%	1.2%	3.7%	0.63%
ROHYPNOL	0.2%	-0.1%	0.5%	0.16%
STEROIDS	0.7%	0.2%	1.3%	0.28%
ECSTASY	3.6%	1.7%	5.4%	0.91%
HEROIN	0.3%	-0.1%	0.7%	0.18%
METHAMPHETAMINE	1.1%	0.6%	1.6%	0.26%
CODEINE	8.0%	4.8%	11.2%	1.61%
OXYCONTIN ETC	1.6%	0.5%	2.7%	0.54%
VICODIN ETC	2.9%	1.5%	4.3%	0.68%
VALIUM ETC	0.5%	0.2%	0.9%	0.19%
XANAX ETC	1.6%	0.8%	2.5%	0.43%
DXM ETC	1.7%	1.0%	2.5%	0.39%

Lifetime Use of Selected Substances - Mexican American Overall				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	33.0%	30.8%	35.2%	1.10%
CIGARETTES	31.9%	29.8%	34.1%	1.08%
SMOKELESS TOBACCO	8.1%	7.0%	9.2%	0.55%
ALCOHOL	65.6%	63.6%	67.6%	1.00%
INHALANTS	19.1%	18.1%	20.0%	0.48%
ANY ILLICIT DRUG	30.6%	27.8%	33.3%	1.38%
MARIJUANA	28.2%	25.3%	31.1%	1.45%
MARIJUANA ONLY	13.6%	11.9%	15.3%	0.86%
COCAINE OR CRACK	8.0%	7.3%	8.6%	0.33%
COCAINE	7.4%	6.7%	8.1%	0.34%
CRACK	2.2%	2.0%	2.5%	0.11%
HALLUCINOGENS	4.6%	3.7%	5.4%	0.42%
UPPERS	5.3%	4.4%	6.2%	0.46%
DOWNERS	6.1%	5.0%	7.3%	0.58%
ROHYPNOL	3.9%	3.2%	4.7%	0.36%
STEROIDS	1.1%	1.0%	1.3%	0.07%
ECSTASY	8.2%	7.1%	9.2%	0.51%
HEROIN	1.7%	1.5%	1.9%	0.10%
METHAMPHETAMINE	3.3%	3.0%	3.5%	0.11%
CODEINE	12.6%	11.0%	14.1%	0.78%
OXYCONTIN ETC	2.6%	2.1%	3.1%	0.25%
VICODIN ETC	4.0%	2.7%	5.3%	0.65%
VALIUM ETC	2.0%	1.6%	2.3%	0.17%
XANAX ETC	4.3%	2.9%	5.7%	0.71%
DXM ETC	5.8%	5.2%	6.3%	0.27%

Lifetime Use of Selected Substances - Mexican American Grade 7						
CATEGORY	Prevalence	Boun	daries	Standard		
		Lower	Upper	Error		
TOBACCO	17.6%	16.3%	18.8%	0.63%		
CIGARETTES	16.7%	15.5%	17.9%	0.59%		
SMOKELESS TOBACCO	4.2%	3.4%	4.9%	0.37%		
ALCOHOL	47.6%	45.8%	49.5%	0.94%		
INHALANTS	22.6%	21.2%	23.9%	0.66%		
ANY ILLICIT DRUG	15.1%	13.7%	16.6%	0.75%		
MARIJUANA	12.3%	10.9%	13.8%	0.71%		
MARIJUANA ONLY	6.5%	5.5%	7.6%	0.52%		
COCAINE OR CRACK	3.6%	3.1%	4.1%	0.26%		
COCAINE	2.7%	2.3%	3.1%	0.21%		

Past 30-day Use of Selec	ted Substances - N	1exican Am	nerican Ove	erall
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	13.1%	11.5%	14.7%	0.79%
CIGARETTES	12.2%	10.7%	13.8%	0.78%
SMOKELESS TOBACCO	3.1%	2.6%	3.6%	0.23%
ALCOHOL	32.4%	30.9%	34.0%	0.78%
INHALANTS	6.8%	6.3%	7.3%	0.24%
ANY ILLICIT DRUG	14.8%	13.2%	16.3%	0.78%
MARIJUANA	12.4%	10.8%	14.0%	0.80%
MARIJUANA ONLY	4.7%	3.9%	5.5%	0.38%
COCAINE OR CRACK	2.5%	2.3%	2.8%	0.12%
COCAINE	2.3%	2.0%	2.5%	0.12%
CRACK	0.7%	0.6%	0.8%	0.05%
HALLUCINOGENS	1.5%	1.3%	1.7%	0.09%
UPPERS	2.2%	1.9%	2.6%	0.18%
DOWNERS	2.5%	2.1%	3.0%	0.24%
ROHYPNOL	1.6%	1.3%	1.9%	0.13%
STEROIDS	0.4%	0.3%	0.5%	0.04%
ECSTASY	3.3%	2.8%	3.8%	0.25%
HEROIN	0.7%	0.6%	0.8%	0.04%
METHAMPHETAMINE	0.8%	0.7%	1.0%	0.06%
CODEINE	5.4%	4.6%	6.2%	0.39%
OXYCONTIN ETC	1.0%	0.8%	1.1%	0.08%
VICODIN ETC	1.5%	0.9%	2.0%	0.27%
VALIUM ETC	0.8%	0.6%	1.0%	0.08%
XANAX ETC	1.8%	1.1%	2.5%	0.35%
DXM ETC	2.3%	2.1%	2.5%	0.12%

Past 30-day Use of Selected Substances - Mexican American Grade 7					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	5.6%	5.1%	6.2%	0.27%	
CIGARETTES	5.1%	4.5%	5.6%	0.28%	
SMOKELESS TOBACCO	1.5%	1.1%	1.9%	0.20%	
ALCOHOL	18.7%	17.4%	19.9%	0.63%	
INHALANTS	9.7%	9.1%	10.4%	0.32%	
ANY ILLICIT DRUG	7.0%	6.2%	7.8%	0.41%	
MARIJUANA	5.1%	4.4%	5.8%	0.35%	
MARIJUANA ONLY	2.1%	1.7%	2.5%	0.19%	
COCAINE OR CRACK	1.5%	1.2%	1.8%	0.14%	
COCAINE	1.1%	0.9%	1.3%	0.12%	

CRACK	2.3%	1.8%	2.8%	0.24%
HALLUCINOGENS	1.7%	1.4%	1.9%	0.13%
UPPERS	1.5%	1.2%	1.8%	0.15%
DOWNERS	1.7%	1.4%	2.0%	0.14%
ROHYPNOL	2.1%	1.7%	2.5%	0.21%
STEROIDS	1.0%	0.8%	1.3%	0.13%
ECSTASY	2.8%	2.3%	3.3%	0.25%
HEROIN	1.9%	1.6%	2.3%	0.18%
METHAMPHETAMINE	2.8%	2.3%	3.2%	0.21%
CODEINE	7.4%	6.6%	8.2%	0.39%
OXYCONTIN ETC	1.1%	0.8%	1.4%	0.13%
VICODIN ETC	1.0%	0.8%	1.2%	0.11%
VALIUM ETC	0.9%	0.7%	1.2%	0.13%
XANAX ETC	1.2%	0.9%	1.5%	0.15%
DXM ETC	4.4%	4.0%	4.8%	0.22%

i	•			_
CRACK	0.8%	0.6%	1.0%	0.11%
HALLUCINOGENS	0.7%	0.5%	0.8%	0.09%
UPPERS	0.6%	0.5%	0.8%	0.09%
DOWNERS	0.6%	0.5%	0.8%	0.08%
ROHYPNOL	0.8%	0.5%	1.0%	0.12%
STEROIDS	0.3%	0.2%	0.4%	0.04%
ECSTASY	1.2%	0.8%	1.5%	0.18%
HEROIN	0.7%	0.5%	0.8%	0.09%
METHAMPHETAMINE	0.9%	0.7%	1.1%	0.11%
CODEINE	3.3%	2.8%	3.8%	0.26%
OXYCONTIN ETC	0.5%	0.4%	0.6%	0.06%
VICODIN ETC	0.4%	0.2%	0.5%	0.08%
VALIUM ETC	0.4%	0.2%	0.6%	0.10%
XANAX ETC	0.6%	0.4%	0.8%	0.11%
DXM ETC	2.0%	1.7%	2.3%	0.16%

Lifetime Use of Selected Substances - Mexican American Grade 8					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	26.8%	25.3%	28.3%	0.76%	
CIGARETTES	25.8%	24.4%	27.1%	0.69%	
SMOKELESS TOBACCO	6.2%	5.0%	7.3%	0.57%	
ALCOHOL	60.4%	58.5%	62.3%	0.97%	
INHALANTS	24.4%	23.0%	25.9%	0.73%	
ANY ILLICIT DRUG	22.7%	20.8%	24.6%	0.94%	
MARIJUANA	19.8%	18.1%	21.6%	0.88%	
MARIJUANA ONLY	9.3%	8.3%	10.4%	0.50%	
COCAINE OR CRACK	5.2%	4.4%	5.9%	0.39%	
COCAINE	4.3%	3.6%	4.9%	0.34%	
CRACK	2.5%	2.2%	2.9%	0.18%	
HALLUCINOGENS	3.6%	3.0%	4.2%	0.29%	
UPPERS	3.1%	2.3%	3.9%	0.40%	
DOWNERS	3.6%	2.8%	4.4%	0.39%	
ROHYPNOL	4.1%	3.3%	4.8%	0.39%	
STEROIDS	1.2%	1.0%	1.3%	0.09%	
ECSTASY	5.9%	4.7%	7.0%	0.58%	
HEROIN	2.3%	1.8%	2.8%	0.25%	
METHAMPHETAMINE	3.4%	2.9%	4.0%	0.27%	
CODEINE	10.4%	9.1%	11.7%	0.65%	
OXYCONTIN ETC	2.0%	1.4%	2.6%	0.31%	
VICODIN ETC	1.9%	1.2%	2.6%	0.34%	

Past 30-day Use of Selected Substances - Mexican American Grade 8					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	9.4%	8.3%	10.4%	0.52%	
CIGARETTES	8.3%	7.4%	9.2%	0.43%	
SMOKELESS TOBACCO	2.6%	2.2%	3.1%	0.23%	
ALCOHOL	27.2%	25.7%	28.6%	0.72%	
INHALANTS	9.9%	9.0%	10.9%	0.49%	
ANY ILLICIT DRUG	11.8%	10.5%	13.1%	0.65%	
MARIJUANA	8.9%	7.9%	10.0%	0.53%	
MARIJUANA ONLY	3.4%	2.9%	4.0%	0.28%	
COCAINE OR CRACK	2.1%	1.7%	2.4%	0.17%	
COCAINE	1.7%	1.4%	2.0%	0.14%	
CRACK	0.9%	0.7%	1.1%	0.10%	
HALLUCINOGENS	1.2%	0.9%	1.5%	0.16%	
UPPERS	1.6%	1.0%	2.1%	0.27%	
DOWNERS	1.7%	1.2%	2.2%	0.26%	
ROHYPNOL	2.0%	1.5%	2.5%	0.24%	
STEROIDS	0.4%	0.3%	0.6%	0.07%	
ECSTASY	2.5%	1.8%	3.2%	0.35%	
HEROIN	1.2%	0.9%	1.5%	0.15%	
METHAMPHETAMINE	1.3%	1.0%	1.7%	0.17%	
CODEINE	5.4%	4.7%	6.2%	0.37%	
OXYCONTIN ETC	0.8%	0.5%	1.0%	0.12%	
VICODIN ETC	0.5%	0.2%	0.7%	0.12%	

VALIUM ETC	1.5%	0.8%	2.2%	0.34%
XANAX ETC	2.6%	1.4%	3.9%	0.62%
DXM ETC	6.4%	5.5%	7.3%	0.45%

Lifetime Use of Selected Substances - Mexican American Grade 9					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	32.8%	30.7%	35.0%	1.07%	
CIGARETTES	31.7%	29.6%	33.9%	1.07%	
SMOKELESS TOBACCO	7.7%	6.5%	8.8%	0.59%	
ALCOHOL	67.5%	65.4%	69.6%	1.05%	
INHALANTS	19.7%	17.8%	21.7%	0.97%	
ANY ILLICIT DRUG	30.8%	28.3%	33.4%	1.27%	
MARIJUANA	28.5%	26.0%	31.1%	1.27%	
MARIJUANA ONLY	13.5%	11.8%	15.3%	0.87%	
COCAINE OR CRACK	7.4%	6.6%	8.1%	0.38%	
COCAINE	6.7%	5.9%	7.6%	0.43%	
CRACK	2.4%	2.1%	2.7%	0.14%	
HALLUCINOGENS	5.2%	4.2%	6.2%	0.50%	
UPPERS	5.5%	4.3%	6.7%	0.59%	
DOWNERS	6.5%	5.2%	7.7%	0.61%	
ROHYPNOL	4.6%	3.9%	5.3%	0.35%	
STEROIDS	1.1%	0.8%	1.5%	0.17%	
ECSTASY	8.8%	7.6%	10.0%	0.61%	
HEROIN	2.0%	1.6%	2.4%	0.20%	
METHAMPHETAMINE	3.5%	3.1%	3.9%	0.20%	
CODEINE	13.0%	11.4%	14.5%	0.76%	
OXYCONTIN ETC	2.8%	2.2%	3.4%	0.29%	
VICODIN ETC	3.6%	2.4%	4.7%	0.59%	
VALIUM ETC	2.0%	1.4%	2.5%	0.28%	
XANAX ETC	4.2%	2.8%	5.6%	0.70%	
DXM ETC	6.3%	5.4%	7.3%	0.46%	

Lifetime Use of Selected Substances - Mexican American Grade 10					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	36.4%	33.9%	38.9%	1.25%	
CIGARETTES	35.3%	32.8%	37.8%	1.26%	
SMOKELESS TOBACCO	9.2%	7.6%	10.8%	0.81%	
ALCOHOL	72.2%	70.0%	74.4%	1.11%	
INHALANTS	17.4%	15.4%	19.3%	0.97%	

VALIUM ETC	0.7%	0.3%	1.0%	0.18%
XANAX ETC	1.1%	0.5%	1.8%	0.34%
DXM ETC	2.7%	2.0%	3.4%	0.35%

Past 30-day Use of Selected Substances - Mexican American Grade 9				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	11.7%	10.1%	13.3%	0.82%
CIGARETTES	11.0%	9.4%	12.6%	0.81%
SMOKELESS TOBACCO	2.7%	2.1%	3.3%	0.30%
ALCOHOL	33.1%	31.5%	34.7%	0.81%
INHALANTS	7.2%	6.4%	8.0%	0.38%
ANY ILLICIT DRUG	15.0%	13.2%	16.8%	0.89%
MARIJUANA	12.9%	11.1%	14.7%	0.92%
MARIJUANA ONLY	4.8%	3.9%	5.8%	0.47%
COCAINE OR CRACK	2.6%	2.2%	3.0%	0.21%
COCAINE	2.4%	2.0%	2.9%	0.23%
CRACK	0.6%	0.4%	0.8%	0.09%
HALLUCINOGENS	2.0%	1.7%	2.3%	0.16%
UPPERS	2.4%	1.8%	2.9%	0.25%
DOWNERS	3.0%	2.3%	3.6%	0.31%
ROHYPNOL	1.8%	1.5%	2.1%	0.13%
STEROIDS	0.4%	0.2%	0.6%	0.09%
ECSTASY	3.7%	3.1%	4.4%	0.31%
HEROIN	0.6%	0.4%	0.9%	0.10%
METHAMPHETAMINE	0.8%	0.6%	1.0%	0.10%
CODEINE	5.5%	4.5%	6.5%	0.49%
OXYCONTIN ETC	1.1%	0.9%	1.4%	0.15%
VICODIN ETC	1.5%	1.0%	2.1%	0.27%
VALIUM ETC	0.8%	0.6%	1.0%	0.10%
XANAX ETC	1.8%	1.1%	2.6%	0.38%
DXM ETC	2.7%	2.3%	3.1%	0.22%

Past 30-day Use of Selected Substances - Mexican American Grade 10					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	14.1%	12.1%	16.2%	1.03%	
CIGARETTES	13.3%	11.3%	15.3%	1.02%	
SMOKELESS TOBACCO	3.6%	2.6%	4.5%	0.47%	
ALCOHOL	36.2%	34.1%	38.2%	1.03%	
INHALANTS	5.0%	4.4%	5.5%	0.28%	

ANY ILLICIT DRUG	36.7%	33.1%	40.4%	1.82%
MARIJUANA	34.5%	31.0%	38.0%	1.74%
MARIJUANA ONLY	17.5%	15.4%	19.5%	1.02%
COCAINE OR CRACK	9.6%	8.3%	10.8%	0.61%
COCAINE	9.1%	8.0%	10.3%	0.57%
CRACK	2.0%	1.7%	2.4%	0.19%
HALLUCINOGENS	4.9%	3.8%	6.0%	0.56%
UPPERS	6.9%	5.2%	8.5%	0.82%
DOWNERS	7.8%	6.1%	9.4%	0.83%
ROHYPNOL	4.4%	3.3%	5.4%	0.54%
STEROIDS	0.9%	0.7%	1.1%	0.11%
ECSTASY	9.5%	8.2%	10.9%	0.68%
HEROIN	1.6%	1.3%	1.9%	0.16%
METHAMPHETAMINE	3.0%	2.5%	3.5%	0.24%
CODEINE	15.2%	12.7%	17.8%	1.27%
OXYCONTIN ETC	3.3%	2.1%	4.5%	0.60%
VICODIN ETC	5.2%	3.0%	7.5%	1.14%
VALIUM ETC	2.1%	1.6%	2.6%	0.24%
XANAX ETC	6.1%	3.7%	8.4%	1.18%
DXM ETC	7.2%	6.0%	8.4%	0.62%

	_		_	_
ANY ILLICIT DRUG	17.9%	15.7%	20.0%	1.07%
MARIJUANA	15.4%	13.2%	17.5%	1.06%
MARIJUANA ONLY	6.4%	5.1%	7.8%	0.66%
COCAINE OR CRACK	2.9%	2.3%	3.4%	0.29%
COCAINE	2.7%	2.1%	3.3%	0.28%
CRACK	0.6%	0.4%	0.8%	0.10%
HALLUCINOGENS	1.3%	0.8%	1.9%	0.26%
UPPERS	2.6%	1.9%	3.2%	0.34%
DOWNERS	3.1%	2.2%	4.0%	0.46%
ROHYPNOL	1.9%	1.3%	2.4%	0.27%
STEROIDS	0.2%	0.1%	0.4%	0.06%
ECSTASY	3.8%	3.0%	4.7%	0.44%
HEROIN	0.8%	0.5%	1.2%	0.16%
METHAMPHETAMINE	0.5%	0.3%	0.7%	0.09%
CODEINE	6.6%	4.6%	8.6%	1.02%
OXYCONTIN ETC	1.2%	0.8%	1.5%	0.19%
VICODIN ETC	1.9%	0.9%	3.0%	0.52%
VALIUM ETC	0.7%	0.5%	1.0%	0.12%
XANAX ETC	2.5%	1.0%	4.0%	0.76%
DXM ETC	2.7%	2.3%	3.2%	0.24%

Lifetime Use of Selected Substances - Mexican American Grade 11				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	42.3%	40.0%	44.6%	1.13%
CIGARETTES	41.1%	39.1%	43.1%	1.01%
SMOKELESS TOBACCO	10.7%	9.2%	12.3%	0.77%
ALCOHOL	73.6%	71.6%	75.6%	1.00%
INHALANTS	15.1%	13.9%	16.2%	0.57%
ANY ILLICIT DRUG	40.3%	36.8%	43.8%	1.76%
MARIJUANA	38.2%	33.9%	42.5%	2.15%
MARIJUANA ONLY	17.9%	15.5%	20.4%	1.24%
COCAINE OR CRACK	11.0%	10.0%	12.0%	0.51%
COCAINE	10.9%	9.9%	11.9%	0.51%
CRACK	2.0%	1.4%	2.5%	0.28%
HALLUCINOGENS	6.4%	5.0%	7.8%	0.71%
UPPERS	7.9%	6.9%	8.8%	0.48%
DOWNERS	9.4%	7.5%	11.3%	0.94%
ROHYPNOL	4.5%	3.2%	5.8%	0.65%
STEROIDS	1.0%	0.7%	1.3%	0.16%
ECSTASY	10.9%	9.3%	12.5%	0.80%
HEROIN	1.2%	0.8%	1.7%	0.23%

Past 30-day Use of Selected Substances - Mexican American Grade 11					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	17.9%	15.4%	20.4%	1.24%	
CIGARETTES	16.9%	14.5%	19.3%	1.19%	
SMOKELESS TOBACCO	4.0%	3.3%	4.8%	0.36%	
ALCOHOL	38.7%	36.9%	40.5%	0.91%	
INHALANTS	4.0%	3.3%	4.7%	0.34%	
ANY ILLICIT DRUG	19.0%	16.9%	21.1%	1.04%	
MARIJUANA	16.6%	14.2%	18.9%	1.17%	
MARIJUANA ONLY	5.8%	5.0%	6.6%	0.42%	
COCAINE OR CRACK	3.4%	2.9%	3.9%	0.24%	
COCAINE	3.3%	2.8%	3.8%	0.24%	
CRACK	0.6%	0.4%	0.8%	0.10%	
HALLUCINOGENS	2.1%	1.6%	2.7%	0.25%	
UPPERS	3.4%	2.8%	4.0%	0.31%	
DOWNERS	3.9%	3.2%	4.5%	0.34%	
ROHYPNOL	1.7%	1.2%	2.3%	0.29%	
STEROIDS	0.4%	0.2%	0.5%	0.08%	
ECSTASY	4.4%	3.7%	5.0%	0.34%	
HEROIN	0.4%	0.2%	0.6%	0.11%	

METHAMPHETAMINE	3.3%	2.8%	3.7%	0.24%
CODEINE	15.6%	13.3%	18.0%	1.19%
OXYCONTIN ETC	3.5%	3.0%	4.0%	0.25%
VICODIN ETC	6.7%	4.6%	8.8%	1.04%
VALIUM ETC	2.7%	2.1%	3.3%	0.29%
XANAX ETC	6.6%	4.3%	8.9%	1.14%
DXM ETC	5.3%	4.6%	6.1%	0.37%

Lifetime Use of Selected	Lifetime Use of Selected Substances - Mexican American Grade 12				
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	48.0%	45.3%	50.7%	1.35%	
CIGARETTES	46.7%	44.1%	49.3%	1.30%	
SMOKELESS TOBACCO	12.6%	11.2%	14.0%	0.68%	
ALCOHOL	76.8%	74.5%	79.1%	1.15%	
INHALANTS	12.7%	11.2%	14.3%	0.77%	
ANY ILLICIT DRUG	43.4%	40.4%	46.3%	1.49%	
MARIJUANA	41.6%	38.4%	44.8%	1.59%	
MARIJUANA ONLY	19.4%	16.8%	22.1%	1.31%	
COCAINE OR CRACK	13.1%	11.8%	14.4%	0.64%	
COCAINE	12.9%	11.6%	14.3%	0.66%	
CRACK	2.2%	1.7%	2.6%	0.23%	
HALLUCINOGENS	6.4%	5.3%	7.5%	0.55%	
UPPERS	8.2%	6.7%	9.8%	0.77%	
DOWNERS	9.4%	7.7%	11.2%	0.89%	
ROHYPNOL	4.2%	2.9%	5.5%	0.63%	
STEROIDS	1.6%	1.1%	2.1%	0.24%	
ECSTASY	12.7%	10.6%	14.8%	1.06%	
HEROIN	0.9%	0.7%	1.1%	0.10%	
METHAMPHETAMINE	3.7%	3.2%	4.2%	0.25%	
CODEINE	15.1%	13.2%	17.1%	0.97%	
OXYCONTIN ETC	3.5%	2.9%	4.2%	0.31%	
VICODIN ETC	6.8%	5.1%	8.4%	0.83%	
VALIUM ETC	3.0%	2.3%	3.7%	0.36%	
XANAX ETC	6.4%	4.5%	8.3%	0.95%	
DXM ETC	4.7%	3.9%	5.6%	0.41%	

METHAMPHETAMINE	0.8%	0.6%	0.9%	0.09%
CODEINE	6.3%	5.3%	7.3%	0.51%
OXYCONTIN ETC	1.3%	1.0%	1.7%	0.17%
VICODIN ETC	2.5%	1.6%	3.3%	0.41%
VALIUM ETC	0.9%	0.6%	1.3%	0.17%
XANAX ETC	2.6%	1.4%	3.7%	0.57%
DXM ETC	2.0%	1.7%	2.4%	0.19%

Past 30-day Use of Selected Substances - Mexican American Grade 12				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	23.9%	21.9%	26.0%	1.02%
CIGARETTES	22.6%	20.5%	24.6%	1.02%
SMOKELESS TOBACCO	4.9%	4.2%	5.6%	0.34%
ALCOHOL	45.5%	43.2%	47.8%	1.14%
INHALANTS	3.4%	2.8%	4.0%	0.32%
ANY ILLICIT DRUG	20.4%	18.5%	22.2%	0.93%
MARIJUANA	18.2%	16.2%	20.1%	0.98%
MARIJUANA ONLY	6.4%	5.1%	7.7%	0.65%
COCAINE OR CRACK	2.9%	2.3%	3.4%	0.29%
COCAINE	2.7%	2.2%	3.3%	0.30%
CRACK	0.4%	0.2%	0.6%	0.09%
HALLUCINOGENS	2.0%	1.6%	2.5%	0.23%
UPPERS	3.4%	2.7%	4.1%	0.35%
DOWNERS	3.4%	2.8%	4.0%	0.31%
ROHYPNOL	1.3%	1.0%	1.7%	0.17%
STEROIDS	0.6%	0.3%	0.8%	0.13%
ECSTASY	4.5%	3.6%	5.4%	0.46%
HEROIN	0.3%	0.2%	0.5%	0.07%
METHAMPHETAMINE	0.6%	0.4%	0.9%	0.12%
CODEINE	5.7%	4.7%	6.6%	0.47%
OXYCONTIN ETC	1.0%	0.7%	1.3%	0.15%
VICODIN ETC	2.4%	1.7%	3.1%	0.34%
VALIUM ETC	1.4%	0.9%	2.0%	0.28%
XANAX ETC	2.5%	1.9%	3.2%	0.34%
DXM ETC	1.4%	1.1%	1.7%	0.16%

Lifetime Use of Selected Substances - Border Overall				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	32.6%	31.1%	34.1%	0.76%
CIGARETTES	31.8%	30.4%	33.3%	0.72%
SMOKELESS TOBACCO	7.5%	6.4%	8.5%	0.53%
ALCOHOL	61.5%	58.1%	64.8%	1.67%
INHALANTS	19.0%	17.8%	20.2%	0.61%
ANY ILLICIT DRUG	28.0%	26.0%	29.9%	0.96%
MARIJUANA	24.7%	23.0%	26.4%	0.84%
MARIJUANA ONLY	9.8%	8.9%	10.7%	0.45%
COCAINE OR CRACK	7.9%	7.3%	8.5%	0.30%
COCAINE	7.5%	6.9%	8.0%	0.28%
CRACK	2.1%	1.9%	2.3%	0.11%
HALLUCINOGENS	4.7%	3.8%	5.5%	0.42%
UPPERS	6.1%	5.2%	7.0%	0.45%
DOWNERS	7.3%	6.0%	8.6%	0.64%
ROHYPNOL	5.8%	4.5%	7.1%	0.66%
STEROIDS	1.2%	0.9%	1.4%	0.12%
ECSTASY	10.7%	8.9%	12.5%	0.91%
HEROIN	1.5%	1.4%	1.7%	0.06%
METHAMPHETAMINE	3.0%	2.8%	3.3%	0.14%
CODEINE	10.5%	9.4%	11.6%	0.56%
OXYCONTIN ETC	2.1%	1.8%	2.4%	0.13%
VICODIN ETC	2.8%	2.3%	3.4%	0.28%
VALIUM ETC	1.8%	1.6%	2.0%	0.08%
XANAX ETC	3.6%	3.1%	4.2%	0.29%
DXM ETC	4.8%	4.1%	5.5%	0.35%

Lifetime Use of Selected Substances - Border Grade 7						
CATEGORY	Prevalence	Boundaries		Standard		
		Lower	Upper	Error		
TOBACCO	15.6%	14.0%	17.2%	0.81%		
CIGARETTES	15.0%	13.3%	16.6%	0.82%		
SMOKELESS TOBACCO	3.3%	2.8%	3.8%	0.24%		
ALCOHOL	43.1%	40.3%	45.8%	1.38%		
INHALANTS	21.2%	19.2%	23.2%	0.99%		
ANY ILLICIT DRUG	12.8%	11.1%	14.4%	0.84%		
MARIJUANA	9.9%	8.2%	11.6%	0.85%		
MARIJUANA ONLY	4.3%	3.4%	5.2%	0.45%		
COCAINE OR CRACK	2.9%	2.4%	3.5%	0.26%		
COCAINE	2.4%	2.0%	2.8%	0.21%		

Past 30-day Use of Selected Substances - Border Overall				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	14.0%	13.3%	14.8%	0.38%
CIGARETTES	13.3%	12.6%	14.0%	0.35%
SMOKELESS TOBACCO	3.1%	2.7%	3.5%	0.18%
ALCOHOL	30.4%	28.4%	32.3%	0.97%
INHALANTS	7.4%	6.5%	8.2%	0.41%
ANY ILLICIT DRUG	14.2%	13.3%	15.1%	0.46%
MARIJUANA	11.0%	10.2%	11.8%	0.39%
MARIJUANA ONLY	3.4%	2.9%	3.8%	0.21%
COCAINE OR CRACK	2.9%	2.7%	3.2%	0.12%
COCAINE	2.7%	2.4%	2.9%	0.13%
CRACK	0.7%	0.6%	0.9%	0.07%
HALLUCINOGENS	1.7%	1.5%	1.9%	0.11%
UPPERS	2.8%	2.3%	3.2%	0.22%
DOWNERS	3.2%	2.5%	3.8%	0.32%
ROHYPNOL	2.3%	1.8%	2.8%	0.26%
STEROIDS	0.5%	0.3%	0.6%	0.07%
ECSTASY	4.9%	4.1%	5.6%	0.37%
HEROIN	0.7%	0.6%	0.8%	0.04%
METHAMPHETAMINE	1.0%	0.8%	1.1%	0.08%
CODEINE	4.7%	4.3%	5.2%	0.24%
OXYCONTIN ETC	0.8%	0.7%	0.9%	0.05%
VICODIN ETC	1.1%	0.9%	1.4%	0.13%
VALIUM ETC	0.8%	0.5%	1.1%	0.14%
XANAX ETC	1.5%	1.2%	1.9%	0.17%
DXM ETC	2.0%	1.8%	2.3%	0.11%

Past 30-day Use of Selected Substances - Border Grade 7					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	5.1%	4.4%	5.8%	0.36%	
CIGARETTES	4.8%	4.1%	5.5%	0.35%	
SMOKELESS TOBACCO	1.3%	1.0%	1.6%	0.13%	
ALCOHOL	16.7%	15.5%	18.0%	0.64%	
INHALANTS	9.7%	8.9%	10.5%	0.41%	
ANY ILLICIT DRUG	6.6%	5.4%	7.9%	0.61%	
MARIJUANA	4.4%	3.5%	5.3%	0.46%	
MARIJUANA ONLY	1.5%	1.1%	1.9%	0.18%	
COCAINE OR CRACK	1.4%	1.0%	1.8%	0.20%	
COCAINE	1.0%	0.7%	1.3%	0.16%	

CRACK	1.5%	1.1%	2.0%	0.21%
HALLUCINOGENS	1.6%	1.2%	2.0%	0.21%
UPPERS	1.7%	1.3%	2.1%	0.21%
DOWNERS	2.0%	1.3%	2.6%	0.30%
ROHYPNOL	2.5%	1.8%	3.1%	0.32%
STEROIDS	0.9%	0.7%	1.1%	0.09%
ECSTASY	3.9%	3.1%	4.7%	0.38%
HEROIN	1.4%	1.1%	1.6%	0.14%
METHAMPHETAMINE	2.4%	1.8%	2.9%	0.28%
CODEINE	5.5%	4.2%	6.7%	0.60%
OXYCONTIN ETC	0.9%	0.6%	1.1%	0.13%
VICODIN ETC	0.8%	0.6%	1.0%	0.10%
VALIUM ETC	0.8%	0.5%	1.0%	0.13%
XANAX ETC	0.9%	0.6%	1.1%	0.13%
DXM ETC	3.7%	3.2%	4.2%	0.25%

i de la companya de				•
CRACK	0.7%	0.4%	1.0%	0.13%
HALLUCINOGENS	0.7%	0.4%	1.0%	0.17%
UPPERS	1.0%	0.7%	1.2%	0.14%
DOWNERS	1.0%	0.7%	1.3%	0.15%
ROHYPNOL	0.8%	0.4%	1.2%	0.20%
STEROIDS	0.4%	0.3%	0.6%	0.08%
ECSTASY	2.0%	1.4%	2.5%	0.27%
HEROIN	0.5%	0.3%	0.7%	0.10%
METHAMPHETAMINE	1.1%	0.9%	1.3%	0.12%
CODEINE	2.7%	2.0%	3.4%	0.35%
OXYCONTIN ETC	0.4%	0.4%	0.5%	0.04%
VICODIN ETC	0.4%	0.2%	0.5%	0.08%
VALIUM ETC	0.4%	0.2%	0.5%	0.08%
XANAX ETC	0.4%	0.3%	0.5%	0.06%
DXM ETC	1.9%	1.5%	2.3%	0.20%

Lifetime Use of Selected Substances - Border Grade 8				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	26.3%	24.9%	27.8%	0.74%
CIGARETTES	25.5%	24.2%	26.9%	0.68%
SMOKELESS TOBACCO	5.6%	4.5%	6.7%	0.55%
ALCOHOL	55.2%	52.3%	58.1%	1.44%
INHALANTS	22.2%	19.8%	24.7%	1.24%
ANY ILLICIT DRUG	20.5%	18.3%	22.7%	1.09%
MARIJUANA	17.1%	15.2%	18.9%	0.92%
MARIJUANA ONLY	7.2%	6.4%	8.1%	0.41%
COCAINE OR CRACK	4.9%	3.8%	6.1%	0.57%
COCAINE	4.1%	3.1%	5.1%	0.51%
CRACK	2.3%	1.8%	2.8%	0.24%
HALLUCINOGENS	3.8%	2.9%	4.7%	0.46%
UPPERS	3.6%	2.7%	4.5%	0.45%
DOWNERS	4.2%	3.2%	5.2%	0.49%
ROHYPNOL	4.6%	3.4%	5.8%	0.62%
STEROIDS	1.2%	0.9%	1.5%	0.13%
ECSTASY	6.7%	4.7%	8.7%	1.01%
HEROIN	2.0%	1.6%	2.5%	0.22%
METHAMPHETAMINE	3.3%	2.6%	4.0%	0.36%
CODEINE	7.9%	6.7%	9.2%	0.62%
OXYCONTIN ETC	1.6%	1.3%	1.9%	0.14%
VICODIN ETC	1.5%	1.2%	1.8%	0.15%
VALIUM ETC	1.2%	0.9%	1.4%	0.13%

Past 30-day Use of Selected Substances - Border Grade 8				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	9.5%	7.7%	11.3%	0.90%
CIGARETTES	8.5%	7.0%	10.0%	0.75%
SMOKELESS TOBACCO	2.7%	2.0%	3.4%	0.34%
ALCOHOL	23.9%	22.3%	25.5%	0.79%
INHALANTS	10.1%	8.5%	11.8%	0.82%
ANY ILLICIT DRUG	10.8%	9.5%	12.2%	0.70%
MARIJUANA	7.7%	6.7%	8.8%	0.53%
MARIJUANA ONLY	2.7%	2.3%	3.2%	0.24%
COCAINE OR CRACK	2.0%	1.4%	2.5%	0.28%
COCAINE	1.7%	1.2%	2.2%	0.24%
CRACK	0.8%	0.5%	1.2%	0.17%
HALLUCINOGENS	1.4%	1.1%	1.6%	0.13%
UPPERS	1.7%	1.2%	2.2%	0.25%
DOWNERS	1.7%	1.2%	2.1%	0.24%
ROHYPNOL	2.1%	1.4%	2.8%	0.36%
STEROIDS	0.6%	0.3%	0.9%	0.14%
ECSTASY	3.4%	2.2%	4.7%	0.63%
HEROIN	1.1%	0.7%	1.5%	0.18%
METHAMPHETAMINE	1.3%	1.0%	1.6%	0.15%
CODEINE	4.2%	3.5%	4.9%	0.35%
OXYCONTIN ETC	0.7%	0.5%	0.8%	0.07%
VICODIN ETC	0.5%	0.4%	0.7%	0.07%
VALIUM ETC	0.6%	0.4%	0.8%	0.09%

XANAX ETC	2.0%	1.6%	2.3%	0.16%
DXM ETC	4.8%	3.7%	5.8%	0.54%

Lifetime Use of Selected Substances - Border Grade 9				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	31.0%	29.7%	32.4%	0.69%
CIGARETTES	30.5%	29.1%	32.0%	0.72%
SMOKELESS TOBACCO	6.8%	5.5%	8.2%	0.67%
ALCOHOL	61.7%	58.2%	65.2%	1.75%
INHALANTS	20.2%	17.6%	22.8%	1.28%
ANY ILLICIT DRUG	28.0%	25.8%	30.3%	1.11%
MARIJUANA	24.7%	22.6%	26.8%	1.06%
MARIJUANA ONLY	9.2%	8.2%	10.3%	0.53%
COCAINE OR CRACK	7.5%	6.5%	8.6%	0.52%
COCAINE	6.9%	5.8%	8.0%	0.54%
CRACK	2.4%	2.0%	2.8%	0.21%
HALLUCINOGENS	4.8%	4.0%	5.7%	0.44%
UPPERS	7.0%	5.9%	8.1%	0.55%
DOWNERS	8.4%	6.9%	9.8%	0.72%
ROHYPNOL	6.8%	5.5%	8.2%	0.67%
STEROIDS	1.3%	0.9%	1.6%	0.19%
ECSTASY	11.4%	9.6%	13.2%	0.89%
HEROIN	2.2%	1.6%	2.8%	0.30%
METHAMPHETAMINE	3.1%	2.7%	3.5%	0.20%
CODEINE	11.1%	9.6%	12.7%	0.79%
OXYCONTIN ETC	2.4%	2.1%	2.7%	0.15%
VICODIN ETC	2.4%	1.6%	3.2%	0.38%
VALIUM ETC	1.9%	1.5%	2.4%	0.22%
XANAX ETC	3.6%	3.1%	4.1%	0.25%
DXM ETC	5.7%	5.0%	6.4%	0.36%

Lifetime Use of Selected Substances - Border Grade 10					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	36.8%	33.3%	40.4%	1.78%	
CIGARETTES	35.7%	32.4%	39.0%	1.67%	
SMOKELESS TOBACCO	8.9%	7.3%	10.4%	0.80%	
ALCOHOL	69.0%	64.6%	73.5%	2.23%	
INHALANTS	18.0%	17.0%	19.0%	0.51%	
ANY ILLICIT DRUG	33.7%	29.3%	38.2%	2.22%	
MARIJUANA	30.2%	25.9%	34.5%	2.16%	

XANAX ETC	0.8%	0.5%	1.2%	0.16%
DXM ETC	2.0%	1.4%	2.6%	0.31%

Past 30-day Use of Selected Substances - Border Grade 9				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	12.4%	11.4%	13.4%	0.50%
CIGARETTES	11.9%	10.9%	12.8%	0.48%
SMOKELESS TOBACCO	2.6%	2.0%	3.2%	0.32%
ALCOHOL	30.9%	28.7%	33.2%	1.12%
INHALANTS	8.3%	7.3%	9.3%	0.51%
ANY ILLICIT DRUG	14.4%	13.0%	15.8%	0.70%
MARIJUANA	11.2%	9.8%	12.6%	0.70%
MARIJUANA ONLY	3.0%	2.4%	3.5%	0.28%
COCAINE OR CRACK	3.0%	2.5%	3.4%	0.21%
COCAINE	2.7%	2.3%	3.1%	0.21%
CRACK	0.7%	0.4%	0.9%	0.12%
HALLUCINOGENS	1.9%	1.6%	2.3%	0.16%
UPPERS	3.4%	2.8%	4.0%	0.31%
DOWNERS	4.0%	2.9%	5.0%	0.51%
ROHYPNOL	2.8%	2.3%	3.3%	0.25%
STEROIDS	0.4%	0.2%	0.5%	0.08%
ECSTASY	5.3%	4.2%	6.4%	0.55%
HEROIN	0.9%	0.6%	1.2%	0.14%
METHAMPHETAMINE	0.9%	0.5%	1.3%	0.18%
CODEINE	5.4%	4.6%	6.3%	0.41%
OXYCONTIN ETC	1.1%	0.9%	1.2%	0.07%
VICODIN ETC	1.2%	0.8%	1.6%	0.19%
VALIUM ETC	1.0%	0.8%	1.1%	0.07%
XANAX ETC	1.6%	1.3%	1.9%	0.16%
DXM ETC	2.7%	2.3%	3.1%	0.20%

Past 30-day Use of Selected Substances - Border Grade 10					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	15.6%	13.6%	17.6%	0.98%	
CIGARETTES	14.7%	12.9%	16.4%	0.86%	
SMOKELESS TOBACCO	4.3%	2.9%	5.7%	0.69%	
ALCOHOL	34.7%	31.9%	37.5%	1.42%	
INHALANTS	6.1%	5.7%	6.5%	0.20%	
ANY ILLICIT DRUG	16.4%	14.2%	18.7%	1.12%	
MARIJUANA	13.1%	11.2%	15.0%	0.93%	

MARIJUANA ONLY	13.2%	10.9%	15.4%	1.13%
COCAINE OR CRACK	7.5%	6.4%	8.7%	0.56%
COCAINE	7.3%	6.3%	8.4%	0.54%
CRACK	1.9%	1.5%	2.4%	0.21%
HALLUCINOGENS	6.0%	4.3%	7.6%	0.82%
UPPERS	7.4%	5.7%	9.1%	0.84%
DOWNERS	8.9%	6.3%	11.4%	1.27%
ROHYPNOL	6.1%	4.5%	7.6%	0.76%
STEROIDS	0.8%	0.6%	1.1%	0.14%
ECSTASY	12.4%	9.4%	15.4%	1.49%
HEROIN	1.3%	0.9%	1.7%	0.20%
METHAMPHETAMINE	2.8%	2.2%	3.5%	0.32%
CODEINE	12.9%	11.8%	14.0%	0.56%
OXYCONTIN ETC	2.2%	1.5%	2.8%	0.32%
VICODIN ETC	3.0%	2.2%	3.8%	0.39%
VALIUM ETC	1.6%	1.3%	1.9%	0.16%
XANAX ETC	4.3%	3.8%	4.8%	0.26%
DXM ETC	5.6%	4.5%	6.6%	0.52%

MARIJUANA ONLY	4.5%	3.7%	5.3%	0.39%
COCAINE OR CRACK	2.8%	2.3%	3.3%	0.26%
COCAINE	2.6%	2.1%	3.1%	0.25%
CRACK	0.7%	0.5%	1.0%	0.11%
HALLUCINOGENS	2.2%	1.2%	3.3%	0.52%
UPPERS	3.2%	2.3%	4.1%	0.46%
DOWNERS	3.8%	2.6%	4.9%	0.58%
ROHYPNOL	2.6%	1.7%	3.5%	0.43%
STEROIDS	0.3%	0.1%	0.4%	0.07%
ECSTASY	5.7%	4.3%	7.0%	0.68%
HEROIN	0.6%	0.4%	0.8%	0.10%
METHAMPHETAMINE	0.7%	0.4%	1.0%	0.15%
CODEINE	5.2%	4.6%	5.7%	0.27%
OXYCONTIN ETC	0.8%	0.6%	1.1%	0.13%
VICODIN ETC	1.0%	0.6%	1.3%	0.18%
VALIUM ETC	0.6%	0.4%	0.8%	0.10%
XANAX ETC	1.5%	1.2%	1.9%	0.18%
DXM ETC	2.4%	2.1%	2.8%	0.18%

Lifetime Use of Selected Substances - Border Grade 11				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	41.4%	39.9%	43.0%	0.78%
CIGARETTES	40.4%	38.8%	42.0%	0.80%
SMOKELESS TOBACCO	9.7%	8.1%	11.3%	0.80%
ALCOHOL	70.1%	65.8%	74.5%	2.15%
INHALANTS	16.6%	14.2%	19.1%	1.23%
ANY ILLICIT DRUG	35.7%	32.8%	38.7%	1.48%
MARIJUANA	32.7%	30.1%	35.2%	1.29%
MARIJUANA ONLY	12.7%	10.7%	14.6%	0.99%
COCAINE OR CRACK	11.5%	10.4%	12.6%	0.56%
COCAINE	11.3%	10.1%	12.4%	0.58%
CRACK	2.1%	1.6%	2.7%	0.27%
HALLUCINOGENS	6.4%	5.1%	7.6%	0.62%
UPPERS	8.9%	6.8%	11.1%	1.08%
DOWNERS	10.7%	8.1%	13.3%	1.31%
ROHYPNOL	7.4%	4.7%	10.1%	1.35%
STEROIDS	1.5%	1.3%	1.8%	0.12%
ECSTASY	14.6%	12.7%	16.5%	0.97%
HEROIN	1.1%	0.8%	1.3%	0.10%
METHAMPHETAMINE	3.1%	2.6%	3.6%	0.26%
CODEINE	12.8%	11.3%	14.2%	0.71%

Past 30-day Use of Selected Substances - Border Grade 11				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	18.6%	17.7%	19.5%	0.46%
CIGARETTES	17.9%	17.1%	18.8%	0.43%
SMOKELESS TOBACCO	3.4%	2.5%	4.3%	0.44%
ALCOHOL	35.3%	32.3%	38.3%	1.50%
INHALANTS	4.7%	3.3%	6.1%	0.69%
ANY ILLICIT DRUG	17.7%	16.1%	19.4%	0.82%
MARIJUANA	14.3%	13.0%	15.7%	0.69%
MARIJUANA ONLY	4.1%	3.4%	4.8%	0.35%
COCAINE OR CRACK	4.3%	3.6%	5.0%	0.36%
COCAINE	4.1%	3.4%	4.8%	0.34%
CRACK	0.7%	0.6%	0.9%	0.08%
HALLUCINOGENS	2.3%	1.8%	2.8%	0.25%
UPPERS	4.2%	3.0%	5.4%	0.60%
DOWNERS	4.7%	3.3%	6.2%	0.71%
ROHYPNOL	2.9%	2.0%	3.8%	0.45%
STEROIDS	0.6%	0.3%	0.9%	0.16%
ECSTASY	6.8%	5.8%	7.8%	0.50%
HEROIN	0.4%	0.2%	0.7%	0.12%
METHAMPHETAMINE	0.7%	0.6%	0.9%	0.09%
CODEINE	5.7%	5.0%	6.5%	0.37%

OXYCONTIN ETC	3.0%	2.5%	3.5%	0.23%
VICODIN ETC	4.2%	3.5%	4.9%	0.36%
VALIUM ETC	2.7%	2.1%	3.2%	0.28%
XANAX ETC	5.7%	4.0%	7.4%	0.83%
DXM ETC	4.4%	3.5%	5.4%	0.49%

Lifetime Use of Se	elected Substances	- Border G	rade 12	
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	48.7%	45.7%	51.8%	1.52%
CIGARETTES	47.9%	45.4%	50.5%	1.28%
SMOKELESS TOBACCO	11.8%	9.7%	13.8%	1.03%
ALCOHOL	73.6%	69.2%	78.1%	2.24%
INHALANTS	14.5%	13.0%	16.0%	0.74%
ANY ILLICIT DRUG	40.7%	37.2%	44.2%	1.75%
MARIJUANA	37.3%	34.2%	40.5%	1.56%
MARIJUANA ONLY	13.8%	11.6%	15.9%	1.07%
COCAINE OR CRACK	14.5%	12.6%	16.4%	0.95%
COCAINE	14.3%	12.2%	16.4%	1.05%
CRACK	2.5%	2.0%	3.0%	0.25%
HALLUCINOGENS	6.1%	5.2%	6.9%	0.43%
UPPERS	8.9%	6.2%	11.6%	1.36%
DOWNERS	11.1%	8.2%	13.9%	1.42%
ROHYPNOL	7.7%	4.9%	10.6%	1.42%
STEROIDS	1.3%	0.6%	2.0%	0.35%
ECSTASY	16.8%	13.6%	20.0%	1.59%
HEROIN	1.1%	0.8%	1.5%	0.17%
METHAMPHETAMINE	3.6%	3.0%	4.2%	0.29%
CODEINE	14.1%	11.6%	16.6%	1.25%
OXYCONTIN ETC	2.9%	2.3%	3.4%	0.26%
VICODIN ETC	5.6%	4.4%	6.8%	0.61%
VALIUM ETC	2.9%	1.9%	4.0%	0.51%
XANAX ETC	6.2%	4.2%	8.2%	1.01%
DXM ETC	4.4%	3.4%	5.4%	0.50%

OXYCONTIN ETC	0.9%	0.6%	1.3%	0.17%	
VICODIN ETC	1.7%	1.3%	2.2%	0.23%	
VALIUM ETC	0.9%	0.3%	1.5%	0.29%	
XANAX ETC	2.2%	1.1%	3.3%	0.53%	
DXM ETC	1.4%	1.2%	1.7%	0.13%	

Past 30-day Use of Selected Substances - Border Grade 12					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	25.7%	22.9%	28.4%	1.38%	
CIGARETTES	24.9%	22.3%	27.5%	1.30%	
SMOKELESS TOBACCO	4.7%	3.9%	5.4%	0.37%	
ALCOHOL	44.0%	40.5%	47.4%	1.74%	
INHALANTS	4.2%	3.0%	5.4%	0.59%	
ANY ILLICIT DRUG	20.9%	18.7%	23.0%	1.07%	
MARIJUANA	16.9%	15.2%	18.6%	0.87%	
MARIJUANA ONLY	4.8%	3.6%	6.1%	0.63%	
COCAINE OR CRACK	4.6%	3.8%	5.4%	0.40%	
COCAINE	4.4%	3.5%	5.4%	0.47%	
CRACK	0.7%	0.4%	0.9%	0.14%	
HALLUCINOGENS	1.8%	1.4%	2.2%	0.21%	
UPPERS	3.5%	2.7%	4.3%	0.40%	
DOWNERS	4.3%	3.3%	5.4%	0.54%	
ROHYPNOL	2.5%	1.5%	3.4%	0.47%	
STEROIDS	0.5%	0.1%	0.9%	0.20%	
ECSTASY	6.6%	5.3%	7.9%	0.65%	
HEROIN	0.5%	0.4%	0.6%	0.06%	
METHAMPHETAMINE	1.0%	0.7%	1.4%	0.17%	
CODEINE	5.4%	4.4%	6.4%	0.50%	
OXYCONTIN ETC	1.0%	0.7%	1.2%	0.13%	
VICODIN ETC	2.3%	1.2%	3.5%	0.57%	
VALIUM ETC	1.5%	0.6%	2.5%	0.48%	
XANAX ETC	2.9%	1.9%	3.9%	0.48%	
DXM ETC	1.8%	1.2%	2.3%	0.28%	

Lifetime Use of Sele	Lifetime Use of Selected Substances - Non-Border Overall					
CATEGORY	Prevalence	Boundaries		Standard		
		Lower	Upper	Error		
TOBACCO	29.9%	27.6%	32.3%	1.17%		
CIGARETTES	27.9%	25.7%	30.1%	1.10%		
SMOKELESS TOBACCO	10.0%	8.6%	11.3%	0.67%		
ALCOHOL	61.9%	59.3%	64.5%	1.31%		
INHALANTS	16.7%	15.7%	17.7%	0.49%		
ANY ILLICIT DRUG	27.9%	25.4%	30.4%	1.24%		
MARIJUANA	26.6%	24.0%	29.3%	1.34%		
MARIJUANA ONLY	15.0%	13.4%	16.5%	0.77%		
COCAINE OR CRACK	4.8%	4.1%	5.4%	0.34%		
COCAINE	4.3%	3.7%	5.0%	0.34%		
CRACK	1.5%	1.3%	1.7%	0.08%		
HALLUCINOGENS	4.6%	3.9%	5.4%	0.37%		
UPPERS	5.0%	4.2%	5.9%	0.44%		
DOWNERS	5.3%	4.3%	6.2%	0.47%		
ROHYPNOL	1.9%	1.5%	2.2%	0.17%		
STEROIDS	1.5%	1.3%	1.6%	0.08%		
ECSTASY	5.8%	5.4%	6.3%	0.22%		
HEROIN	1.4%	1.2%	1.6%	0.09%		
METHAMPHETAMINE	3.0%	2.7%	3.2%	0.12%		
CODEINE	12.8%	11.5%	14.0%	0.63%		
OXYCONTIN ETC	3.3%	2.8%	3.9%	0.27%		
VICODIN ETC	5.9%	4.6%	7.2%	0.67%		
VALIUM ETC	2.5%	2.1%	3.0%	0.23%		
XANAX ETC	4.8%	3.7%	5.9%	0.55%		
DXM ETC	5.5%	5.1%	6.0%	0.23%		

Lifetime Use of Selected Substances - Non-Border Grade 7					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	15.0%	13.3%	16.6%	0.84%	
CIGARETTES	13.9%	12.3%	15.5%	0.79%	
SMOKELESS TOBACCO	4.1%	3.4%	4.8%	0.35%	
ALCOHOL	41.8%	38.6%	45.1%	1.63%	
INHALANTS	18.9%	17.4%	20.3%	0.75%	
ANY ILLICIT DRUG	11.2%	9.4%	13.0%	0.88%	
MARIJUANA	9.4%	7.7%	11.0%	0.83%	
MARIJUANA ONLY	5.4%	4.3%	6.4%	0.53%	
COCAINE OR CRACK	2.4%	1.8%	2.9%	0.26%	

Past 30-day Use of Selected Substances - Non-Border Overall					
CATEGORY	Prevalence	Bound	daries	Standard	
		Lower	Upper	Error	
TOBACCO	12.1%	10.8%	13.4%	0.65%	
CIGARETTES	10.7%	9.5%	11.9%	0.59%	
SMOKELESS TOBACCO	3.8%	3.2%	4.4%	0.28%	
ALCOHOL	28.7%	26.9%	30.5%	0.92%	
INHALANTS	5.6%	5.2%	6.0%	0.22%	
ANY ILLICIT DRUG	12.9%	11.5%	14.2%	0.68%	
MARIJUANA	11.5%	10.1%	12.9%	0.70%	
MARIJUANA ONLY	5.1%	4.4%	5.8%	0.36%	
COCAINE OR CRACK	1.3%	1.2%	1.5%	0.09%	
COCAINE	1.2%	1.0%	1.4%	0.09%	
CRACK	0.4%	0.3%	0.5%	0.03%	
HALLUCINOGENS	1.5%	1.3%	1.6%	0.09%	
UPPERS	2.0%	1.7%	2.4%	0.17%	
DOWNERS	2.2%	1.8%	2.6%	0.20%	
ROHYPNOL	0.7%	0.6%	0.9%	0.08%	
STEROIDS	0.5%	0.4%	0.6%	0.04%	
ECSTASY	2.0%	1.7%	2.2%	0.13%	
HEROIN	0.5%	0.4%	0.6%	0.04%	
METHAMPHETAMINE	0.7%	0.7%	0.8%	0.04%	
CODEINE	4.9%	4.2%	5.5%	0.31%	
OXYCONTIN ETC	1.0%	0.8%	1.2%	0.09%	
VICODIN ETC	2.0%	1.5%	2.5%	0.27%	
VALIUM ETC	0.7%	0.6%	0.9%	0.07%	
XANAX ETC	1.8%	1.2%	2.3%	0.29%	
DXM ETC	1.9%	1.7%	2.1%	0.11%	

Past 30-day Use of Selected Substances - Non-Border Grade 7				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	4.7%	4.0%	5.3%	0.33%
CIGARETTES	4.2%	3.6%	4.8%	0.31%
SMOKELESS TOBACCO	1.3%	1.0%	1.7%	0.18%
ALCOHOL	14.6%	12.7%	16.5%	0.93%
INHALANTS	8.0%	7.2%	8.8%	0.40%
ANY ILLICIT DRUG	4.8%	3.8%	5.7%	0.47%
MARIJUANA	3.7%	2.9%	4.5%	0.41%
MARIJUANA ONLY	1.7%	1.3%	2.1%	0.18%
COCAINE OR CRACK	0.8%	0.6%	1.0%	0.11%

COCAINE	1.7%	1.3%	2.1%	0.20%
CRACK	1.5%	1.2%	1.9%	0.18%
HALLUCINOGENS	1.5%	1.2%	1.8%	0.14%
UPPERS	1.1%	0.8%	1.4%	0.16%
DOWNERS	1.1%	0.8%	1.3%	0.13%
ROHYPNOL	1.1%	0.8%	1.4%	0.15%
STEROIDS	1.4%	1.1%	1.6%	0.11%
ECSTASY	1.8%	1.5%	2.2%	0.18%
HEROIN	1.5%	1.1%	1.8%	0.16%
METHAMPHETAMINE	2.0%	1.7%	2.3%	0.16%
CODEINE	8.2%	7.4%	9.0%	0.41%
OXYCONTIN ETC	1.0%	0.8%	1.1%	0.09%
VICODIN ETC	1.1%	0.8%	1.4%	0.15%
VALIUM ETC	0.9%	0.8%	1.1%	0.09%
XANAX ETC	1.3%	1.1%	1.6%	0.13%
DXM ETC	4.0%	3.4%	4.5%	0.26%

		_		_
COCAINE	0.5%	0.4%	0.7%	0.08%
CRACK	0.5%	0.4%	0.6%	0.06%
HALLUCINOGENS	0.6%	0.4%	0.8%	0.10%
UPPERS	0.4%	0.3%	0.5%	0.06%
DOWNERS	0.4%	0.3%	0.5%	0.06%
ROHYPNOL	0.4%	0.2%	0.6%	0.08%
STEROIDS	0.4%	0.2%	0.5%	0.06%
ECSTASY	0.5%	0.4%	0.7%	0.07%
HEROIN	0.5%	0.4%	0.6%	0.06%
METHAMPHETAMINE	0.5%	0.4%	0.7%	0.08%
CODEINE	3.4%	2.9%	3.9%	0.26%
OXYCONTIN ETC	0.4%	0.3%	0.5%	0.06%
VICODIN ETC	0.4%	0.2%	0.6%	0.10%
VALIUM ETC	0.4%	0.2%	0.5%	0.07%
XANAX ETC	0.6%	0.5%	0.8%	0.08%
DXM ETC	1.6%	1.2%	1.9%	0.18%

Lifetime Use of Selected Substances - Non-Border Grade 8					
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	22.3%	20.0%	24.7%	1.19%	
CIGARETTES	20.6%	18.5%	22.8%	1.09%	
SMOKELESS TOBACCO	6.4%	5.2%	7.6%	0.59%	
ALCOHOL	55.0%	51.8%	58.3%	1.61%	
INHALANTS	20.8%	19.4%	22.2%	0.70%	
ANY ILLICIT DRUG	18.5%	15.9%	21.2%	1.33%	
MARIJUANA	16.3%	13.6%	19.0%	1.34%	
MARIJUANA ONLY	9.4%	7.9%	11.0%	0.78%	
COCAINE OR CRACK	2.9%	2.3%	3.6%	0.33%	
COCAINE	2.4%	1.8%	2.9%	0.28%	
CRACK	1.6%	1.3%	2.0%	0.17%	
HALLUCINOGENS	3.0%	2.3%	3.6%	0.31%	
UPPERS	2.4%	1.8%	2.9%	0.28%	
DOWNERS	2.4%	1.8%	3.0%	0.30%	
ROHYPNOL	2.1%	1.4%	2.7%	0.32%	
STEROIDS	1.6%	1.4%	1.9%	0.13%	
ECSTASY	3.9%	3.1%	4.7%	0.41%	
HEROIN	1.7%	1.3%	2.1%	0.22%	
METHAMPHETAMINE	2.4%	2.0%	2.9%	0.23%	
CODEINE	9.9%	8.6%	11.3%	0.66%	
OXYCONTIN ETC	1.8%	1.4%	2.3%	0.23%	
VICODIN ETC	2.3%	1.6%	2.9%	0.32%	

Past 30-day Use of Selected Substances - Non-Border Grade 8					
CATEGORY	Prevalence	Boundaries		Standard	
		Lower	Upper	Error	
TOBACCO	7.0%	5.9%	8.0%	0.52%	
CIGARETTES	6.0%	5.2%	6.9%	0.43%	
SMOKELESS TOBACCO	2.1%	1.7%	2.5%	0.22%	
ALCOHOL	21.2%	18.5%	23.9%	1.35%	
INHALANTS	8.0%	7.1%	8.8%	0.41%	
ANY ILLICIT DRUG	8.4%	6.8%	10.0%	0.78%	
MARIJUANA	6.6%	5.3%	7.9%	0.66%	
MARIJUANA ONLY	3.0%	2.3%	3.7%	0.33%	
COCAINE OR CRACK	1.1%	0.8%	1.4%	0.13%	
COCAINE	0.9%	0.6%	1.1%	0.11%	
CRACK	0.5%	0.4%	0.7%	0.09%	
HALLUCINOGENS	1.0%	0.6%	1.3%	0.17%	
UPPERS	1.1%	0.7%	1.5%	0.20%	
DOWNERS	1.2%	0.7%	1.7%	0.24%	
ROHYPNOL	0.9%	0.5%	1.3%	0.21%	
STEROIDS	0.4%	0.3%	0.5%	0.05%	
ECSTASY	1.4%	0.9%	1.9%	0.25%	
HEROIN	0.7%	0.4%	0.9%	0.12%	
METHAMPHETAMINE	0.8%	0.5%	1.1%	0.16%	
CODEINE	4.4%	3.8%	5.0%	0.29%	
OXYCONTIN ETC	0.5%	0.4%	0.7%	0.09%	
VICODIN ETC	0.8%	0.5%	1.0%	0.14%	

VALIUM ETC	1.3%	0.9%	1.8%	0.23%
XANAX ETC	2.3%	1.3%	3.2%	0.47%
DXM ETC	4.9%	4.0%	5.8%	0.45%

Lifetime Use of Sele	Lifetime Use of Selected Substances - Non-Border Grade 9				
CATEGORY	Prevalence	Boun	daries	Standard	
		Lower	Upper	Error	
TOBACCO	28.4%	26.0%	30.8%	1.18%	
CIGARETTES	26.5%	24.3%	28.7%	1.11%	
SMOKELESS TOBACCO	8.6%	7.3%	9.9%	0.65%	
ALCOHOL	63.1%	60.5%	65.7%	1.29%	
INHALANTS	17.1%	15.2%	19.0%	0.94%	
ANY ILLICIT DRUG	26.4%	24.1%	28.7%	1.16%	
MARIJUANA	25.3%	23.1%	27.6%	1.12%	
MARIJUANA ONLY	14.3%	12.8%	15.8%	0.75%	
COCAINE OR CRACK	4.1%	3.3%	4.8%	0.36%	
COCAINE	3.7%	3.0%	4.3%	0.35%	
CRACK	1.4%	1.1%	1.7%	0.15%	
HALLUCINOGENS	4.5%	3.6%	5.4%	0.44%	
UPPERS	4.2%	3.2%	5.1%	0.46%	
DOWNERS	4.3%	3.4%	5.2%	0.45%	
ROHYPNOL	2.0%	1.5%	2.5%	0.24%	
STEROIDS	1.3%	1.0%	1.6%	0.16%	
ECSTASY	5.5%	4.6%	6.3%	0.43%	
HEROIN	1.4%	1.0%	1.7%	0.16%	
METHAMPHETAMINE	2.9%	2.5%	3.3%	0.20%	
CODEINE	12.6%	11.5%	13.7%	0.56%	
OXYCONTIN ETC	3.0%	2.4%	3.6%	0.28%	
VICODIN ETC	4.6%	3.6%	5.6%	0.51%	
VALIUM ETC	2.1%	1.6%	2.6%	0.25%	
XANAX ETC	4.1%	2.9%	5.3%	0.61%	
DXM ETC	5.7%	4.9%	6.5%	0.40%	

Lifetime Use of Selec	Lifetime Use of Selected Substances - Non-Border Grade 10					
CATEGORY	Prevalence	Boundaries		Standard		
		Lower	Upper	Error		
TOBACCO	34.4%	31.9%	36.8%	1.22%		
CIGARETTES	32.1%	29.8%	34.4%	1.15%		
SMOKELESS TOBACCO	11.9%	10.0%	13.7%	0.92%		
ALCOHOL	69.4%	67.1%	71.7%	1.13%		
INHALANTS	16.2%	14.3%	18.0%	0.93%		
ANY ILLICIT DRUG	33.6%	30.5%	36.8%	1.58%		

VALIUM ETC	0.6%	0.3%	0.8%	0.12%
XANAX ETC	1.0%	0.5%	1.5%	0.26%
DXM ETC	2.2%	1.6%	2.7%	0.27%

Past 30-day Use of Selected Substances - Non-Border Grade 9				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	10.6%	9.2%	12.0%	0.70%
CIGARETTES	9.2%	7.9%	10.6%	0.67%
SMOKELESS TOBACCO	3.3%	2.7%	3.9%	0.30%
ALCOHOL	28.2%	26.1%	30.3%	1.05%
INHALANTS	5.8%	5.0%	6.6%	0.41%
ANY ILLICIT DRUG	12.1%	10.7%	13.5%	0.72%
MARIJUANA	11.1%	9.7%	12.5%	0.70%
MARIJUANA ONLY	4.9%	4.1%	5.8%	0.43%
COCAINE OR CRACK	1.4%	1.1%	1.7%	0.15%
COCAINE	1.3%	0.9%	1.6%	0.16%
CRACK	0.4%	0.3%	0.5%	0.05%
HALLUCINOGENS	1.7%	1.4%	2.0%	0.15%
UPPERS	1.9%	1.5%	2.2%	0.16%
DOWNERS	2.0%	1.6%	2.4%	0.20%
ROHYPNOL	0.7%	0.5%	0.9%	0.12%
STEROIDS	0.5%	0.3%	0.6%	0.08%
ECSTASY	2.0%	1.6%	2.3%	0.16%
HEROIN	0.4%	0.2%	0.5%	0.05%
METHAMPHETAMINE	0.8%	0.6%	1.0%	0.09%
CODEINE	4.9%	4.1%	5.7%	0.39%
OXYCONTIN ETC	1.1%	0.8%	1.3%	0.12%
VICODIN ETC	1.7%	1.3%	2.1%	0.21%
VALIUM ETC	0.6%	0.5%	0.8%	0.08%
XANAX ETC	1.6%	0.9%	2.3%	0.35%
DXM ETC	2.1%	1.6%	2.5%	0.20%

Past 30-day Use of Selected Substances - Non-Border Grade 10				
CATEGORY	Prevalence	Boundaries		Standard
		Lower	Upper	Error
TOBACCO	13.5%	12.2%	14.8%	0.64%
CIGARETTES	11.8%	10.6%	13.0%	0.60%
SMOKELESS TOBACCO	4.6%	3.8%	5.3%	0.38%
ALCOHOL	32.6%	31.1%	34.2%	0.78%
INHALANTS	4.3%	3.7%	4.8%	0.25%
ANY ILLICIT DRUG	16.0%	14.2%	17.8%	0.89%

MARIJUANA	32.7%	29.3%	36.0%	1.67%
MARIJUANA ONLY	18.2%	16.3%	20.1%	0.95%
COCAINE OR CRACK	6.0%	4.9%	7.2%	0.56%
COCAINE	5.6%	4.5%	6.7%	0.56%
CRACK	1.6%	1.2%	1.9%	0.16%
HALLUCINOGENS	5.5%	4.5%	6.5%	0.49%
UPPERS	6.9%	5.5%	8.4%	0.74%
DOWNERS	6.6%	5.5%	7.8%	0.59%
ROHYPNOL	2.4%	1.9%	2.8%	0.23%
STEROIDS	1.4%	1.0%	1.7%	0.17%
ECSTASY	6.9%	6.2%	7.6%	0.35%
HEROIN	1.5%	1.1%	1.8%	0.18%
METHAMPHETAMINE	3.1%	2.7%	3.5%	0.21%
CODEINE	15.1%	13.0%	17.1%	1.02%
OXYCONTIN ETC	4.4%	3.5%	5.3%	0.45%
VICODIN ETC	7.8%	5.7%	9.9%	1.06%
VALIUM ETC	3.2%	2.5%	3.8%	0.32%
XANAX ETC	6.4%	4.5%	8.4%	0.97%
DXM ETC	6.6%	5.8%	7.4%	0.39%

MARIJUANA	14.5%	12.7%	16.3%	0.89%
MARIJUANA ONLY	6.6%	5.5%	7.8%	0.56%
COCAINE OR CRACK	1.6%	1.3%	2.0%	0.17%
COCAINE	1.5%	1.2%	1.9%	0.17%
CRACK	0.3%	0.1%	0.4%	0.07%
HALLUCINOGENS	1.5%	1.0%	2.0%	0.25%
UPPERS	2.8%	2.1%	3.5%	0.35%
DOWNERS	2.9%	2.1%	3.6%	0.38%
ROHYPNOL	0.9%	0.7%	1.1%	0.11%
STEROIDS	0.4%	0.3%	0.6%	0.09%
ECSTASY	2.3%	1.6%	2.9%	0.31%
HEROIN	0.6%	0.4%	0.8%	0.09%
METHAMPHETAMINE	0.8%	0.6%	0.9%	0.08%
CODEINE	5.9%	4.4%	7.5%	0.78%
OXYCONTIN ETC	1.3%	1.0%	1.7%	0.18%
VICODIN ETC	2.8%	1.8%	3.7%	0.47%
VALIUM ETC	0.8%	0.6%	1.0%	0.11%
XANAX ETC	2.7%	1.3%	4.0%	0.67%
DXM ETC	2.1%	1.7%	2.4%	0.18%

Lifetime Use of Selected Substances - Non-Border Grade 11				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	39.3%	37.1%	41.5%	1.09%
CIGARETTES	36.4%	34.3%	38.5%	1.06%
SMOKELESS TOBACCO	14.7%	13.0%	16.4%	0.83%
ALCOHOL	70.2%	68.0%	72.3%	1.07%
INHALANTS	14.3%	12.9%	15.7%	0.70%
ANY ILLICIT DRUG	38.8%	36.6%	41.1%	1.12%
MARIJUANA	37.9%	35.1%	40.6%	1.37%
MARIJUANA ONLY	20.9%	19.1%	22.8%	0.93%
COCAINE OR CRACK	6.3%	5.4%	7.2%	0.46%
COCAINE	6.2%	5.3%	7.1%	0.47%
CRACK	1.4%	1.1%	1.8%	0.18%
HALLUCINOGENS	6.5%	5.6%	7.4%	0.44%
UPPERS	7.9%	6.9%	8.9%	0.49%
DOWNERS	8.7%	7.3%	10.0%	0.66%
ROHYPNOL	1.9%	1.4%	2.4%	0.26%
STEROIDS	1.5%	1.1%	1.9%	0.21%
ECSTASY	8.1%	7.1%	9.1%	0.48%
HEROIN	1.2%	0.9%	1.6%	0.19%
METHAMPHETAMINE	3.7%	3.1%	4.3%	0.30%

Past 30-day Use of Selected Substances - Non-Border Grade 11				
CATEGORY	Prevalence	Bound	daries	Standard
		Lower	Upper	Error
TOBACCO	17.5%	15.8%	19.3%	0.87%
CIGARETTES	15.7%	14.2%	17.2%	0.76%
SMOKELESS TOBACCO	5.8%	4.8%	6.7%	0.49%
ALCOHOL	35.9%	33.9%	38.0%	1.01%
INHALANTS	4.0%	3.4%	4.6%	0.30%
ANY ILLICIT DRUG	17.7%	16.3%	19.0%	0.68%
MARIJUANA	16.3%	14.7%	17.8%	0.77%
MARIJUANA ONLY	6.8%	5.8%	7.7%	0.49%
COCAINE OR CRACK	1.6%	1.3%	2.0%	0.16%
COCAINE	1.5%	1.2%	1.9%	0.16%
CRACK	0.4%	0.3%	0.5%	0.05%
HALLUCINOGENS	1.9%	1.6%	2.3%	0.19%
UPPERS	3.0%	2.5%	3.4%	0.22%
DOWNERS	3.5%	2.9%	4.1%	0.29%
ROHYPNOL	0.8%	0.5%	1.1%	0.15%
STEROIDS	0.5%	0.2%	0.7%	0.10%
ECSTASY	2.7%	2.2%	3.2%	0.24%
HEROIN	0.5%	0.3%	0.7%	0.09%
METHAMPHETAMINE	0.9%	0.7%	1.1%	0.11%

CODEINE	4 - 70/	14 20/	17 20/	0.750/
CODEINE	15.7%	14.2%	17.2%	0.75%
OXYCONTIN ETC	4.8%	4.2%	5.5%	0.35%
VICODIN ETC	9.9%	8.2%	11.5%	0.82%
VALIUM ETC	3.5%	2.8%	4.2%	0.35%
XANAX ETC	7.3%	6.2%	8.4%	0.56%
DXM ETC	6.0%	5.3%	6.7%	0.35%

Lifetime Use of Selected Substances - Non-Border Grade 12				
CATEGORY	Prevalence	Boun	daries	Standard
		Lower	Upper	Error
TOBACCO	44.1%	41.3%	46.8%	1.37%
CIGARETTES	41.5%	39.0%	44.0%	1.24%
SMOKELESS TOBACCO	16.0%	14.2%	17.8%	0.91%
ALCOHOL	75.4%	73.3%	77.5%	1.04%
INHALANTS	12.0%	10.4%	13.5%	0.77%
ANY ILLICIT DRUG	42.9%	40.3%	45.5%	1.30%
MARIJUANA	42.6%	39.8%	45.4%	1.40%
MARIJUANA ONLY	24.0%	21.9%	26.1%	1.05%
COCAINE OR CRACK	7.6%	6.7%	8.6%	0.48%
COCAINE	7.5%	6.5%	8.5%	0.51%
CRACK	1.4%	1.1%	1.7%	0.14%
HALLUCINOGENS	7.6%	6.4%	8.7%	0.57%
UPPERS	9.1%	7.7%	10.4%	0.68%
DOWNERS	9.9%	8.3%	11.4%	0.78%
ROHYPNOL	1.6%	1.2%	2.1%	0.23%
STEROIDS	1.8%	1.5%	2.2%	0.17%
ECSTASY	9.7%	8.4%	11.0%	0.67%
HEROIN	1.1%	0.9%	1.4%	0.13%
METHAMPHETAMINE	3.8%	3.3%	4.3%	0.25%
CODEINE	15.9%	14.3%	17.6%	0.82%
OXYCONTIN ETC	5.6%	4.8%	6.3%	0.37%
VICODIN ETC	11.1%	9.5%	12.7%	0.79%
VALIUM ETC	4.4%	3.7%	5.2%	0.40%
XANAX ETC	7.9%	6.6%	9.3%	0.67%
DXM ETC	6.0%	5.1%	7.0%	0.48%

CODEINE	5.3%	4.6%	6.1%	0.37%
OXYCONTIN ETC	1.5%	1.1%	1.8%	0.17%
VICODIN ETC	3.3%	2.5%	4.1%	0.40%
VALIUM ETC	1.0%	0.7%	1.4%	0.19%
XANAX ETC	2.7%	2.0%	3.3%	0.33%
DXM ETC	1.9%	1.6%	2.2%	0.17%

Past 30-day Use of Selected Substances - Non-Border Grade 12							
CATEGORY	Prevalence	Boundaries		Standard			
		Lower	Upper	Error			
TOBACCO	21.9%	19.8%	24.1%	1.07%			
CIGARETTES	19.8%	17.9%	21.8%	0.97%			
SMOKELESS TOBACCO	6.6%	5.6%	7.7%	0.52%			
ALCOHOL	43.2%	41.0%	45.4%	1.09%			
INHALANTS	2.9%	2.4%	3.4%	0.25%			
ANY ILLICIT DRUG	20.2%	18.4%	21.9%	0.89%			
MARIJUANA	18.8%	16.9%	20.7%	0.95%			
MARIJUANA ONLY	8.1%	7.2%	9.0%	0.47%			
COCAINE OR CRACK	1.5%	1.2%	1.9%	0.16%			
COCAINE	1.5%	1.1%	1.8%	0.17%			
CRACK	0.3%	0.1%	0.4%	0.07%			
HALLUCINOGENS	2.1%	1.7%	2.6%	0.23%			
UPPERS	3.5%	2.8%	4.1%	0.32%			
DOWNERS	3.7%	3.0%	4.4%	0.36%			
ROHYPNOL	0.6%	0.4%	0.8%	0.11%			
STEROIDS	0.7%	0.5%	0.9%	0.10%			
ECSTASY	3.2%	2.3%	4.1%	0.45%			
HEROIN	0.4%	0.3%	0.6%	0.07%			
METHAMPHETAMINE	0.7%	0.5%	0.9%	0.10%			
CODEINE	5.4%	4.5%	6.3%	0.45%			
OXYCONTIN ETC	1.5%	1.1%	1.8%	0.15%			
VICODIN ETC	3.4%	2.7%	4.0%	0.34%			
VALIUM ETC	1.1%	0.7%	1.5%	0.20%			
XANAX ETC	2.2%	1.7%	2.7%	0.25%			
DXM ETC	1.6%	1.2%	1.9%	0.17%			